Meta-Analysis of Genome-Wide Association Studies of Attention- Deficit/Hyperactivity Disorder

Benjamin M. Neale, Ph.D., Sarah E. Medland, Ph.D., Stephan Ripke, M.D.,
Philip Asherson, M.R.C.Psych., Ph.D., Barbara Franke, Ph.D., Klaus-Peter Lesch, M.D.,
Stephen V. Faraone, Ph.D., Thuy Trang Nguyen, Dipl. Math. oec.,
Helmut Schäfer, Ph.D., Peter Holmans, Ph.D., Mark Daly, Ph.D.,
Hans-Christoph Steinhausen, M.D., Ph.D., D.M.Sc., Christine Freitag, M.D., M.A.,
Andreas Reif, M.D., Tobias J. Renner, M.D., Marcel Romanos, M.D.,
Jasmin Romanos, M.D., Susanne Walitza, M.D., Andreas Warnke, M.D., Ph.D.,
Jobst Meyer, Ph.D., Haukur Palmason, Ph.D., Jan Buitelaar, M.D.,
Alejandro Arias Vasquez, Ph.D., Nanda Lambregts-Rommelse, Ph.D.,
Michael Gill, Mb B.Ch. B.A.O., M.D., M.R.C.Psych., F.T.C.D., Richard J.L. Anney, Ph.D.,
Kate Langley, Ph.D., Michael O'Donovan, F.R.C.Psych., Ph.D., Nigel Williams, Ph.D.,
Michael Owen, Ph.D., F.R.C.Psych., Anita Thapar, M.D., Lindsey Kent, M.D., Ph.D.,
Joseph Sergeant, Ph.D., Herbert Roeyers, M.D., Ph.D., Eric Mick, Sc.D.,
Joseph Biederman, M.D., Alysa Doyle, Ph.D., Susan Smalley, Ph.D.,
Sandra Loo, Ph.D., Hakon Hakonarson, M.D., Ph.D., Josephine Elia, M.D.,
Alexandre Todorov, Ph.D., Ana Miranda, M.D., Fernando Mulas, M.D., Ph.D.,
Richard P. Ebstein, Ph.D., Aribert Rothenberger, M.D., Ph.D.,
Tobias Banaschewski, M.D., Ph.D., Robert D. Oades, Ph.D.,
Edmund Sonuga-Barke, Ph.D., James McGough, M.D., Laura Nisenbaum, Ph.D.,
Frank Middleton, Ph.D., Xiaolan Hu, Ph.D., Stan Nelson, M.D., for the Psychiatric
GWAS Consortium: ADHD Subgroup

Objective: Although twin and family studies have shown attention-deficit/hyperactivity disorder (ADHD) to be highly heritable, genetic variants influencing the trait at a genomewide significant level have yet to be identified. As prior genome-wide association studies (GWAS) have not yielded significant results, we conducted a meta-analysis of existing studies to boost statistical power. **Method:** We used data from four projects: a) the Children's Hospital of Philadelphia (CHOP); b) phase I of the International Multicenter ADHD Genetics project (IMAGE); c) phase II of IMAGE (IMAGE II); and d) the Pfizer-funded study from the University of California, Los Angeles, Washington University, and Massachusetts General Hospital (PUWMA). The final sample size consisted of 2,064 trios, 896 cases, and 2,455 controls. For each study, we imputed HapMap single nucleotide polymorphisms, computed association test statistics and transformed them to z-scores, and then combined weighted z-scores in a meta-analysis. **Results:** No genome-wide significant associations were found, although an analysis of candidate genes suggests that they may be involved in the disorder. **Conclusions:** Given that ADHD is a highly heritable disorder, our negative results suggest that the effects of common ADHD risk variants must, individually, be very small or that other types of variants, e.g., rare ones, account for much of the disorder's heritability. *J. Am. Acad. Child Adolesc. Psychiatry*, 2010;49(9):884–897. **Key Words:** ADHD, meta-analysis, association, GWAS, genetics

Attention-deficit/hyperactivity disorder (ADHD) is one of the most common psychiatric phenotypes affecting children and adolescents. It is characterized by an inability to focus, high levels of impulsivity and age-inappropriate hyperactivity, and increased rates of antisocial, anxiety, mood, and substance use disorders. With estimates of prevalence ranging from 5% to 12%^{1,2} and 2% to 5% in adults,^{3,4} ADHD is a substantial public health concern.

Previous work in ADHD has established a strongly heritable component to the phenotype, with genetic factors, across 20 twin studies, explaining approximately 76% of the phenotypic variance.^{5–7} There appears to be substantial overlap between the two ADHD symptom dimensions hyperactivity/impulsivity and inattention.⁸ The heritability of ADHD is high, regardless of whether it is measured as a dimensional trait or a categorical disorder. Genome-wide linkage and hypothesis-driven candidate gene association studies have failed to unequivocally identify specific genetic variation that predisposes to ADHD, although genome-wide levels of significance were derived in a meta-analysis for variants within or close to the dopamine D4 (DRD4) and D5 receptor (DRD5) genes.⁹ Similarly, genome-wide association studies (GWAS) of ADHD in children have not yet yielded genome-wide significant association to common variation. The first genome-wide association scan of 909 trios from the International Multicenter ADHD Genetics (IMAGE) study did not find any significant associations,¹⁰ although an

analysis of ADHD-related quantitative traits in this data set revealed (trait-specific) genome-wide significance for two variants, in the CDH13 gene encoding the neural adhesion protein cadherin 13 and the GFOD1 gene encoding a protein of unknown function.¹¹ The other GWAS of childhood ADHD ^{12,13} did not reveal any genome-wide significant associations, nor did a small, pooled GWAS of 343 cases with persistent, adult ADHD and 304 controls did not provide statistically significant association findings.¹⁴ The recently reported first genome-wide study of copy number variants in approximately 400 trios from Children's Hospital of Philadelphia likewise did not result in genome-wide significant results.¹⁵

To determine whether a combination of the samples from studies in childhood ADHD would be sufficient to identify genes underlying the heritable component of ADHD, we conducted a meta-analysis of these genome-wide association scans. Although we failed to identify any genome-wide significant association, there is some evidence that true associations are present in the top end of our distribution of results.

Method

Samples

Our total data set comprises four projects: a) the Children's Hospital of Philadelphia (CHOP); b) phase I of the International Multisite ADHD Genetics Project (IMAGE); c) phase II of IMAGE (IMAGE II); and d) the Pfizer-funded study from the University of California, Los Angeles, Washington University, and Massachusetts General Hospital (PUWMA). Each of these datasets has been described elsewhere.^{10,12,13,15} For this meta-analysis, we attempted to include the same individual-level data as those included in the original publication, but amended the analysis with additional quality control at the single nucleotide polymorphism (SNP) and individual level to maintain quality control through the necessary imputation step (described below). We also restricted the analysis to one affected offspring per family (which eliminated three individuals from analysis). In addition, we imputed all missing data, so the results from the imputation are fully complete for all individuals, which slightly changed the results for each subsample from the published results.

The IMAGE trio samples were collected using a common protocol with centralized training and reliability testing of raters and centralized data management. Family members were of white European origin from countries including Belgium, Germany, Ireland, the Netherlands, Spain, Switzerland, the United Kingdom, and Israel. At the IMAGE sites, parents of children were interviewed with the Parental Account of Childhood Symptom (PACS), a semi-structured, standardized, investigator-based interview developed as an instrument to provide an objective measure of child behavior. Both parents and teachers completed the respective versions of the Conners ADHD rating scales and the Strengths and Difficulties Questionnaire. The assessment procedures have been described in more detail elsewhere.^{16–20} Probands had been referred for assessment of hyperactive, disruptive, or disorganized behavior and had been clinically diagnosed with ADHD (or hyperkinetic disorder, the most closely equivalent category in the ICD-10 nomenclature used at some of the clinics). Most probands (N = 868) met criteria for DSM-IV combined-type ADHD, and additional probands met criteria for

DSM-IV inattentive subtype (N = 13) or hyperactive subtype (N = 33) or missed one of the ADHD diagnoses by a single item on the structured interview (N = 19). We retained these latter families because, upon review of the medical record and structured interview data, the ADHD diagnosis was confirmed. Exclusion criteria were autism, epilepsy, IQ <70, brain disorders, and any genetic or medical disorder associated with externalizing behaviors that might mimic ADHD. All data were collected with informed consent of parents and with the approval of the site's institutional review board or ethical committee.

The IMAGE II ADHD samples included some samples from the original IMAGE project along with samples provided by colleagues at other sites, using similar but not identical methods. In Germany, families were recruited in order of clinical referral in the outpatient clinics in Wuerzburg, Homburg, and Trier. Families were of white German ancestry. All cases met DSM-IV criteria for ADHD. The index child was 6 years or more of age, further affected siblings were included when at least 6 years of age. All children were assessed by full semi-structured interview (Kiddie-Sads-PL-German Version or Kinder-DIPS) and parent and teacher ADHD DSM-IV–based rating scales to ensure pervasiveness of symptoms. Exclusion criteria were IQ <80, comorbid autistic disorders or somatic disorders (e.g., hyperthyroidism, epilepsy, neurological diseases, severe head trauma), primary affective disorders, Tourette syndrome, psychotic disorders or other severe primary psychiatric disorders, and birth weight <2,000 g.

At the Cardiff site, children ages 6 to 16 years of white British ancestry were assessed by interviewing parents with the Parent Child and Adolescent Psychiatric Assessment (CAPA), a semi-structured research diagnostic interview, and a telephone interview with the teacher using the Child ADHD Teacher Telephone Interview. All cases met diagnostic criteria for DSM-IV ADHD or ICD-10 hyperkinetic disorder or DSM-III-R ADHD and had IQ test scores >70. Exclusion criteria were pervasive developmental disorder, Tourette syndrome, psychosis or any neurological conditions. At the Scottish site, children ages 6 to 16 years, of white British ancestry were assessed by interviewing parents with the CAPA. To confirm pervasiveness, teachers completed the Conners Teacher Rating Scale. All cases met diagnostic criteria for DSM-IV ADHD. Children with an IQ <70, autistic spectrum disorder, head injury, known chromosomal abnormality, encephalitis, or significant medical conditions such as epilepsy were excluded. At the Dutch site, assessment data are available for 112 subjects 3 to 18 years of age with DSM-IV ADHD and related comorbidities, including mood, anxiety, oppositional defiant disorder (ODD) and conduct disorder (CD). Most of the sample was collected as part of a sibling pair genome-wide linkage study in ADHD.²¹ Subjects were assessed using the DSM-IV version of the Diagnostic Interview Schedule for Children (DISC-P) with both parents, supplemented by Conners' Questionnaires (old versions), the Childhood Behavior Checklist (CBCL), and Teacher Report Form (TRF). In addition, we obtained ADHD symptom severity scores from the parents on the Strengths and Weaknesses of ADHD symptoms and Normal behaviors (SWAN) questionnaire.

The IMAGE II control samples (2,653 population controls of European ancestry) were collected for an institutional review board–approved GWAS of schizophrenia and have been described elsewhere.²² Briefly, the

control participants were drawn from a US nationally representative survey panel (of approximately 60,000 adult individuals at any one time, with constant turnover) ascertained via random digit dialing. Participants were screened for psychosis and bipolar disorder but not ADHD. Control participants were not screened for ADHD. A blood sample was collected via a US national phlebotomy service. Control participants gave written consent for their biological materials to be used for medical research at the discretion of the National Institute of Mental Health.

The PUWMa samples were collected independently at three sites using similar but slightly different methods: Massachusetts General Hospital (MGH), Washington University, and University of California, Los Angeles (UCLA). At MGH, 309 families were recruited from clinics at. Screening and recruitment for some subjects (N = 121) occurred before publication of DSM-IV. Initial affection status for those subjects was based on DSM-III-R criteria; however lifetime DSM-IV-TR criteria was asked at follow-up, and only those subjects endorsing a lifetime DSM-IV-TR diagnosis of ADHD were enrolled. Remaining subjects were screened and assessed according to DSM-IV-TR criteria (N = 188). Psychiatric assessments were made with K-SADSE (Epidemiologic Version). Potential probands were excluded if they had major sensorimotor handicaps (deafness, blindness), psychosis/schizophrenia, autism, inadequate command of the English language, or a Full Scale IQ <80. At Washington University, 272 families were selected from a population-representative sample identified through birth records of the state of Missouri, for a genetic epidemiologic study of the prevalence and heritability of ADHD. The original sample included 812 complete male and female twin pairs and six individual twins 7 to 19 years of age at the time of interview identified from the Missouri Family Registry from 1996 to 2002. Families were invited into the study if at least one child exhibited three or more inattentive symptoms on a brief screening interview. Parents reported on their children and themselves, and the youths on themselves, using the Missouri Assessment of Genetics Interview for Children (MAGIC), a semi-structured psychiatric interview. DSM-IV diagnoses of ADHD were based on parental reports (most of the time, maternal). Families were excluded if a parent/guardian reported mental retardation or if the parent/guardian and twins could not speak English. At UCLA, 156 subjects were drawn from 540 children and adolescents 5 to 18 years of age and 519 of their parents ascertained from 370 families with ADHD-affected sibling pairs. Lifetime psychiatric diagnoses were based on semi-structured diagnostic interviews conducted by master's degree-level clinical psychologists or highly trained interviewers with extensive experience and reliability training in psychiatric diagnoses. Children and adolescents were assessed using the Schedule for Affective Disorders and Schizophrenia for School-Age Children-Present and Lifetime version (K-SADS-PL). Adult parents were assessed using the Schedule for Affective Disorders and Schizophrenia-Lifetime version (SADS-LA-IV), supplemented with the K-SADS Behavioral Disorders module for diagnosis of ADHD and disruptive behavior disorders. Direct interviews were supplemented with parent and teacher versions of the Swanson, Nolan, and Pelham, version IV (SNAP-IV) rating scale, as well as a parent-completed CBCL and Teacher Report Form. Parents also completed current ratings of self and spouse behavior with the ADHD Rating Scale IV. Subjects were excluded from participation if they were positive for any of the following: neurological disorder, head injury resulting in concussion, lifetime diagnoses of schizophrenia or autism, or estimated Full Scale IQ < 70. Subjects on stimulant medication were

asked to discontinue use for 24 hours before their visit.

The CHOP ADHD trio families were recruited from pediatric and behavioral health clinics in the Philadelphia area.¹⁵ Inclusion criteria included families of European descent with an ADHD proband (6–18 years of age). Exclusionary criteria included prematurity (<36 weeks), mental retardation, major medical and neurological disorders, pervasive developmental disorder, psychoses, and major mood disorders. Diagnostic data were collected using the K-SADS interview.

Table 1 presents the breakdown of samples that were included in the analysis. Three of these samples are proband–parent trio-based samples, with the fourth being a case-only sample. To analyze these controls, we selected genomically matched, pre-existing controls, available through the genetics data repository of the National Institute of Mental Health. The final sample size consisted of 2,064 trios, 896 cases, and 2,455 controls.

TABLE 1 Breakdown of Samples Included in the Analysis

Sample	Cases	Controls	Trios	SNPs
CHOP	—	—	423	469,283
IMAGE	—	—	909	438,784
IMAGE II	896	2,455	—	294,811
PUWMA	—	—	732	645,995
Total	896	2,455	2,064	1,206,462 ^a

Note: CHOP = Children's Hospital of Philadelphia; IMAGE = International Multicenter ADHD Genetics Project; PUWMA = Pfizer-funded study from the University of California, Los Angeles, Washington University, and Massachusetts General Hospital; SNP = single-nucleotide polymorphism.

^a*Imputes SNPs using Beagle 3.0.6.*

Imputation and Association Analysis

Each of the studies had been conducted on different platforms (Illumina 550K for CHOP, Perlegen 600K for IMAGE, Affymetrix 500K for IMAGE II, and Illumina 1M Duo for PUWMA). From these genome-wide association datasets, we imputed untyped loci across the genome using the HapMap Phase III European CEU and TSI samples as the reference panel.^{23,24} The IMAGE II sample had already undergone imputation, which we carried through to this meta-analysis. We had to impute SNPs for the trio samples before the meta-analysis. As

of this writing, and to the best of our knowledge, none of the imputation programs handle trios and supply posterior probabilities of the offspring's transmitted and non-transmitted alleles at each imputed locus. To resolve this, we phased the trios using Beagle, which outputs the most likely haplotypes for each parent split by transmitted and nontransmitted haplotypes. We then created case pseudo-control individuals consistent with the Haplotype Relative Risk (HRR) test.^{25,26} We then passed these case and pseudo-controls through Beagle^{27,28} to conduct imputation and analyzed the samples using MACH2DAT,²⁹ using a logistic regression model. The p value from this analysis was then transformed into a z-score. The p value represents the probability of observing a test statistic equally or more deviant than the observed test statistic from the analysis. By the same token, we can consider the normal distribution and identify the critical value for which the ratio of the area under the normal distribution above the critical value plus the area under the normal distribution below the negative critical value to the total area under the normal distribution is equal to the p value. Considered another way, we are aiming to identify the z-score that is as deviant as the test statistic from the regression analysis. Conditional on this transformation we can then conduct meta-analysis of z-scores, which is described by de Bakker et al.³⁰ Any SNP that was imputed with an Symbol2 of <0.6 was excluded from each individual dataset before meta-analysis.

Meta-Analysis

To conduct the meta-analysis, we first calculated the computed an association z-score corresponding to the association test statistic for each sample and then combined z's after weighting each by the inverse of the variances of the estimate of the odds ratios from each analysis. We then calculated the p value from this z-score. Formula 1 shows the analytic method:

where w_i is the weight of each individual study, as defined by the variance of the estimate from the logistic regression, w_t is the sum of these weights, and Z_i is the z-score for each study, as defined by the p value and direction.

Regional Association Plotting

We highlight the most significant regions from the meta-analysis by use of the SNP annotation and Proxy (SNAP) Search website.³¹ SNAP displays and annotates association results, such that the LD patterns of the genetic variation in the region are included along with the association results and recombination rate.

Results

Genome-Wide Results

The full distribution of results can be seen in the QQ plot in [Figure 1](#). Under the null hypothesis of no association, these points should fall along the diagonal line. The dotted line plots the 95% confidence interval. The QQ plot and the lambda statistic (1.025) show no appreciable inflation of the test statistic. The lambdas for each individual study are 1.085 for IMAGE II, 1.012 for IMAGE, 0.970 for PUWMA, and 1.047 for CHOP, which yields an expected lambda of 1.028 based on the average lambda, weighted by case size. The lambda statistic is the ratio of the observed median chi-square test statistic to the theoretical expectation of the median chi-square.³² This approach capitalizes on the observation that an increase in the variance of the distribution of the test statistic will increase the median of the chi-square. This increase in variance can be caused by population stratification, technical bias, or other confounders. The reason that stratification can yield lambda is that the null conditions that the allele frequencies in cases and controls are the same are violated, and thus the test statistic is expected to increase. This slightly lower than expected lambda from the meta-analysis is indicative of no particular correlated bias across these studies, enhancing our confidence in the quality of results. These results show that our quality control procedures removed most association signals that could be attributed to either technical artifacts or population stratification, and that there is little evidence for correlated biases across the studies, as correlated biases will yield a further inflation of lambda.

FIGURE 1 Quantile– quantile (QQ) plot of the metaanalysis of four attention-deficit/hyperactivity disorder genome-wide associations studies. Note: The QQ plot shows the distribution of expected p -values based against observer distribution. There is slight inflammation in the distribution of results, as indicated by the lambda of 1.025. The red dashed line represents the 95% confidence interval for the distribution of results.

Table 2 presents the top 50 findings of the meta-analysis. None of our results achieved genome-wide significance, defined as a p value of 5×10^{-8} .^{33,34} We present the regional association plots for our top three findings across the meta-analysis, using SNAP ³¹ in Figures 2a, 2b, and 2c. Our top hit is rs1464807 (p = 1.10E-06), which is in a gene-poor region. The SNP lies 230 kb 5' to SHFM1. The product of this gene may be involved in proteolysis and the regulation of the cell cycle. According to the NCBI Entrez Gene database (<http://www.ncbi.nlm.nih.gov/gene/>), mutations in the product of SHFM1 cause split hand/foot malformation type I. Eight SNPs in the region were among the top 50 findings in Table 2, which lends credibility to this association finding. Our second hit, rs177290098 (p = 1.68E-06), located in an intergenic region on chromosome 20, appears to be a false-positive result, as it is sufficiently poorly imputed in the three trio samples to be filtered from inclusion in the meta-analysis (Table 1) and SNPs in the region, which are in linkage disequilibrium (LD) with this SNP, do not show any association signal (Figure 2b). Given the low levels of recombination around this SNP, these surrounding SNPs are more likely to be reflective of the true association evidence in this region rather than the single large association observed. The third best hit, rs7463256, shows strong regional association, indicating that this is not likely to be a technical artifact. This association signal is close to the 5' end of the CHMP7 gene, with a number of additional SNPs in the top 50 list present located within the gene. CHMP7 is involved in the endosomal sorting pathway. Another three genes are present within 100 kb in this gene-rich area: TNFRSF10D, which protects against apoptosis; TNFRSF10A, which transduces cell death signals to start apoptosis; and LOXL2, which is involved in the biogenesis of connective tissue.

TABLE 2 Top 50 Hits from The Genome-Wide Attention-Deficit/Hyperactivity Disorder (ADHD) Meta-Analysis

Chr	BP ^a	SNP ^b	A1 ^c	A2 ^d	Weight ^e	z-Score ^f	p Value ^g	Direction ^h	Gene ⁱ
7	96200280	rs1464807	T	G	23.54	4.874	1.10E-06	+++	
20	58511043	rs17729098	T	C	2.85	4.788	1.68E-06	???	
7	96238981	rs12375086	T	C	23.33	4.704	2.56E-06	+++	
7	96215976	rs12673393	A	G	23.41	4.665	3.09E-06	+++	
8	23157565	rs7463256	T	C	36.67	-4.659	3.17E-06	---	CHMP7
8	23160140	rs2294123	T	G	36.67	4.608	4.77E-06	+++	CHMP7
8	23166448	rs11135712	T	C	35.75	4.575	4.77E-06	+++	CHMP7
8	23152539	rs13268919	A	G	35.78	4.551	5.33E-06	+++	
8	70971815	rs12680109	T	C	36.13	-4.502	6.73E-06	---	
8	94154715	rs1027730	A	G	21.6	4.497	6.89E-06	+++	
7	96215498	rs12673272	T	C	23.54	4.462	8.14E-06	+++	
20	51017390	rs10485813	A	G	20.41	4.46	8.20E-06	+++	
8	94152003	rs16915515	A	G	21.91	-4.456	8.37E-06	---	
20	51015479	rs16997358	A	T	20.41	4.436	9.17E-06	+++	
3	15321704	rs7340679	T	C	23	4.409	1.04E-05	++?	SH3BP5
20	58235038	rs7271202	A	G	39.25	-4.406	1.05E-05	---	
8	94153512	rs1384769	A	G	22.37	4.393	1.12E-05	+++	
15	85242738	rs7176964	A	T	17.95	-4.384	1.17E-05	---	AGBL1
19	54116059	rs9676447	T	C	2.48	-4.369	1.25E-05	??-	NUCB1
8	94150207	rs1472747	A	G	22	-4.368	1.25E-05	---	
15	89221977	rs1573643	T	C	21.93	4.357	1.32E-05	++?	FURIN
8	23162610	rs2280934	A	G	36.77	-4.355	1.33E-05	---	CHMP7
7	151104746	rs7809589	T	G	27.1	4.347	1.38E-05	+++	PRKAG2
7	96243316	rs17499178	T	G	17.29	4.342	1.41E-05	+++	
16	3307062	rs17611827	A	G	37.06	-4.336	1.45E-05	---	ZNF75A
7	96207549	rs17167761	A	C	23.49	4.327	1.51E-05	+++	
6	16858880	rs649266	T	C	46.63	-4.325	1.52E-05	---	ATXN1
11	70829906	rs736894	T	C	39.28	-4.317	1.58E-05	---	DHCR7
11	70836489	rs3750997	A	C	42.89	-4.312	1.62E-05	---	DHCR7
6	16857546	rs607138	A	G	46.97	-4.303	1.69E-05	---	ATXN1
3	61578705	rs36051446	T	C	47.5	-4.299	1.72E-05	+	PTPRG
14	97563832	rs8019853	A	G	13.85	-4.285	1.83E-05	---	
20	58232796	rs2865793	T	G	39.22	4.276	1.91E-05	+++	
20	58234268	rs6100796	T	C	38.79	-4.262	2.03E-05	---	
5	50381012	rs4866023	A	G	45.29	4.261	2.03E-05	+++	
11	70832468	rs12422045	A	G	42.85	4.261	2.03E-05	+++	DHCR7
11	70838711	rs7928249	A	G	42.6	-4.259	2.05E-05	---	
16	10540666	rs11074889	A	G	7.87	4.251	2.13E-05	?	EMP2
11	70831107	rs11603330	A	C	42.95	4.249	2.14E-05	+++	DHCR7
14	97566932	rs12892356	T	G	13.64	-4.243	2.21E-05	---	
11	70848651	rs12800438	A	G	42.58	4.24	2.23E-05	+++	NADSYN1
7	96262263	rs1004561	A	T	17.29	4.237	2.26E-05	+++	
8	75041641	rs16938747	C	G	8.88	-4.234	2.30E-05	---	TCEB1
15	50953701	rs2120445	A	G	22.04	-4.231	2.33E-05	?	
11	70828168	rs1790324	T	G	39.27	4.228	2.36E-05	+++	DHCR7
7	96235685	rs1449591	T	C	35.29	4.225	2.39E-05	+++	
20	58247823	rs1535028	T	C	44.04	4.216	2.48E-05	+++	
21	22305480	rs9982805	A	C	12.43	-4.206	2.60E-05	---	

Chr	BP ^a	SNP ^b	A1 ^c	A2 ^d	Weight ^e	z-Score ^f	p Value ^g	Direction ^h	Gene ⁱ
10	12630181	rs4747989	T	C	12.39	4.194	2.74E-05	++??	CAMK1D
11	70828433	rs1630498	A	C	39.15	4.191	2.77E-05	++++	DHCR7

Note: A = allele; BP = base pair; Chr = chromosome.
^aBase pair position as annotated in build36 of the human genome.
^brs Number for the single-nucleotide polymorphism (SNP).
^cReference allele for analysis.
^dSecond allele at locus.
^eRepresents sum of weights for results that are included in analysis based on inverse of variance of estimate of odds ratio.
^fz Used for analysis.
^gp Value associated with z score.
^hDirection of effects for each of the four studies that entered the meta-analysis, with + indicating increasing risk for A1, - indicating decreasing risk for A1, and ? indicating that the SNP was filtered from inclusion into the meta-analysis.
ⁱDefined as variant falling within gene or showing $r^2 > 0.5$ with functional variant in gene (nonsense, missense, or frame shift).

FIGURE 2 Three region association plots are shown here. Note: On the x-axis is the base pair position based on the human genome 18 build. On the left y axis, the $-\log_{10}(\text{P-value})$ is reported. On the right y-axis, the recombination rate in cM per Mb is shown. The points are each individual single-nucleotide polymorphism (SNP), genotyped and imputed, color-coded by r^2 to the most significant SNP in each region with dark red indicating an r^2 between 0.8 and 1 (high linkage disequilibrium [LD]), light red indicating an r^2 between 0.5 and 0.8 (moderate LD), light pink indicating an r^2 between 0.2 and 0.5 (low LD), and white indicating an r^2 between 0 and 0.2 (no LD).

Candidate Region Results

We also generated the distribution of results for a previously defined set of ADHD candidate genes selected by the IMAGE team ³⁵: ADRA1A, ADRA1B, ADRA2A, ADRA2C, ADRB2, ADRBK2, ARRB1, BDNF, CDH13, CHRNA4, COMT, CSNK1E, DBH, DDC, DRD1, DRD2, DRD3, DRD4, FADS1, FADS2, HES1, HTR1B, HTR1E, HTR2A, HTR2C, HTR3B, MAOA, MAOB, NFIL3, NR4A2, PER1, PER2, SLC18A2, SLC6A1, SLC6A2, SLC6A3, SLC6A4, SLC9A9, SNAP25, STX1A, SYT1, TPH1, and TPH2. These results from the 2752 SNPs in these candidate genes can be seen in [Figure 3](#). The marked deviation of this SNP set from the null expectation (diagonal line) suggests that some of these variants are associated with ADHD, although none achieve genome-wide significance. Finally, we present the top 50 results from the candidate gene SNPs in [Table 3](#).

FIGURE 3 Quantile–quantile (QQ) plot of candidate genes. Note: The QQ plot shows the distribution of expected p values against the observed distribution. The red dashed line represents the 95% confidence interval for the distribution of results. These p values are uncorrected.

TABLE 3 Top 50 Results from Candidate Genes

Chr	BP ^a	SNP ^b	A1 ^c	A2 ^d	Weight ^e	z-Score ^f	p Value ^g	Direction ^h	Gene ⁱ
16	81819599	rs8045365	A	G	11.03	-4.043	5.28E-05	----	CDH13
3	11028517	rs6778281	A	C	20.97	3.959	7.52E-05	++++	SLC6A1
16	81822613	rs12917991	A	C	10.97	3.829	0.0001286	++++	CDH13
16	81813774	rs8045006	A	C	15.84	3.778	0.000158	++++	CDH13
16	81280799	rs13334902	T	C	24.53	-3.279	0.001043	----	CDH13
16	81828063	rs7184058	A	G	22.73	3.258	0.001121	++++	CDH13
16	81298757	rs10514564	T	G	34.52	3.248	0.001161	++++	CDH13
16	81277337	rs8060937	T	C	23.89	-3.17	0.001526	----	CDH13
16	81273739	rs8048202	A	C	25	-3.126	0.001771	----	CDH13
16	81254805	rs12444015	T	C	29.01	-3.125	0.001775	----	CDH13
16	81384008	rs17192646	C	G	21.29	-3.12	0.001808	----	CDH13
16	81283227	rs13336758	A	G	24.32	-3.082	0.002054	----	CDH13
3	11041876	rs9822125	A	T	20.06	-3.068	0.002152	----	SLC6A1
16	81271375	rs4081995	T	C	23.59	3.066	0.002166	++++	CDH13
16	81267473	rs8054295	T	C	18.54	-3.029	0.002451	----	CDH13
8	26773734	rs17426222	T	C	31.51	3.006	0.002645	++++	ADRA1A
16	81469875	rs17676039	T	C	24.23	-2.941	0.003268	----	CDH13
16	81281593	rs12932203	A	G	24.63	2.929	0.0034	++++	CDH13
3	11018019	rs1919075	A	G	33.96	-2.899	0.003749	----	SLC6A1
7	50502889	rs11575522	T	C	9.84	-2.895	0.003795	----	FGNL1(+17.31kb) DDC
7	50498363	rs11575543	A	G	9.68	-2.866	0.004158	----	FGNL1(+12.78kb) DDC
16	81370055	rs17276565	T	G	23.12	2.864	0.004178	++++	CDH13
7	50503662	rs11575510	A	G	9.68	-2.864	0.004188	----	FGNL1(+18.08kb) DDC
16	81468190	rs7192135	A	C	23.78	2.851	0.004362	++++	CDH13
7	50501821	rs11575527	C	G	9.73	-2.836	0.004569	----	FGNL1(+16.24kb) DDC
16	81264849	rs9936363	T	C	18.08	-2.834	0.004598	----	CDH13
7	50575188	rs11575302	A	G	10.21	-2.832	0.004619	----	DDC
7	50503581	rs11575511	T	C	9.68	-2.821	0.004783	----	FGNL1(+18kb) DDC
3	11025563	rs11710097	A	G	19.22	-2.814	0.004896	----	SLC6A1
7	50505546	rs11575494	A	G	9.67	-2.797	0.005165	----	FGNL1(+19.96kb) DDC
7	50499488	rs11575535	T	C	9.72	-2.791	0.005248	----	FGNL1(+13.91kb) DDC
16	81497693	rs16958840	C	G	25.06	2.767	0.005651	++++	CDH13
7	50504792	rs10499693	A	G	9.71	-2.763	0.005729	----	FGNL1(+19.21kb) DDC
3	11025518	rs17466478	T	C	14.51	-2.756	0.00586	----	SLC6A1
7	50563965	rs11575340	T	G	9.71	-2.74	0.00613	----	DDC
16	81349970	rs1870847	T	C	29.09	-2.715	0.006625	----	CDH13
7	50504601	rs11575501	A	G	9.66	2.713	0.006677	++++	FGNL1(+19.1kb) DDC
Chr	BP ^a	SNP ^b	A1 ^c	A2 ^d	Weight ^e	z-Score ^f	p Value ^g	Direction ^h	Gene ⁱ
12	70629554	rs11179003	T	C	12.45	2.707	0.006791	++++	TFH2
16	81854762	rs16960006	T	G	19.76	2.698	0.006982	++++	CDH13
7	50499379	rs11575537	T	C	9.69	-2.696	0.007019	----	FGNL1(+13.88kb) DDC
3	11038545	rs11712912	A	G	23.37	2.693	0.007086	++++	SLC6A1
16	81495401	rs17674776	T	C	25.11	2.681	0.007334	++++	CDH13
16	81490402	rs16958834	T	C	25.15	-2.629	0.008564	----	CDH13
16	81491933	rs9646331	C	G	25.06	-2.61	0.009053	----	CDH13
3	11024133	rs1710891	T	C	35.51	2.599	0.009352	++++	SLC6A1
3	11015439	rs9990174	T	G	33.21	-2.589	0.009613	----	SLC6A1
16	81275531	rs8055389	A	G	24.72	-2.587	0.009683	----	CDH13
22	24309234	rs713921	A	G	5.95	2.555	0.01063	+++	ADRBK2
16	81483050	rs16958826	T	G	30.61	2.552	0.01071	++++	CDH13
22	24308356	rs738344	A	G	5.94	2.542	0.01101	+++	ADRBK2

^a Note: A = allele, BP = base pair, Chr = chromosome.
^b Basepair position as annotated in build30 of the human genome.
^c N = Number for the single nucleotide polymorphism (SNP).
^d Reference allele for analysis.
^e Second allele of locus.
^f Represents sum of weights for results that are included in analysis based on inverse of variance of estimate of odds ratio.
^g Used for analysis.
^h p-Value associated with z-score.
ⁱ Direction of effect for each of the four studies that entered the meta-analysis, with + indicating increasing risk for A1, - indicating decreasing risk for A1, and ? indicating that SNP was filtered from inclusion in meta-analysis.
 Defined as variant falling within 2.5 kb of a gene or showing $r^2 > 0.5$ with a functional variant in the gene (promoter, intron, or frame shift with the distance from the gene included in parenthesis).

Discussion

We have presented the first meta-analysis of genome-wide association datasets for childhood ADHD, including a total of 2,064 trios, 896 cases, and 2,455 controls. Genome-wide significant effects still elude detection for this disorder, suggesting that the effect sizes for the common variants influencing risk for ADHD are likely to be very small. However, these results include a number of promising regions, for which replication is an essential next step. Moreover, our analysis of candidate genes suggests that some of these loci are worth studying further.

The most strongly implicated region is on chromosome 7, where eight SNPs feature among the top 50 association findings. The chromosome 7 region is gene-poor. The gene most near to the findings is the SHFM1 gene, which is still more than 200 kb away from the association findings and is not an obvious candidate for ADHD. According to Entrez, the product of SHFM1 may be involved in proteolysis and the regulation of the cell

cycle. Mutations in the gene cause split hand/foot malformation, indicating that it functions from an early time point in embryonic development. SHFM1 is also expressed in the brain, which points to a possibly yet unknown function for this gene in the context of ADHD. Genetic variation in the same region was previously found to be associated with major depressive disorder and bipolar disorder at p values of 10^{-4} in GWAS.[29,36](#) In addition to the finding on chromosome 7, a region on chromosome 20 contains the second best hit in our study. It includes four other SNPs from the top 50 list. This region does not lie close to known genes, however.

A region on chromosome 8, featuring five SNPs in the top list, also seems interesting. The genes in this region play a role in immunity and cardiovascular processes, but the CHMP7 gene, implicated most stringently by our findings, also shows expression in the brain and has a function in endosomal sorting and vesicular transport.[28](#) Mutations in a related gene, CHMP2B, led to frontotemporal dementia, disinhibition, and executive dysfunction.[37](#) A second (small) region on chromosome 8 implicated through four association findings in the top 50 does not contain any genes, but lies less than 100 kb 5' to C8ORF83, a gene of unknown function.

In addition to those, a region on chromosome 11 contains 8 of the 50 top findings. This region contains the 5' ends of two genes, DHCR7 and NADSYN1. The former encodes 7-dehydrocholesterol reductase, and mutations in this gene cause Smith–Lemli–Opitz syndrome. In adults, the ubiquitously transcribed DHCR7 is most abundant in adrenal gland, liver, testis, and brain. NADSYN1 encodes nicotinamide adenine dinucleotide (NAD) synthase 1, which catalyzes the final step in the biosynthesis of NAD from nicotinic acid adenine dinucleotide (NaAD). NAD is a coenzyme in metabolic redox reactions, a precursor for several cell signaling molecules, and a substrate for protein posttranslational modifications. Replication efforts will be necessary to be certain that these putative associations are actually related to ADHD.

Clearly, this sample still lacks power to be able to unequivocally identify genome-wide significant associations of common genetic variants (SNPs) with ADHD. There are a number of possible explanations for why this lack of power still persists. First, it may be that the true effect sizes of individual genetic variants for ADHD are extremely small, accounting for less than 0.51% of the variance. As such, our findings remain in line with other psychiatric phenotypes. Recent work on psychiatric phenotypes such as schizophrenia and bipolar disorder have identified bona fide associations, but the combined sample size required to do is two to three times as large as the results presented in this study. As a result, further genome-wide association studies and replication efforts are essential. We note that at similar sample sizes for schizophrenia and bipolar disorder, there likewise were no unequivocal associations. Essentially, this meta-analysis imposes a ceiling on the effect size for common variation in the sample. We have 98% power to detect an effect accounting for 0.5% of the variance of the phenotype. The power to detect is 48% when we decrease the effect size to 0.25% of the variance and drops to 2% power when we consider an effect size of 0.1% of the phenotypic variance explained. These power calculations are all conducted assuming an alpha level of 5×10^{-8} , consistent with the genome-wide significance threshold, and are based on the postcleaning sample size.

Beyond the question of effect size, there are other possible explanations for our negative findings. Although each data collection site provided diagnoses of ADHD based on structured, systematic interviews, there may be variability in measurement across data collection sites. Such variability would induce additional phenotypic (and probably genotypic) heterogeneity. Alternatively, a similar problem would arise if differences in local referral patterns led to increased heterogeneity between the datasets. In addition, it may be that much of the genetic variation that predisposes to ADHD is extremely rare and deleterious, rather than common and conferring modest risk. A number of studies in other psychiatric disorders recently suggested that a significant (albeit undefined) proportion of cases may be caused by such rare variants, in this case copy number variants.³⁸ A first study in ADHD by members of this group also pointed to a role of such rare variants in this disorder.¹⁵ We also note that we may lose power from incomplete tagging of the putative causal variation. The expected non/centrality parameter for an SNP in LD with a functional variant is equal to the product of the r^2 and the noncentrality parameter at the causal locus.³⁹ Furthermore, a gene $\times$ environment interaction may also yield a reduction in power, as might a gene $\times$ gene interaction. Although much of our sample is drawn from populations of western European descent, there is likely a great deal of environmental variability between the countries of origin that are included in this study, in terms of diet, exposure to toxins, and so forth. This variation may interact and influence the genetic mechanisms that predispose to ADHD and, as a consequence, limit power to detect an association. In all likelihood, each of these possible explanations may partially contribute to the difficulty in finding ADHD susceptibility genes through GWAS.

This work should be viewed in the context of some limitations. Although we have conducted the largest genetic study of ADHD to date, our sample size is still relatively small compared with successful GWAS of complex disorders. Because creating this large sample required a collaborative effort among many sites, differences in ascertainment approaches and clinical assessments could have added noise to our sample and reduced power. Moreover, although using repository controls is a cost-effective strategy, because our control sample was not screened for ADHD, our power was less than it would have been for a screened sample.

In conclusion, we amassed a sizeable genome-wide association dataset for meta-analysis consisting of 2,960 individual childhood ADHD cases plus parental or independent control samples. Such numbers are indicative of a strong commitment by the ADHD genetics community to share data and to collaborate in the identification of significant associations to gain biological insight. Additional collaborative efforts should return true associations that will start to unravel the complex biological underpinnings of ADHD.

We thank the patients and the family members who provided data for this project.

REFERENCES

1. Faraone SV, Sergeant J, Gillberg C, Biederman J. The worldwide prevalence of ADHD: Is it an American Condition? *World Psychiatry*. 2003;2:104-113.
2. Polanczyk G, de Lima MS, Horta BL, Biederman J, Rohde LA. The worldwide prevalence of ADHD: A systematic review and metaregression analysis. *Am J Psychiatry*. 2007;164:942-948.
3. Kessler RC, Adler L, Barkley R, *et al*. The prevalence and correlates of adult ADHD in the United States: Results from the National Comorbidity Survey Replication. *Am J Psychiatry*. 2006; 163:716-723.
4. Faraone SV, Biederman J. What is the prevalence of adult ADHD? Results of a population screen of 966 adults. *J Attention Disord*. 2005;9:384-391.
5. Freitag CM, Rohde LA, Lempp T, Romanos M. Phenotypic and measurement influences on heritability estimates in childhood ADHD. *Eur Child Adolesc Psychiatry*. 2010;19:311-323.
6. Faraone SV, Perlis RH, Doyle AE, *et al*. Molecular genetics of attention-deficit/hyperactivity disorder. *Biol Psychiatry*. 2005;57: 1313-1323.
7. Faraone SV, Mick E. Molecular genetics of attention deficit hyperactivity disorder. *Psychiatr Clin North Am*. 2010;33:159- 180.
8. McLoughlin G, Ronald A, Kuntsi J, Asherson P, Plomin R. Genetic support for the dual nature of attention deficit hyperactivity disorder: Substantial genetic overlap between the inattentive and hyperactive-impulsive components. *J Abnorm Child Psychol*. 2007;35:999-1008.
9. Li D, Sham PC, Owen MJ, He L. Meta-analysis shows significant association between dopamine system genes and attention deficit hyperactivity disorder (ADHD). *Hum Mol Genet*. 2006;15:2276- 2284.
10. Neale BM, Lasky-Su J, Anney R, *et al*. Genome-wide association scan of attention deficit hyperactivity disorder. *Am J Med Genet B Neuropsychiatr Genet*. 2008;147B:1337-1344.
11. Lasky-Su J, Anney RJ, Neale BM, *et al*. Genome-wide association scan of the time to onset of attention deficit hyperactivity disorder. *Am J Med Genet B Neuropsychiatr Genet*. 2008;147B:1355- 1358.
12. Mick E, Todorov A, Smalley S, *et al*. Family-based genome-wide association scan of attention-deficit/hyperactivity disorder. *J Am Acad Child Adolesc Psychiatry*. 2010;49:xxx-xxx.
13. Consortium II. Case control genome-wide association of attention deficit/hyperactivity disorder. *J Am Acad Child Adolesc Psychiatry*. 2010;49:xxx-xxx.
14. Lesch KP, Timmesfeld N, Renner TJ, *et al*. Molecular genetics of adult ADHD: Converging evidence from genome-wide association and extended pedigree linkage studies. *J Neural Transm*. 2008;115:1573-1585.
15. Elia J, Gai X, Xie HM, *et al*. Rare structural variants found in attention-deficit hyperactivity disorder are preferentially associated with neurodevelopmental genes. *Mol Psychiatry*. 2010;15: 637-646.
16. Neale BM, Lasky-Su J, Anney R, *et al*. Genome-wide association scan of attention deficit hyperactivity disorder. *Am J Med Genet B Neuropsychiatr Genet*. 2008;147B:1337-1344.
17. Anney RJ, Hawi Z, Sheehan K, *et al*. Parent of origin effects in attention/deficit hyperactivity disorder (ADHD): Analysis of data from the International Multicenter ADHD Genetics (IMAGE) program. *Am J Med Genet B Neuropsychiatr Genet*. 2007;147B: 1495-1500.
18. Lasky-Su J, Neale BM, Franke B, *et al*. Genome-wide association scan of quantitative traits for attention deficit hyperactivity disorder identifies novel associations and confirms candidate gene associations. *Am J Med Genet B Neuropsychiatr Genet*. 2008; 147B:1345-1354.
19. Lasky-Su J, Anney RJ, Neale BM, *et al*. Genome-wide association scan of the time to onset of attention deficit hyperactivity disorder. *Am J Med Genet B Neuropsychiatr Genet*. 2008;147B:1355- 1358.
20. Anney RJ, Lasky-Su J, O'Dushlaine C, *et al*. Conduct disorder and ADHD: Evaluation of conduct problems as a categorical and quantitative trait in the international multicentre ADHD genetics study. *Am J Med Genet B Neuropsychiatr Genet*. 2008;147B:369- 1378.

21. Bakker SC, van der Meulen EM, Buitelaar JK, *et al.* A wholegenome scan in 164 Dutch sib pairs with attention-deficit/ hyperactivity disorder: Suggestive evidence for linkage on chromosomes 7p and 15q. *Am J Hum Genet.* 2003;72:1251-1260.
22. O'Donovan MC, Craddock N, Norton N, *et al.* Identification of loci associated with schizophrenia by genome-wide association and follow-up. *Nat Genet.* 2008;40:1053-1055.
23. Thorisson GA, Smith AV, Krishnan L, Stein LD. The International HapMap Project Web site. *Genome Res.* 2005;15:1592-1593.
24. The International HapMap Consortium. The International Hap- Map Project. *Nature.* 2003;426:789-796.
25. Knapp M, Seuchter SA, Baur MP. The haplotype-relative-risk (HRR) method for analysis of association in nuclear families. *Am J Hum Genet.* 1993;52:1085-1093.
26. Terwilliger JD, Ott J. A haplotype-based 'haplotype relative risk' approach to detecting allelic associations. *Hum Hered.* 1992;42: 337-346.
27. Browning BL, Browning SR. A unified approach to genotype imputation and haplotype-phase inference for large data sets of trios and unrelated individuals. *Am J Hum Genet.* 2009;84: 210-223.
28. Browning SR, Browning BL. Rapid and accurate haplotype phasing and missing-data inference for whole-genome association studies by use of localized haplotype clustering. *Am J Hum Genet.* 2007;81:1084-1097.
29. Moskvina V, Craddock N, Holmans P, *et al.* Gene-wide analyses of genome-wide association data sets: Evidence for multiple common risk alleles for schizophrenia and bipolar disorder and for overlap in genetic risk. *Mol Psychiatry.* 2009;14:252-260.
30. de Bakker PI, Ferreira MA, Jia X, Neale BM, Raychaudhuri S, Voight BF. Practical aspects of imputation-driven meta-analysis of genome-wide association studies. *Hum Mol Genet.* 2008;17: R122-R128.
31. Johnson AD, Handsaker RE, Pulit SL, Nizzari MM, O'Donnell CJ, de Bakker PI. SNAP: A Web-based tool for identification and annotation of proxy SNPs using HapMap. *Bioinformatics.* 2008; 24:2938-2939.
32. Devlin B, Roeder K. Genomic control for association studies. *Biometrics.* 1999;55:997-1004.
33. Pe'er I, Yelensky R, Altshuler D, Daly MJ. Estimation of the multiple testing burden for genomewide association studies of nearly all common variants. *Genet Epidemiol.* 2008;32:381-385.
34. Dudbridge F, Gusnanto A. Estimation of significance thresholds for genomewide association scans. *Genet Epidemiol.* 2008;32:227- 234.
35. Brookes K, Xu X, Chen W, *et al.* The analysis of 51 genes in DSM-IV combined type attention deficit hyperactivity disorder: Association signals in DRD4, DAT1 and 16 other genes. *Mol Psychiatry.* 2006;11:934-953.
36. Konneker T, Barnes T, Furberg H, Losh M, Bulik CM, Sullivan PF. A searchable database of genetic evidence for psychiatric disorders. *Am J Med Genet B Neuropsychiatr Genet.* 2008;147B:671- 675.
37. Urwin H, Ghazi-Noori S, Collinge J, Isaacs A. The role of CHMP2B in frontotemporal dementia. *Biochem Soc Trans.* 2009; 37:208-212.
38. Weiss LA, Shen Y, Korn JM, *et al.* Association between microdeletion and microduplication at 16p11.2 and autism. *N Engl J Med.* 2008;358:667-675.
39. Sham PC, Zhao JH, Cherny SS, Hewitt JK. Variance-components QTL linkage analysis of selected and non-normal samples: Conditioning on trait values. *Genetic Epidemiol.* 2000;19(Suppl 1): S22-S28.