

## 12. Sediment transport in the Geba River system

Amanuel Zenebe<sup>1,\*</sup>, Matthias Vanmaercke<sup>2,3</sup>, Jean Poesen<sup>2</sup>, Gert Verstraeten<sup>2</sup>, Jan Nyssen<sup>4</sup>, Gebreyohannes Girmay<sup>1,2</sup>, Mitiku Haile<sup>1</sup>, Hans Bauer<sup>1,2</sup>, Jozef Deckers<sup>2</sup>

<sup>1</sup>Mekelle University, Department of Land Resources Management and Environmental Protection, POBox 231, Mekelle, Ethiopia

<sup>2</sup>Department of Earth and Environmental Sciences, K.U.Leuven, Celestijnenlaan 200E, B-3001 Heverlee, Belgium

<sup>3</sup>Fund for Scientific Research—Flanders, Belgium

<sup>4</sup>Department of Geography, Ghent University, Krijgslaan 281(S8), B-9000 Gent, Belgium

\*Corresponding author: (amanuelzenebe@gmail.com)

### 1 Introduction

Soil erosion by water is one of the most important geomorphological processes in the Northern Ethiopian highlands. Studies on river discharge and sediment yield at the intermediate catchment scale (100 – 10 000 km<sup>2</sup>) are however hardly available in the Ethiopian highlands. Nevertheless, huge investments are made for dam construction works in this region. Due to a lack of reliable river discharge and sediment yield data many dams and reservoirs have been over-dimensioned (Haregeweyn, et al., 2006). Likewise, many reservoirs fill up with sediments at an alarming rate because of the underestimation of sediment yield. This research, therefore, aims at a better understanding the river discharge and sediment yield dynamics in the Geba River system.

### 2 Study area

This study was conducted in the catchment of Geba River, a tributary of Tekeze (Atbara) and Nile River (Fig. 1). The Geba drains an area (A) of 5133 km<sup>2</sup> in the highlands of Tigray, northern Ethiopia. The Geba catchment is located between 38°38'E and 39°48'E and between 13°18'N and 14°15'N, and is representative for the northern Ethiopian highlands regarding its bio-physical characteristics, i.e. climate, geology, soils, topography, and land use. Ten tributaries of the Geba were monitored in this study: Suluh (SU), Genfel (GE), Agula (AG), Ilala (IL), Upper Geba (UG), May Gabat (MY), Endasselassie (EN), Middle Geba (MG), Upper Tankwa (UT) and Lower Tankwa (LT). Several of these catchments are nested. SU, GE, AG and IL are tributaries of UG. UG, MY and EN are tributaries of MG. Furthermore, UT is a tributary of LT, which flows in to Geba a few hundred meters downstream of the Middle Geba gauging station (Fig. 1).

The long term average yearly rainfall in the catchment is 646 mm, strongly concentrated in July-September. During the rainy season, convective clouds are generally formed at the end of the morning, leading to rain showers in the afternoon. The Geba catchment lies between altitudes of 936 and 3314 m.a.s.l. The topography is mountainous with plateaus that are deeply incised by river gorges. The geology of the Geba catchment consists of a basement complex plateau having an upper sedimentary rock layer with some doleritic intrusions and capped by basalt trap series. Alluvium occurs along narrow incised river valleys. The catchment is extensively cultivated including steep and stony valley sides. Effective soil and water conservation measures (stone bunds) were constructed throughout the catchment since the 1970s (Munro et al., 2008).




Fig. 1. River and rainfall monitoring stations of Geba catchment (after Zenebe et al., submitted). Red dot indicates excursion viewpoint.

### 3. Methodology

River discharge and suspended sediment concentration measurements were conducted during the rainy seasons (July – September) of 2004-2007 in ten monitoring stations of Geba catchment (Fig. 1). Discharge was determined using the velocity-area method, and depth-discharge rating curves were developed to estimate the continuous discharge from automatically recorded 10-minute interval depth series using pressure transducers (Zenebe, 2009; Vanmaercke et al., 2010). Suspended sediment samples (n = 2846) were taken at the monitoring stations in the Geba catchment from 2004 to 2007 (Vanmaercke et al., 2010). Simple and multiple regression models were applied for identifying the catchment controlling factors of river discharge and area-specific sediment yield (SSY) (Vanmaercke et al., 2010; Zenebe et al., subm.).



### 4.3 Flash floods and associated sediment yield

The largest part of the runoff in all monitored catchments took place in the form of flash floods (Fig. 3), i.e. high discharge events that occur in a very short period of time. The hydrologic regime is characterized by flashfloods due to intense convective rainfall, combined with the steep topography and often poor soil cover (Vanmaercke, et al., 2010). These floods mostly occur in the evening or night and often have a steep rising limb with flow depths rising from several centimeters to 3–8 m in less than 1 h (Zenebe, 2009). With regard to sediment transport, both positive and negative hysteresis occurs in the Geba catchment. Mostly there is a positive hysteresis, which is generally attributed to sediment depletion in upper slopes of a basin, sometimes even before the runoff has peaked (Williams, 1989; Vanmaercke et al., 2010).




Fig. 3 Hydrograph of Upper Geba and its monitored tributaries during the flash flood of August 26-27 2006 (Zenebe et al., submitted).

### 4.4 SSY and catchment area

The relationship between catchment area and SSY is complex, as it is scale dependent and, furthermore, depends strongly on active erosion processes (de Vente and Poesen, 2005; Vanmaercke et al., 2010). Nyssen et al. (2004) found a decreasing significant negative

relationship between A and SSY in Ethiopia. However, this relationship included almost no observations for medium-sized catchments (100–10,000 km<sup>2</sup>), and it would underestimate the SSY values of this study. The higher than expected SSY for medium-sized catchments (Fig. 4) can probably be explained by the geomorphologic characteristics of the Ethiopian highlands (Vanmaercke et al., 2010): in catchments up to 4000 km<sup>2</sup> there is little possibility for redeposition as they are generally steep with well confined rivers, whereas larger catchments include extensive floodplains.




Fig. 4. Yearly sediment yield (SSY) in relation to catchment area (A). Available sediment yield data from other studies were also included (Nyssen et al., 2004; Haregeweyn et al., 2008). Data for Koka are probably overestimated and were not included in the regression analysis of Nyssen et al. (2004). Trend line is obtained by free hand curve fitting (Vanmaercke et al., 2010).

Cumulative distribution analyses of the continuous sediment export data indicate that roughly 50% of the sediment was exported during <5% of the measuring period (Fig. 5). This clearly illustrates the high temporal variability in sediment export (Vanmaercke et al., 2010). Hence, expensive measurement campaigns should focus on sampling flash floods.




Fig. 5. Relations between the suspended sediment yield (SSY) of the largest one to three floods and the seasonal sediment yield for all data of the ten monitored sub-catchments ( $n=21$ ) (Vanmaercke et al., 2010)

Soil and water conservation measures have a positive impact on the reduction of runoff and SSY of small catchments in the Ethiopian highlands (Haregeweyn et al., 2008; Nyssen et al., 2009). In contrast, this study showed no significant correlation between RC and estimated stone bund density (SBD) of each sub-catchment (Zenebe, 2009; Zenebe et al., subm.). Similar results were also found between SBD and SSY (Vanmaercke et al., 2010). Both cases (RC and SSY) tend to decrease with SBD although the relationship is insignificant. The lack of complete data on all types of soil and water conservation measures (e.g. soil bund and check dams are not considered) may result in less correlation of SBD with RC or SSY. Moreover, as stone bunds are constructed in all sub-catchments, this may result in less contrast among the catchments to generate a difference in RCs and SSY. Taking into consideration the high spatial variation in runoff and SSY, integrated watershed management would be crucial to reduce the runoff and sediment yield responses in the Geba sub-catchments.

## 5 Conclusions

The sub-catchments of Geba show large temporal and spatial variation in river discharge due to the variation of rainfall and other bio-physical characteristics in the catchments. Flash floods are intense with large volumes of river discharge and sediment yield causing casualties, and damage to infrastructures. Such events are difficult to predict since these floods depend mainly on local rain storms. The runoff depth is controlled by rainfall and proportion of limestone in the catchment. Runoff coefficient decreases with catchment area, as well as with the proportion of limestone in the catchment and tends to decrease with increasing density of soil conservation structures. Measured sediment yields of medium-sized catchments in the northern Ethiopian highlands are higher than estimations of previous studies. Increasing vegetation cover and strengthening on-going soil and water conservation measures have a positive effect by decreasing runoff and sediment yield. Insight of temporal and spatial variability of river

discharge and sediment yield is important to prevent casualties, damage to infrastructure and allow correct design of water storage facilities.

### **Acknowledgments**

This study was conducted in the framework of VLIR-Mekelle University's IUC-Land Project (Belgium-Ethiopia). Amanuel Zenebe received postdoc financial support from VLIR and was hosted at K.U. Leuven. The help of Hailay Hagos, Kedir Mohammed, Berhane Berhe, Kim Vanhulle, Annelies Beel, Isabelle Neyskens, Elke Soumillion and all other students and guards who helped with the collection and processing of the field data is gratefully acknowledged.

### **References:**

- de Vente, J., Poesen, J., 2005. Predicting soil erosion and sediment yield at the basin scale: scale issues and semi-quantitative models. *Earth Science Reviews*, 71: 95-125.
- Haregeweyn, N., Poesen, J., Nyssen, J., De Wit, J., Haile, M., Govers, G., Deckers, J., 2006. Reservoir sedimentation in Tigray (Northern Ethiopia): Characteristics and sediment deposition problems. *Land Degradation and Development*, 17: 211-230.
- Haregeweyn, N., Poesen, J., Nyssen, J., Govers, G., Verstraeten, G., de Vente, J., Deckers, J., Moeyersons, J., Haile, M., 2008. Sediment yield variability in northern Ethiopia: a quantitative analysis of its controlling factors. *Catena* 75:65–76
- Munro R.N., Deckers, J., Mitiku Haile, Grove, A.T., Poesen, J., Nyssen, J., 2008. Soil landscapes, land cover change and erosion features of the Central Plateau region of Tigray, Ethiopia: Photo-monitoring with an interval of 30 years. *Catena*, 75: 55–64
- Nyssen, J., Poesen, J., Moeyersons, J., Deckers, J., Haile, M., Lang, A., 2004. Human impact on the environment in the Ethiopian and Eritrean highlands—a state of the art. *Earth-Sci. Rev.*, 64: 273–320.
- Nyssen, J., Clymans, W., Descheemaeker, K., Poesen, J., Vandecasteele, I., Vanmaercke, M., Mitiku Haile, Nigussie Haregeweyn, Moeyersons, J., Martens, K., Amanuel Zenebe, Van Camp, M., Tesfamichael Gebreyohannes, Deckers, J., Walraevens, K., 2009. Impact of soil and water conservation on catchment hydrological response - a case in northern Ethiopia. *Hydrological processes* 24: 1880-1895.
- Vanmaercke, M., Zenebe, A., Poesen, J., Nyssen, J., Verstraeten, G., Deckers, J., 2010. Sediment dynamics and the role of flash floods in sediment export from medium-sized catchments: a case study from the semi-arid tropical highlands in northern Ethiopia. *Journal of Soils and Sediments*, 10: 611-627.
- Williams, G.P., 1989. Sediment concentration versus water discharge during single hydrologic events in rivers. *Journal of Hydrology*, 111: 89–106.
- Zenebe, A., 2009. Assessment of spatial and temporal variability of river discharge, sediment yield and sediment-fixed nutrient export in Geba River catchment, northern Ethiopia, PhD dissertation, Department of Geography, K.U. Leuven.
- Zenebe, A., Vanmaercke, M., Poesen, J., Nyssen, J., Verstraeten, G., Haregeweyn, N. Mitku Haile, Kassa Amare, Deckers, S (submitted). Spatial and temporal variability of river discharge in degraded semi-arid tropical catchments of the Northern Ethiopian Highlands. *Land Degradation and Development*.