

Morphine decreases 5-HT_{2A} receptor binding measured with SPECT in the canine frontal cortex.

A. Adriaens^a, S. Vermeire^a, T. Waelbers^a, L. Duchateau^a, S. Sys^a, S. Van Dorpe^b, J. Eersels^c, A. Dobbeleir^{a,d}, B. De Spiegeleer^b, K. Audenaert^d, I. Polis^a, K. Peremans^a

^aFaculty of Veterinary Medicine, Ghent University, Merelbeke, Belgium; ^bDrug Quality & Registration (DruQuaR) group, Faculty of Pharmaceutical Sciences, Ghent University, Ghent, Belgium; ^cNuclear Medicine and PET research, VU University Medical Center, Amsterdam, The Netherlands; ^dFaculty of Medicine, Ghent University, Gent, Belgium

Both the serotonergic and the opioid neurotransmitter systems play an important role in mood disorders and pain regulation. Electrophysiological and behavioral studies have previously demonstrated a physiological interaction between the serotonin-2A and μ -opioid receptors.

Aim: to investigate the influence of a single injection of morphine on cerebral serotonin-2A receptor (5-HT_{2A}) binding in dogs with ¹²³I-5I-R91150, a selective 5-HT_{2A} receptor radioligand, and SPECT.

Material and Methods: 5-HT_{2A} binding was estimated with (M) and without (control) morphine pretreatment (0.5 mg kg⁻¹ intravenously (IV), 30 minutes prior radioligand injection) in eight 5-year-old female beagles. Scans were carried out with a triple head gamma camera (Triad, Trionix) 90 minutes after ¹²³I-5I-R91150 injection (15.07 ± 2.69 MBq kg⁻¹ IV). Dogs were premedicated with dexmedetomidine and anesthesia was induced with propofol and maintained with isoflurane in oxygen. Semiquantification, with the cerebellum (a region void of 5-HT_{2A} receptors) as a reference region, was performed to calculate the 5-HT_{2A} receptor binding index (BI) in the frontal, parietal, temporal and occipital cortex and the subcortical region. Data were analyzed by mixed-model ANOVA. Significance was set at $p < 0.05$.

Results: A significantly decreased 5-HT_{2A} receptor BI was found after morphine administration in the right and left frontal cortices (resp. 1.41 ± 0.06 and 1.44 ± 0.08) compared to the blank scan (resp. 1.53 ± 0.10 and 1.55 ± 0.11) with $p = 0.012$ and 0.040 resp. No significant differences were noted for the other regions.

Conclusion: morphine administration decreases frontocortical 5-HT_{2A} receptor availability. This confirms an interaction between the serotonergic and the opioid neurotransmitter system. Whether the decreased radioligand binding is the consequence of decreased receptor density due to downregulation/internalization or the result of indirect actions, such as increased release of endogenous serotonin, remains to be elucidated.

This study was supported by the Ghent University Special Research Fund (grant n°01J06109).