Conditioned fear modulates visual selection

Manon Mulckhuyse¹, Geert Crombez¹ and Stefan Van der Stigchel²

**1. Department of Experimental-Clinical and Health Psychology, Ghent University,
Ghent, Belgium**

**2. Experimental Psychology, Helmholtz Institute Utrecht University, Utrecht, The
Netherlands**

Address for correspondence:

Manon Mulckhuyse

Department of Experimental-Clinical and Health Psychology

Ghent University

Henri Dunantlaan 2

B-9000 Ghent

Belgium

Email: Manon.Mulckhuyse@UGent.be

fax: +32 9 264 64 89

Abstract

Eye movements reflect the dynamic interplay between top-down and bottom-up driven processes. For example, when we voluntarily move our eyes across the visual field, salient visual stimuli in the environment may capture our attention, our eyes, or modulate the trajectory of an eye movement. Previous research showed that the behavioral relevance of a salient stimulus modulates these processes. The current study investigates whether a stimulus signaling an aversive event modulates saccadic behavior. Using a differential fear conditioning procedure, we presented a threatening (Conditioned Stimulus: CS+) and a non-threatening stimulus distractor (CS-) during an oculomotor selection task. The results show that short-latency saccades deviated stronger towards the CS+ than to the CS- distractor, whereas long-latency saccades deviated stronger away from the CS+ than from the CS- distractor. Moreover, the CS+ distractor captured the eyes more often than the CS- distractor. Together these results demonstrate that conditioned fear has a direct and immediate influence on visual selection. The findings are interpreted in terms of a neurobiological model of emotional visual processing.

Keywords: FEAR-CONDITIONING, SACCADIC CURVATURE, EMOTION, VISUAL SELECTION, EYE-MOVEMENT

Introduction

Imagine walking in a crowded underground trying to find your way out. Your eyes search for the green exit signs while you attempt to ignore distracting information. Nevertheless, your eyes are sometimes captured by irrelevant information such as flickering neon lights or screaming billboards. Such oculomotor captures are due to the high visual salience of a stimulus in relation to its background (Itti & Koch, 2000), either in terms of color, luminance, motion or orientation (Nothdurft, 2000). Attentional or oculomotor capture may occur irrespective of the task at hand (Theeuwes, 1992; 1994, but see Folk et al., 1992; 1994), but top-down processes, such as task or behavioral relevance enhance this so-called priority processing (Fecteau & Munoz, 2006). Likewise, emotional stimuli are also assumed to be prioritized by the visual system (see for review Mather & Sutherland, 2011). In line with this idea, research has shown that threatening stimuli such as an angry face in a crowd (Hansen & Hansen, 1988; Öhman et al., 2001b, but see Becker et al., 2011), a snake in the grass (Öhman et al., 2001a, but see Lipp et al., 2004; Tipples et al., 2002), or a man holding a knife (Blanchet, 2006) are detected faster than non-emotional stimuli (see for review Yiend, 2010). However, the questions how and when in time emotional stimuli modulate visual selection are still a matter of debate (Pessoa & Adolphs 2009; de Gelder et al., 2011; Pourtois et al., 2012).

One method to index the effect of emotion on attentional and visual selection is to investigate eye movement behavior. Despite the strong link between visual spatial attention and eye movements (Rizzolatti et al., 1987), not many studies on emotional attention have used eye movements. When available, they have mainly focused upon saccade latency (e.g., Bannerman et al., 2009; 2010a ; Kissler & Keil, 2008) or viewing patterns, such as first fixations or fixation duration in free-viewing or visual search

paradigms (e.g., Calvo & Lang, 2004; Humprey et al., 2012; Nummenmaa et al., 2006). These studies have found faster saccades towards and more fixations on emotional relative to non-emotional pictures. However, the emotional information in these studies is never completely task-irrelevant, leaving open the question whether emotional visual stimuli are automatically prioritized by the visual system (Hodsoll et al., 2011). For example, in the study of Calvo and Lang (2004) the emotional stimulus was also a target stimulus, and was therefore task-relevant. In the study of Bannerman et al. (2010b) the emotional stimulus is presented at the same location as the target stimulus and was therefore also task-relevant to some extent.

A well-validated eye movement paradigm to investigate the influence of task irrelevant information on visual selection is the oculomotor selection task (Van der Stigchel et al., 2006; Van der Stigchel, 2010). In this task, participants are instructed to make a simple saccade from a fixation cross to a target stimulus while a visual distractor is presented close to the target. Research has consistently revealed that salient distractors modulate the trajectory of the saccade to the target stimulus (Doyle & walker, 2001; Godijn & Theeuwes, 2004; Ludwig & Gilchrist 2003; Van der Stigchel et al., 2006). Such modulation is believed to occur because spatial attention is allocated to the distractor location (Sheliga, Riggio & Rizzolatti, 1994; Van der Stigchel & Theeuwes, 2005). Noteworthy, this modulation shows a linear relation with saccade latency: saccades that are initiated shortly after stimulus onset tend to deviate towards a distractor, whereas saccades initiated later (after 200ms) tend to deviate away from the distractor (McSorley et al., 2006). These phenomena have been explained by competing activity between the target and distractor locations in the oculomotor system. Presaccadic activity of neurons signaling the target location is averaged with activity of

neurons signaling the distractor location. Therefore, a saccade will deviate towards the distractor when a saccade is generated early after stimuli onset (McPeck, 2006; McPeck et al., 2003). Subsequently, a top-down process suppresses the activity at the distractor location (McSorley et al., 2006). Due to this inhibition, the saccade vector deviates away from the distractor location and hence, saccades generated later after stimuli onset will deviate away from a distractor (Van der Stigchel et al., 2006; Walker et al., 2006; Walker and McSorley, 2008). Erroneous saccades that land on the distractor stimulus (oculomotor capture) are explained by faster bottom-up driven activity at the distractor location relative to the slower top-down driven activity at the target location (Godijn & Theeuwes, 2002). Two studies (Nummenmaa et al., 2009, Exp. 3; Schmidt et al., 2012) have used the oculomotor selection task with an emotional distractor. Both reported that the trajectory of the saccades deviated more strongly away from an emotional distractor than from a neutral distractor. As the mean saccade latencies in the Nummenmaa et al. (2009) and Schmidt et al. (2012) studies were above 200ms, the to be expected result is indeed a deviation away from the distractor.

The main objective of our study was to investigate the time course of emotional modulation on saccade trajectories. More specifically, we were interested whether saccades that are initiated shortly after stimulus onset (short-latency saccades) result in stronger deviation towards the emotional-laden distractor than the neutral distractor, or even result in oculomotor capture, and whether long-latency saccades result in stronger deviation away from the emotional-laden distractor. To generate short-latency and long-latency saccades we varied the gap between the offset of the fixation cross and the onset of the target stimulus (McSorley et al., 2006; Munoz et al., 2000; Saslow, 1967). Previous research has revealed that saccade latencies decrease when the time between

fixation offset and target onset becomes longer, and saccade latencies increase when the fixation cross and target presentation overlap in time. To create a threatening distractor, we used a differential fear-conditioning procedure (Mackintosh, 1983). This procedure has proven successful in emotional attention research. For example, behavioral studies have shown that fear-conditioned stimuli are detected faster than neutral stimuli (Öhman et al., 2001) and are associated with enhanced activation of a fronto-parietal network that is also involved in spatial attention (Armony & Dolan, 2002). Moreover, fear-conditioned stimuli are associated with enhanced processing in visual cortex (Lim et al., 2009; Stolarova, Keil & Moratti, 2005) and with enhanced amygdala activation (Davis & Whalen, 2001; Lim et al., 2009). The amygdala is an important subcortical brain structure in emotion processing and is believed to bias perception in favor of emotional stimuli via direct connections to the visual cortex (Vuilleumier, 2005; Vuilleumier, Armony & Dolan, 2003). For example, Stolarova et al. (2005) showed that the earliest visual component (C1), which is thought to originate from striate cortex, is modulated by fear-conditioned stimuli (see also Pourtois, 2004). Moreover, in binocular rivalry fear-conditioned stimuli predominate neutral stimuli, a process that is believed to occur through direct connections from the amygdala to the visual cortex (Alpers, Ruhleder, Walz, Mühlberger, & Pauli, 2005). It is of interest that the latter two studies (Alpers et al., 2005; Stolarova et al., 2005) used conditioned stimuli that were initially neutral (e.g. gratings), suggesting that there is no need to use biologically relevant stimuli as conditioned stimuli to observe effects of fear on attention procedure. Also in our conditioning procedure, we use initially neutral stimuli: One of two colored circles (conditioned stimulus: CS+) was associated with an aversive noise (Unconditioned Stimulus: US), whereas the other colored circle (CS-) was not.

The use of a fear conditioning paradigm has two advantages in comparison with previous research. First, emotional distractors were signals of a genuine threat instead of symbolic representations of threat (e.g. pictures). Second, the stimulus features of the distractor are under experimenter control allowing for a comparison that is not influenced by potential visual differences between stimuli (e.g., Calvo et al., 2008; Hunt et al., 2007). As saccade deviation reflects the amount of competition within the oculomotor system (Godijn & Theeuwes, 2004), we expected that the threatening distractor (CS+) would be a stronger competitor for visual selection than the non-threatening distractor (CS-), and, hence, we expected that saccades generated early in time (short-latency saccades) would deviate stronger towards the threatening distractor than the non-threatening distractor and saccades generated later in time (long-latency saccades) would deviate stronger away from the threatening distractor than the non-threatening distractor. In addition, we expected the threatening distractor to capture the eyes more often than the non-threatening distractor.

Method Experiment 1

Participants, apparatus and design

Eleven participants were seated 65 cm from a computer screen with their head positioned on a chin rest. Eye movements were registered by means of a video-based eye tracker (SR Research Ltd., Canada), which has a 1000 Hz temporal and a 0.025° spatial resolution. Participants fixated a center fixation cross on a black background (see Figure 1a). After 1000 ms, the target, a light gray filled circle with a diameter of 1.44° appeared either 9.3° above or below fixation point. The fixation cross was removed at different stimulus onset asynchronies (SOA's) of: -150, -50, 0, 50, 150 ms relative to

target onset to manipulate saccade latency. In 80% of the trials a distractor, either a red or a green filled circle (1.44° in diameter) was presented simultaneously at 8.2° from the target. The vertical distance of the distractor from fixation was 5.39° , whereas it was 3.91° from the target. The horizontal distance of the distractor from fixation and from the target was 7.18° . After 2000 ms, the stimuli were removed from the screen and the next trial started. Figure 1a shows the sequence of a trial with a distractor, and fixation offset at target onset.

Procedure

Before the experiment, a differential fear conditioning procedure was started in which the distractors were presented one by one. One of the colors (Conditioned Stimulus: CS+) was paired with an aversive noise (Unconditioned Stimulus: US) whereas the other color (CS-) was not. Five participants were presented with a red CS+ and a green CS- and six participants vice versa. The US consisted of a 200 ms 100 dB white noise for half of the participants and (due to technical problems) a 50 ms 100 dB white noise for the other halfⁱ. The US was presented 250 ms after CS+ onset. When participants were able to correctly identify the color that predicted the noise, the experiment started.

In the experiment, participants were required to make a speeded saccade to the target while ignoring the presence of a distractor. Each participant performed 200 experimental trials. Furthermore, 8 additional reinforcement trials in which the CS+ was presented together with the US were randomly mixed with the experimental trials to avoid extinction (Mackintosh, 1974). The timing of the US was the same as in the fear-conditioning procedure, which was 250 ms after CS+ onset. The reinforcement trials were not further analyzed. Afterwards, ratings of fear of the CS+ and CS- and ratings of

the intensity, unpleasantness and expectancy of the US were obtained using Likert scales (1 = not at all to 9 = extremely).

Data analyses

The initial saccade starting position had to be within 1-degree horizontal and 2 degrees vertical from the center fixation point. The initial saccade was assigned to a target if the endpoint of the initial saccade was within 4° of the center of the target position and to a distractor if the endpoint was within 4° of the center of the distractor position. Too small saccades ($<3^\circ$) were considered as errors and excluded from analyses. Saccade latencies shorter than 80 ms or longer than 600 ms were also considered as errors and excluded from analyses. Latencies shorter or longer than 2.5 standard deviations away from the mean latency were considered as outliers and excluded from analyses.

Saccade trajectories were examined by calculating the mean angle of the actual saccade path relative to the mean angle of a straight line between the starting point of the saccade and the saccadic target. The angle of the actual saccade was calculated for each 2 ms sample point by examining the angle of the straight line between fixation and the current sample point. Angles were averaged across the whole saccade and subtracted from the angle of the straight line between fixation and the target location. To compute the influence of the distractor on saccade trajectories, we compared each saccade in a trial with a distractor to that of the averaged mean-path-angles of all trials without a distractor, to determine whether the saccade in the presence of a distractor deviated towards or away from the location of the distractor. Deviations were signed so that a positive value indicated deviation towards the distractor and a negative value

deviation away. All deviations are given in radians (for a review of the various measures used in the literature, see Van der Stigchel et al. 2006).

To examine the time-course of oculomotor competition, we calculated individual cumulative distribution functions of the saccade latency for each participant and each condition (CS+ or CS-). These were averaged in four bins using the vincentizing procedure (Ratcliff, 1979). Therefore, the first bin contained the fastest saccades whereas the last bin contained the slowest saccades. For each participant and condition, each bin contained a minimum of 12 trials and a maximum of 20 trials, depending on the number of correct trials for each participant and condition. When the number of trials was not even, the last bins could contain one trial more than the others.

Results

Excluded trials

Two participants were removed from analyses because of bad calibration or too many excluded trials based on the described definitions in the data-analyses (>35% excluded trials). Of the remaining nine participants, 19.3% of all trials were removed. There was no significant difference between percentage of excluded trials in the threatening condition (18.5%) compared to the non-threatening condition (20%; $p = .44$).

Fear-conditioning

Mean ratings of intensity, unpleasantness and fear of the US were 7.7, 7.1 and 5.8 respectively. Participants rated fear of the CS+ ($M = 5.1$) higher than of the CS- ($M = 1.6$; two-related Wilcoxon test, $p < 0.01$). Furthermore, expectancy rating of the US after

a CS+ ($M = 4.8$) was higher than after a CS- ($M = 1.9$; two-related Wilcoxon test, $p < 0.01$). These results suggest that our fear conditioning procedure was successful.

Eye movement behavior

Figure 2 shows the mean saccade latency and deviation per condition per bin. An ANOVA on saccade latency with bin (1, 2, 3, and 4) and signal (CS+ or CS-) as factors revealed a main effect of bin ($F(3,24) = 85.2$, $p < 0.01$, with Greenhouse–Geisser correction), but no main effect of signal ($F < 1$) and no interaction between signal and bin ($p > 0.3$).

A similar ANOVA on saccade deviation revealed no main effect of bin ($p = 0.1$), no interaction between signal and bin ($p > 0.2$), but a significant main effect of signal ($F(1,8) = 9.02$, $p < 0.05$). As can be seen in Figure 2, the saccades deviated more strongly towards the CS+ distractor than to the CS- distractor.

Furthermore, significantly more saccades landed on the CS+ distractor (9.3%) than on the CS- distractor (3.3%; two-related Wilcoxon test, $p < 0.05$).

Discussion Experiment 1

Our results clearly indicate a modulation of visual selection induced by a threatening distractor. The saccade deviation towards the threatening distractor (CS+) was stronger relative to the non-threatening distractor (CS-). Moreover, the threatening distractor captured the eyes more often than the non-threatening distractor. Together, these results suggest that threatening information has a direct low-level impact on the oculomotor system modulating visual selection early in time.

In contrast to previous studies that found deviation away from a threatening distractor (Nummenmaa et al., 2009; Schmidt et al., 2012), we found deviations towards the distractor stimuli. As discussed in the introduction, previous research showed a linear relation between trajectory deviation and saccade latency; a deviation towards a distractor is followed by a deviation away after approximately 200 ms (McSorley, 2006). Indeed, the mean latencies in the Nummenmaa et al. (2009) and Schmidt et al. (2012) studies were above 200ms. Our results show that the slowest saccades (bin 4) curved less towards the distractor than the fastest saccades (bin 1), although the interaction was not significant. Moreover, the mean saccade latencies in all bins in Experiment 1 were relatively short. Probably, this was due to the highly salient signal at the top or the bottom of the visual field of the two sudden onsets signaling target and distractor location. Sudden onsets (luminance increments) are known to be highly salient (Yantis & Jonides, 1984). Therefore, in Experiment 2 we wanted to increase saccade latency by decreasing the visual salience at the target location. We therefore expected to find slower saccade latencies and hence to find deviations away from the distractor in the last bin. Similar to Experiment 1, we expected the threatening distractor to be a stronger competitor for visual selection than the non-threatening distractor, and therefore evoke stronger deviations and capture the eyes more often than the non-threatening distractor. Furthermore, we wanted to replicate the results from Experiment 1 with a US that was the same for all participants.

Method Experiment 2

Participants, apparatus and design

Twenty participants were seated 65 cm from a computer screen with their head positioned on a chin rest. Participants fixated a center fixation cross on a black background. At a distance of 9.3° above and below the fixation point a placeholder, a light gray filled circle with a diameter of 1.44° , was presented (see Figure 1b). After 1000 ms, one of the placeholders changed to white indicating the target location. The fixation cross was removed at different stimulus onset asynchronies (SOA's): -150, -50, 0, 50, 150 ms relative to target onset. In 80% of the trials a distractor, either a red or a green filled circle (1.44° in diameter) was presented simultaneously at 8.2° from the target. The vertical distance of the distractor from fixation was 5.39° , whereas it was 3.91° from the target. The horizontal distance of the distractor from fixation and from the target was 7.18° . After 2000 ms, the stimuli were removed from the screen and the next trial started. Figure 1b shows the sequence of a trial with a distractor, and fixation offset at target onset.

Procedure and data-analyses

The same procedure and data-analyses as in Experiment 1 were used, except that the US consisted of a 200ms 100 dB white noise for all participants.

Results

Excluded trials

Three participants were removed from analyses because of bad calibration or too many excluded trials based on the described definitions (>35% excluded trials). Of the remaining seventeen participants, 22.9% of all trials were removed. A two-related

Wilcoxon test showed that more trials were excluded in the threatening condition (24.8%) compared to the non-threatening condition (21%; $p < .05$).

Fear-conditioning

Mean ratings of intensity, unpleasantness and fear of the US were 6.8, 6.9 and 6.1 respectively. Participants rated fear of the CS+ ($M = 4.9$) higher than of the CS- ($M = 1.2$; two-related Wilcoxon test, $p < 0.01$). Furthermore, expectancy rating of the US after a CS+ ($M = 4.9$) was higher than after a CS- ($M = 1.3$; two-related Wilcoxon test, $p < 0.01$). These results indicate that our fear conditioning procedure was successful.

Eye movement behavior

Figure 3 shows the mean saccade latency and deviation per condition per bin. An ANOVA on saccade latency with bin (1, 2, 3, and 4) and signal (CS+ and CS-) as factors revealed a main effect of bin ($F(3,48) = 288.86$, $p < 0.01$, with Greenhouse–Geisser correction), but no main effect of signal ($p > 0.29$) and no interaction between signal and bin ($F < 1$).

A similar ANOVA on saccade deviation revealed a main effect of bin ($F(3,48) = 7.29$, $p < 0.01$, with Greenhouse-Geisser correction), no main effect of signal ($F < 1$), but a significant interaction between signal and bin ($F(3,48) = 2.9$, $p < 0.05$, with Greenhouse-Geisser correction). A trend analysis with signal (CS+ and CS-) and bin (1, 2, 3 and 4) as factors revealed no main effect of signal ($F < 1$), a significant main effect of bin ($F(1, 16) = 25.3$, $p < .01$) and a significant interaction between signal and bin ($F(1, 16) = 7.52$, $p < .05$). As can be seen in Figure 3, the linear trend is steeper for the threatening distractor condition ($F(1,16) = 19.9$, $p < .01$) than for the non-threatening

distractor condition ($F(1,16) = 3.7, p = .07$), indicating that the threatening distractor has a stronger effect over time on saccade deviation than the non-threatening distractor.

Subsequent planned comparisons showed a marginally significant difference in the first bin ($t(16) = 1.7, p = .06$, one-tailed) due to stronger deviation towards a threatening distractor relative to a non-threatening distractor and a significant difference in the last bin ($t(16) = 2.1, p < .05$, one-tailed) due to stronger deviations away from the threatening distractor relative to a non-threatening distractor.

The two related Wilcoxon test on percentage of oculomotor capture between the non-threatening (3.4%) and the threatening distractor (4.1%) did not reach significance.

Discussion Experiment 2

The results of Experiment 2 show a modulation of visual selection induced by a threatening distractor: short-latency saccades deviated stronger towards the threatening distractor (CS+), whereas long-latency saccades deviated away stronger from the threatening distractor relative to the non-threatening distractor (CS-). The stronger deviations towards the threatening distractor (CS+) for the short-latency saccades are consistent with the results of Experiment 1, indicating that the effects do not seem to be dependent upon particular US characteristics. Furthermore, the lack of an effect of threat on saccade latency is also consistent with Experiment 1. However, as predicted, saccade latencies in Experiment 2 were increased relative to Experiment 1. Probably this was due to the less salient stimulus signaling the target location; a color change instead of a sudden onset. Consistent with previous studies on saccadic behavior with a mean latency of more than 200 ms (Nummenmaa et al., 2009; Schmidt et al., 2012), we

found emotional modulation of the saccade deviation away from the distractor for the slowest saccades.

Discussion

The present study investigated the time-course of emotional modulation on saccade trajectories. Either a threatening or a non-threatening distractor was presented close to a neutral target to induce oculomotor competition. It was predicted that the threatening distractor (CS+) would be a stronger competitor for visual selection than the non-threatening distractor (CS-) and therefore would result in stronger deviations. Consistent with our prediction, results from Experiment 1 and 2 together show that threatening information modulates short latency as well as long-latency saccades. Short-latency saccades deviated more strongly towards a threatening distractor, whereas long-latency saccades deviated more strongly away from a threatening distractor relative to a non-threatening distractor. In addition, in Experiment 1 the threatening distractor captured the eyes more often than the non-threatening distractor. This pattern of results is consistent with earlier studies that manipulated the salience of the distractor by making it more or less relevant (Hickey & van Zoest, 2012; Mulckhuyse et al., 2009). The reason why we did not find a significant difference in oculomotor capture in Experiment 2 was probably due to the increased latencies of the saccades. Previous research showed that oculomotor capture occurs more often when saccades are initiated shortly after the distractor is presented (Godijn & Theeuwes, 2002; Mulckhuyse et al., 2008).

In both experiments, saccade latency was not affected by threatening information, suggesting that the stronger deviations in the threatening condition cannot be explained by differences in saccade latency. The absence of an effect on saccade latency is not

consistent with a previous study investigating emotional modulation of saccade trajectory and latency (West et al., 2011). In contrast, West et al. found emotional modulation of saccade latency but not on saccade trajectory. Consequently, they concluded that temporal rather than spatial programming in the oculomotor system is affected by threatening information. However, in their study the threatening stimulus, an angry face, was presented at fixation, and both the target and the distractor consisted of neutral stimuli. Therefore, the different stimulus configuration is probably the cause for the discrepant findings. Future research with the use of threatening stimuli in different stimulus configurations may further elucidate the temporal and spatial aspects of saccade programming in the presence of threat.

The results from Experiment 1 revealed that a task-irrelevant threatening distractor has an immediate and direct influence on visual selection. This indicates that very early in time the oculomotor system differentiates between a threatening and a non-threatening stimulus. In contrast to the more complex emotional stimuli used in the studies by Nummenmaa et al. (2006) and Schmidt et al. (2012) our stimuli did not request high-level processing. Visual information is processed from lower to higher cortical areas during the so-called feed-forward sweep (Lamme & Roelfsema, 2000). Moreover, visual information is processed via several pathways including a subcortical pathway. The subcortical pathway is supposed to subserve automatic detection of threat (LeDoux, 2000; Liddell et al., 2005; Morris et al., 1999) and is thought to be involved in affective blindsight (Jolij & Lamme, 2005; de Gelder et al., 1999). From an evolutionary viewpoint, it is believed that this pathway processes biological relevant stimuli, such as angry faces or spiders (Öhman et al., 2001a) but it has also been implicated in fear conditioning (Hamm et al., 2003; LeDoux, 2000). For instance, in a study by Hamm et al.

(2003) a patient with bilateral cortical damage to the visual cortex was able to acquire a fear-response to a fear-conditioned visual stimulus that he could not consciously perceive. The subcortical pathway includes structures such as the superior colliculus and the pulvinar, essential for oculomotor behavior (see for review Schall, 1995) and the amygdala, which is known to be vital in the acquisition and expression of conditioned fear (Davidson & Irwin, 1999; Davis, 1992a; 1992b). Activity at the distractor location in the oculomotor system, which caused the eyes to deviate stronger towards the threatening stimulus, may thus be enhanced directly due to the reciprocal connection between the superior colliculus, pulvinar and amygdala (Linke, 1999; see also West et al., 2011). However, the superior colliculus is not the only sub-cortical structure that is essential in oculomotor behavior. For instance, the pulvinar has been associated with saccadic behavior (Robinson & McClurkin, 1989) and in addition with the prioritization processing of emotional visual stimuli (Pessoa & Adolphs, 2009). Furthermore, higher visual areas, such as the frontal eye fields (FEFs) also play an essential role in oculomotor programming (Schall, 1995) and in saccade deviations away from a distractor (McSorley et al., 2006). Moreover, the FEFs are believed to be involved in the activation of possible target locations (McPeck, 2006; Walker, Techawachirakul & Haggard, 2009) and are part of the so-called saliency map (Munoz 2002; Thompson & Bichot, 2005) or priority map (Fecteau & Munoz, 2006). Fecteau and Munoz (2006) argued that top-down information, such as behavioral relevant information integrates with bottom-up saliency information onto a priority map. The underlying mechanisms of the priority map may be the oculomotor network including areas such as the superior colliculus, the pulvinar and the FEF. The current study shows that conditioned fear has a direct and immediate influence on visual selection processes, reflected by stronger

deviations towards and away from a threatening distractor. However, how emotional stimuli modulate these visual selection processes is still unclear. More research is needed to elucidate the underlying neural mechanisms in visual processing of emotional stimuli.

References

- Alpers, G. W., Ruhleder, M., Walz, N., Muhlberger, A., & Pauli, P. (2005). Binocular rivalry between emotional and neutral stimuli: A validation using fear conditioning and EEG. *International Journal of Psychophysiology*, 57, 25–32.
- Armony, J. L., & Dolan, D. J. (2002). Modulation of spatial attention by fear-conditioned stimuli: an event-related fMRI study. *Neuropsychologia*, 40, 817–826.
- Bannerman, R. L., Milders, M., & Sahraie, A. (2010). Attentional bias to brief threat-related faces revealed by saccadic eye movements. *Emotion*, 10, 733–738.
- Bannerman, R. L., Milders, M., de Gelder, B., & Sahraie, A. (2009). Orienting to threat: Faster localization of fearful facial expressions and body postures revealed by saccadic eye movements. *Proceedings of the Royal Society B: Biological Sciences*, 276, 1635–1641.
- Becker, D. V., Anderson, U. S., Mortensen, C. R., Neufeld, S. L., & Neel, R. (2011). The face in the crowd effect unconfounded: Happy faces, not angry faces, are more efficiently detected in single- and multiple-target visual search tasks. *Journal of Experimental Psychology: General*, 140 (4), 637
- Blanchette, I. (2006). Snakes, spiders, guns, and syringes: How specific are evolutionary constraints on the detection of threatening stimuli? *The Quarterly Journal of Experimental Psychology*, 59, 1484–1504.
- Calvo, M. G., and Lang, P. J. (2004). Gaze patterns when looking at emotional pictures: motivationally biased attention. *Motivation and Emotion* 28, 221–243.
- Calvo, M. G., & Nummenmaa L. (2008). Detection of emotional faces: Salient physical features guide effective visual search. *Journal of Experimental Psychology-General*, 137(3), 471–494.
- Davidson, R. J. and W. Irwin (1999). The functional neuroanatomy of emotion and affective style. *Trends in Cognitive Sciences*, 3(1), 11-21.
- Davis, M. (1992a). The role of the amygdala in fear and anxiety. *Annual Review of Neuroscience*, 15, 353-375.
- Davis M. (1992b). The Role of the Amygdala in Conditioned Fear. In: The amygdala, ed. J. P. Aggleton. Wiley-Liss, pp. 255–306
- Davis, M. and P. J. Whalen (2001). The amygdala: vigilance and emotion. *Molecular Psychiatry*, 6(1), 13-34.

Eastwood, J. D., D. Smilek, et al. (2001). Differential attentional guidance by unattended faces expressing positive and negative emotion. *Perception & Psychophysics*, 63(6), 1004-1013.

Fecteau, J. H. & Munoz, D. P. (2006) Saliency, relevance, and firing: A priority map for target selection. *Trends in Cognitive Sciences* 10, 382–90.

Findlay JM, & Walker R. (1999). A model of saccade generation based on parallel processing and competitive inhibition. *Behavioral and Brain Sciences*, 22, 661-721.

Folk CL, Remington RW, & Johnston JC (1992) Involuntary covert orienting is contingent on attentional control settings. *Journal of Experimental Psychology: Human Perception and Performance* 18(4),1030–1044.

Folk CL, Remington RW, & Wright JH (1994) The structure of attention control: contingent attentional capture by apparent motion, abrupt onset, and color. *Journal of Experimental Psychology: Human Perception and Performance* (2), 317–329

de Gelder B, Vroomen J, Pourtois G, Weiskrantz L (1999). Non-conscious recognition of affect in the absence of striate cortex. *Neuroreport*, 10, 3759-3763.

de Gelder, B., van Honk, J. & Tamietto, M. (2011). Emotion in the brain: of low roads, high roads and roads less travelled. *Nature Reviews Neuroscience* 11, 773-783.

Godijn R, Theeuwes J (2002) Programming of endogenous and exogenous saccades: evidence for a competitive integration model. *Journal of Experimental Psychology: Human Perception and Performance*, 28(5),1039-1054.

Godijn, R., & Theeuwes, J. (2004). The relationship between inhibition of return and saccade trajectory deviations. *Journal of Experimental Psychology: Human Perception and Performance*, 30, 538-554.

Hansen, C. H., & Hansen, R. D. (1988). Finding the face in the crowd: An anger superiority effect. *Journal of Personality and Social Psychology*, 54, 917-924.

Hamm, A. O., Weike, A.I., Schupp, H.T., Treig, T, Dressel, A., & Kessler, C. (2003). Affective blindsight: intact fear conditioning to a visual cue in a cortically blind patient. *Brain*, 126, 267-275

Hickey, C., & van Zoest, W. (2012). Reward creates oculomotor saliency. *Current Biology*, 22 (7) R219–R220.

Hodsoll S, Viding E, Lavie N (2011) Attentional capture by irrelevant emotional distractor faces. *Emotion* 11, 346–353.

Humphrey, K., Underwood, G., Lambert, T. (2012). Saliency of the lambs: A

test of the saliency map hypothesis with pictures of emotive objects. *Journal of Vision*, 12 (1).

Hunt, A. R., R. M. Cooper, et al. (2007). The effect of emotional faces on eye movements and attention. *Visual Cognition*, 15(5), 513-531.

Itti L, Koch C. (2000). A saliency-based search mechanism for overt and covert shifts of visual attention. *Vision Research* 40, 1489–1506.

Jolij, J., & Lamme, V. A. F. (2005). Repression of unconscious information by conscious processing: Evidence from affective blindsight induced by transcranial magnetic stimulation. *Proceedings of the National Academy of Sciences of the USA*, 102, 10747–10751.

Kissler, J., and Keil, A. (2008). Look-don't look! How emotional pictures affect pro- and anti saccades. *Experimental Brain Research*, 188, 215-222.

Lamme, V. A. F., & Roelfsema, P. R. (2000). The distinct modes of vision offered by feedforward and recurrent processing. *Trends in Neurosciences*, 23(11), 571_579.

Lim, S., L., Padmala, S., and Pessoa, L. (2009). Segregating the significant from the mundane on a moment-to-moment basis via direct and indirect amygdala contributions. *Proceedings of the National Academy of Sciences, U.S.A.*, 106, 16841–16846.

Linke R, De Lima AD, Schwegler H, & Pape HC (1998) Direct synaptic connections of axons from the superior colliculus with identified thalamo-amygdaloid projection neurons in the rat: possible substrates of a subcortical visual pathway to the amygdala. *The Journal of Comparative Neurology* 403, 158–170

LeDoux, J. E. (2000). Emotion circuits in the brain. *Annual Review of Neuroscience*, 23, 155-184.

LeDoux, J. (2003). The emotional brain, fear, and the amygdala. *Cellular and Molecular Neurobiology*, 23(4-5), 727-738.

Liddell, B. J., K. J. Brown, et al. (2005). A direct brainstem-amygdala-cortical 'alarm' system for subliminal signals of fear. *Neuroimage* ,24(1), 235-243.

Lipp, O., Derakshan, N., Waters, A.M., & Logies, S. (2004). Snakes and cats in the flower bed: Fast detection is not specific to pictures of fear-relevant animals. *Emotion*, 4, 233–250.

Ludwig, C., J., H., & Gilchrist, I., D. (2003). Target similarity affects saccade curvature away from irrelevant onsets. *Experimental Brain Research*, 152, 60-69.

Mackintosh, N. J. (1983). Conditioning and associative learning. Oxford: University Press.

Mackintosh, N. J. (1979). *The psychology of animal learning*. New-York, NY: Academic Press.

Mather, M., & Sutherland, M. R. (2011). Arousal-biased competition in perception and memory. *Perspectives on Psychological Science*, 6, 114–133.

McPeck RM. (2006). Incomplete suppression of distractor-related activity in the frontal eye field results in curved saccades. *Journal of Neurophysiology* 96, 2699–2711.

McPeck RM, Han JH, Keller EL. (2003). Competition between saccade goals in the superior colliculus produces saccade curvature. *Journal of Neurophysiology*, 89, 2577–2590.

McSorley E, Haggard P, Walker R. (2006). Time-course of oculomotor inhibition revealed by saccade trajectory modulation. *Journal of Neurophysiology* 96, 1420–1424.

Morris, J. S., A. Ohman, et al. (1999). A subcortical pathway to the right amygdala mediating unseen fear. *Proceedings of the National Academy of Sciences of the United States of America*, 96(4), 1680-1685.

Mulckhuysen, M., Van Zoest, W., & Theeuwes, J. (2008). Capture of the eyes by relevant and irrelevant onsets. *Experimental Brain Research*, 186, 225-235.

Mulckhuysen, M., Van der Stigchel, & Theeuwes, J. (2009). Early and late modulation of saccade deviations by target distractor similarity. *Journal of Neurophysiology* 102(3), 1451-1458

Munoz DP, Dorris MC, Pare M, Everling S (2000) On your mark, get set: brainstem circuitry underlying saccadic initiation. *Canadian Journal of Physiology and Pharmacology*, 78(11), 934–944

Munoz, D., P. (2002). Commentary: saccadic eye movements: overview of neural circuitry. *Progress in Brain Research*, 140 (2002), 89-96.

Nothdurft H (2000) Salience from feature contrast: additivity across dimensions. *Vision Research* 40, 1183–1201.

Nummenmaa, L., Hyona, J., & Calvo, M. G. (2006). Eye movement assessment of selective attentional capture by emotional pictures. *Emotion*, 6, 257 – 268.

Nummenmaa, L., J. Hyona, et al. (2009). Emotional Scene Content Drives the Saccade Generation System Reflexively. *Journal of Experimental Psychology: Human Perception and Performance*, 35(2), 305-323.

Öhman, A., A. Flykt, et al. (2001a). Emotion drives attention: Detecting the snake in the grass. *Journal of Experimental Psychology: General*, 130(3), 466-478.

- Öhman, A., Lundqvist, D., & Esteves, F. (2001b). The face in the crowd revisited: An anger superiority effect with schematic stimuli. *Journal of Personality and Social Psychology*, 80, 381-396.
- Pessoa, L. & Adolphs, R. (2010) Emotion processing and the amygdala: from a 'low road' to 'many roads' of evaluating biological significance. *Nature Review Neuroscience*, 11, 773–783.
- Pourtois, G., Grandjean, D., Sander, D., & Vuilleumier, P. (2004). Electrophysiological correlates of rapid spatial orienting towards fearful faces. *Cerebral Cortex*, 14(6), 619–633.
- Pourtois G., Schettino A., Vuilleumier P. (in press). Brain mechanisms for emotional influences on perception and attention: what is magic and what is not. *Biological Psychology*
- Ratcliff, R. (1979). Group reaction-time distributions and an analysis of distribution statistics. *Psychological Bulletin* 86(3), 446–461.
- Rizzolatti, G., Riggio, L., Dascola, I., & Umiltà, C. (1987). Reorienting attention across the horizontal and vertical meridians: Evidence in favor of a premotor theory of attention. *Neuropsychologia*, 25(1A), 31-40.
- Robinson, D., L., & McClurkin, J., W. (1989). The visual superior colliculus and Pulvinar. *Reviews of oculomotor research*, Vol. 3, 337–360.
- Saslow, M.,G. (1967). Effects of components of displacement-step stimuli upon latency for saccadic eye movement. *Journal of the Optical Society of America A* 57, 1024–1029.
- Schall JD (1995) Neural basis of saccade target selection. *Reviews in the Neurosciences*, 6(1), 63–85
- Schmidt, L.J., Belopolsky, A.V. & Theeuwes, J. (in press), The presence of threat affects saccade trajectories. *Visual Cognition*
- Stolarova M, Keil A, & Moratti S (2006). Modulation of the C1 visual event-related component by conditioned stimuli: Evidence for sensory plasticity in early affective perception. *Cerebral Cortex*:16, 876–887.
- Theeuwes, J. (1992). Perceptual selectivity for color and form. *Perception & Psychophysics* 51(6), 599–606
- Theeuwes J (1994b) Endogenous and exogenous control of visual selection. *Perception* 23(4), 429–440
- Tipples, J., Young, A.W., Quinlan, P., Broks, P., & Ellis, A.W. (2002). Searching for

threat. *Quarterly Journal of Experimental Psychology: Human Experimental Psychology*, 55A, 1007–1026.

Van der Stigchel, S. (2010). Recent advances in the study of saccade trajectory deviations, *Vision Research* 50(17), 1619-1627.

Van der Stigchel S, Meeter M, Theeuwes J. (2006). Eye movement trajectories and what they tell us. *Neuroscience & Biobehavioral Review* 30, 666–679.

Vuilleumier, P. (2005). How brains beware: Neural systems for emotional attention. *Trends in Cognitive Sciences*, 9(12), 585–594,

Walker R, McSorley E. (2008). The influence of distractors on saccade target selection: saccade trajectory effects. *Journal of Eye Movement Research* 2, 1–13.

Walker, R., McSorley, E., Haggard, P. (2006). The control of saccade trajectories: direction of curvature depends upon prior knowledge of target location and saccade latency. *Perception and Psychophysics* 68 (1), 129-138.

Walker R, Techawachirakul P, Haggard P (2009). Frontal eye field stimulation modulates the balance of salience between target and distractors. *Brain Research* 1270, 54–63.

West, G. L., N. Al-Aidroos, et al. (2011). Emotion and action: the effect of fear on saccadic performance. *Experimental Brain Research*, 209(1), 153-158.

Acknowledgments: This research was funded by a Rubicon grant from NWO

(Netherlands Organization for Scientific Research) to M.M.

Figure 1a and b. From bottom to top, succession of events in a trial in which the target is presented above fixation point (on the vertical meridian) while a colored circle distractor (either a CS+ or CS-) is presented. On the left (panel a) Experiment 1 and on the right (panel b) Experiment 2.

Figure 2. Mean saccade deviation and mean saccade latency for the CS+ and CS- distractor per bin. Error bars represent normalized standard errors (Loftus & Masson, 1994).

Figure 3. Mean saccade deviation and mean saccade latency for the CS+ and CS- distractor per bin. Error bars represent normalized standard errors.

ⁱA repeated measures ANOVA on saccade deviation with US (200ms or 50ms) as between subjects factor and bin (1, 2, 3 or 4) and signal (CS+, CS-) as within subjects factor revealed no interaction with saccade deviation ($p = .16$) or saccade latency ($p = .7$) Therefore, data from both groups were collapsed.