

# Molecular Characterization of *Salmonella* Enteritidis: Comparison of an Optimized Multi-Locus Variable-Number of Tandem Repeat Analysis (MLVA) and Pulsed-Field Gel Electrophoresis

Isabelle Dewaele,<sup>1</sup> Geertrui Rasschaert,<sup>1</sup> Sophie Bertrand,<sup>2</sup> Christa Wildemaue,<sup>2</sup> Pierre Wattiau,<sup>3</sup> Hein Imberechts,<sup>3</sup> Lieve Herman,<sup>1</sup> Richard Ducatelle,<sup>4</sup> Koen De Reu,<sup>1</sup> and Marc Heyndrickx<sup>1,4</sup>

## Abstract

*Salmonella* Enteritidis (SE) is a genetically homogenous serovar, which makes optimal subtype discrimination crucial for epidemiological research. This study describes the development and evaluation of an optimized multiple-locus variable number tandem-repeat assay (MLVA) for characterization of SE. The typeability and discriminatory power of this MLVA was determined on a selected collection of 60 SE isolates and compared with pulsed-field gel electrophoresis (PFGE) using restriction enzymes *Xba*I, *Not*I, or *Sfi*I. In addition, the estimated Wallace coefficient (*W*) was calculated to assess the congruence of the typing methods. Selection of epidemiologically unrelated isolates and more related isolates (originating from layer farms) was also based on the given phage type (PT). When targeting six loci, MLVA generated 16 profiles, while PFGE produced 10, 9, and 16 pulsotypes using *Xba*I, *Not*I, and *Sfi*I, respectively, for the entire strain collection. For the epidemiologically unrelated isolates, MLVA had the highest discriminatory power and showed good discrimination between isolates from different layer farms and among isolates from the same layer farm. MLVA performed together with PT showed higher discriminatory power compared to PFGE using one restriction enzyme together with PT. Results showed that combining PT with the optimized MLVA presented here provides a rapid typing tool with good discriminatory power for characterizing SE isolates of various origins and isolates originating from the same layer farm.

## Introduction

**S**ALMONELLA ENTERITIDIS (SE) is a major cause of food-borne illness in humans, in part because of its relation to eggs (EFSA, 2010; Gantois *et al.*, 2009). Epidemiological studies have been performed to study the relatedness of isolates from human infections to contaminated eggs. Typing is a powerful tool to investigate outbreaks and to study the sources and transmission routes in the human and veterinary context (Lapuz *et al.*, 2007; Much *et al.*, 2009). However, availability of large number of genotypic and phenotypic methods (Foley *et al.*, 2007; Kang *et al.*, 2009) complicates selection of the most appropriate technique for characterizing SE. Because SE is one of the most genetically homogenous serotypes of *Salmonella* (Saeed *et al.*, 2006), methods with high discriminatory power are needed. Traditionally, SE isolates have been characterized by phage typing (PT), a universally

applied phenotypic method (De Lappe *et al.*, 2009; Pang *et al.*, 2005). The major advantage of PT is that it is a globally accepted method and specific phage type numbers can be assigned to isolates, which makes comparison between isolates possible on a worldwide scale. In addition, PT has good intra-laboratory reproducibility (Majtanova *et al.*, 2011). However, some strains are non-typeable, and possible phage type conversion (Brown *et al.*, 1999; Chart *et al.*, 1989; Tankouo-Sandjong *et al.*, 2012; Threlfall *et al.*, 1989) can occur within the serotype. Some phage types can also predominate in a geographical area, which can limit the utility of PT for investigating local outbreaks (Lukinmaa *et al.*, 1999). Another disadvantage is that only a limited number of reference laboratories perform PT (Cho *et al.*, 2008; Majtanova *et al.*, 2011). Pulsed-field gel electrophoresis (PFGE) using *Xba*I is another standard method for genotyping SE (Lacsoncha *et al.*, 2000; Rivoal *et al.*, 2009). The advantages of PFGE are its relatively

<sup>1</sup>Technology and Food Science Unit, Institute for Agricultural and Fisheries Research, Melle, Belgium.

<sup>2</sup>National Reference Centre for *Salmonella* and *Shigella*, Bacterial Diseases Division, Scientific Institute of Public Health, Brussels, Belgium.

<sup>3</sup>Veterinary and Agrochemical Research Centre, Brussels, Belgium.

<sup>4</sup>Department of Pathology, Bacteriology, and Poultry Diseases, Faculty of Veterinary Medicine, Merelbeke, Ghent University, Belgium.

good discriminatory power and good reproducibility. The PFGE method is labor-intensive and time-consuming (Foxman *et al.*, 2005), which makes it less suitable for typing a large number of isolates. More recently, multilocus variable number of tandem repeat analysis (MLVA) involving amplification and fragment size analysis of the number of repeats in the variable number tandem repeat (VNTR) regions has been documented (Van Belkum, 2007). Good reproducibility, good discriminatory power, and the ease of performance and interpretation make MLVA a valuable technique (Kruy *et al.*, 2011).

The aim of the present study was to compare an optimized MLVA with PFGE for typing SE isolates of various origins and different isolation years as well as typing isolates originating from the same layer farm within the same timeframe. If MLVA has comparable discriminatory power to PFGE, this user-friendly technique could replace the elaborate PFGE method performed together with PT. To make this comparison, we first optimized MLVA using a selection of primers from three existing MLVA systems described in literature. We then selected 60 SE isolates previously characterized by PT and used them to compare the optimized MLVA technique and PFGE using restriction enzymes *Xba*I, *Not*I, and *Sfi*I. Typeability and discriminatory power were determined for each method separately, and the Wallace coefficient combining the different methods was calculated.

## Materials and Methods

### Development and optimization of MLVA

Eight characterized SE strains (strains 1–8) of different origins, year of isolation, and/or phage type, and four characterized related outbreak strains (strains 9–12) (Table 1, panel 1) were used to evaluate the typeability and discriminatory power of 25 previously described SE MLVA primer sets (Beranek *et al.*, 2009; Boxrud *et al.*, 2007; Cho *et al.*, 2007, 2008; Malorny *et al.*, 2008; Ross and Heuzenroeder, 2009).

Strains were grown overnight on tryptone soy agar (TSA) plates (Oxoid, Basingstoke, UK) at 37°C. A small loopful of cells were resuspended in 200  $\mu$ L of high-pressure liquid chromatography (HPLC) water. After incubation during 17 min at 90°C, lysates were stored at –20°C until further use. Lysates were centrifuged for 2 min at 14,000  $\times$  g before use in polymerase chain reaction (PCR).

Primers defined by Beranek *et al.* (2009), Boxrud *et al.* (2007) Cho *et al.* (2007, 2008), Malorny *et al.* (2008), and Ross and Heuzenroeder (2009) were tested separately using the described corresponding PCR protocol to evaluate the typeability and discriminatory power of each primer pair. PCR products were analyzed by electrophoresis in 1.5% Seakem LE agarose (Lonza, Rockland, ME) with 0.5  $\times$  Tris-acetate-EDTA (TAE) for 240 min at 120 V using a 100-bp DNA size standard (Invitrogen, Carlsbad, CA).

Primer pairs generating none or multiple amplicons for each of the 12 SE isolates of panel 1 were excluded. The final MLVA included six primer pairs. Each pair discriminated among the nine strains of different origin, year, phase type, or all three (as expected, the outbreak isolates showed no difference in band size). Each pair also generated only one specific amplicon. One primer in each pair was labeled with one of the following dyes: PET, 6-FAM, or VIC. This ensured accurate assignment of PCR products to a specific VNTR locus after capillary electrophoresis. Table 2 lists the selected VNTR

loci and forward primers with their corresponding fluorescent label.

The optimized MLVA protocol was obtained as follows. Template DNA was prepared as described above. PCR was performed using the Qiagen Type-it Microsatellite PCR Kit (model 206243; Qiagen, Hilden, Germany) in two mixes, each in a total volume of 25  $\mu$ L. The first PCR reaction contained 12.5  $\mu$ L of mastermix, 2.5  $\mu$ L of Q-solution, 3.2  $\mu$ M of primer SE7b, 0.04  $\mu$ M of primer SE9, 0.08  $\mu$ M of primer ENTR13, 0.12  $\mu$ M of primer SENTR6, and 1  $\mu$ L of template DNA. The second PCR reaction contained 12.5  $\mu$ L of mastermix, 2.5  $\mu$ L of Q-solution, 0.16  $\mu$ M of primer SE5, 0.12  $\mu$ M of primer SENTR1, and 1  $\mu$ L of template DNA. PCR reactions were performed in a GeneAmp 9700 PCR system (Applied Biosystems, Foster City, CA). Cycling conditions for the first PCR reaction were 94°C for 5 min, followed by 30 cycles of 94°C for 30 sec, 62°C for 1 min, and 72°C for 1 min. A final extension of 72°C for 5 min was employed. Cycling conditions for the second PCR reaction were 94°C for 5 min, followed by 20 cycles of 94°C for 30 sec, 60°C for 1 min, and 72°C for 1 min with a final extension of 72°C for 5 min. Both PCR products were mixed in equal amounts before capillary electrophoresis on ABI PRISM<sup>®</sup> 3130 Genetic Analyzer (Applied Biosystems) with the GENESCAN<sup>™</sup>-1200 LIZ<sup>®</sup> Size Standard. Fragment sizes/repeat numbers were assigned for each locus for analysis with BioNumerics software version 6.5 using the MLVA plugin (Applied Maths, Sint-Martens-Latem, Belgium).

### Isolate collection for comparison of MLVA and PFGE

This study included 60 SE isolates used to compare the typeability and discriminatory power of the optimized MLVA and PFGE. The selection included the ATCC 13076 strain plus 47 isolates of various origins, year of isolation, and phage types considered to be epidemiologically unrelated isolates. In addition, we examined 12 isolates (FODSE) from four layer farms (Farms A, B, C, and H), representing four sets of possibly closely related isolates (Table 1, panel 2). PT of the SE isolates was performed according to the PT scheme of (Ward *et al.*, 1987) at the National Reference Centre for *Salmonella* (Scientific Institute of Public Health, Brussels, Belgium).

### PFGE

Preparation of agarose plugs, cell lysis, and washing of agarose plugs was performed according to the PulseNet protocol ([www.cdc.gov/pulsenet/](http://www.cdc.gov/pulsenet/)). Plug slices were digested for 18 h with 30 U of *Xba*I, *Not*I, or *Sfi*I (New England BioLabs, Ipswich, MA) with a digestion temperature of 37°C for *Xba*I and *Not*I and 50°C for *Sfi*I. DNA fragments were separated by the CHEF mapper (Bio-Rad, La Jolla, CA) in a 1% Seakem gold agarose (Lonza, Rockland, MA). The running conditions were 6V/cm at 14°C in 0.5  $\times$  Tris/borate/EDTA (TBE) buffer for 19 h with a ramping time from 2.16 to 63.8 s for the *Xba*I enzyme, 24 h with a ramping time from 2 to 10 s for the *Not*I enzyme, and 24 h with a ramping time from 2 to 12 s for the *Sfi*I enzyme. Gels were stained with ethidium bromide, destained in water, and digitally captured under ultraviolet light. PFGE profiles were clustered with BioNumerics version 6.5 (Applied Maths) using *Salmonella* Braenderup H9812 digested with *Xba*I as a normalization reference. Similarities between the fingerprints were calculated using the Dice coefficient (with an optimization of 1% and a position

TABLE 1. *SALMONELLA* ENTERITIDIS ISOLATES USED TO EVALUATE THE TYPEABILITY AND DISCRIMINATORY POWER OF THE MULTIPLE-LOCUS VARIABLE NUMBER TANDEM-REPEAT ASSAY PRIMERS TESTED (PANEL 1) AND *SALMONELLA* ENTERITIDIS ISOLATES

Strain ID	Origin	Year	MLVA type	PT	PFGE type		
					XbaI	NotI	SfiI
<b>Panel 1</b>							
1. MB 1535	Deer, Belgium	1999		RDNC/P20			
2. MB 1717	Nerve node of pig, Belgium	2001		PT 4			
3. KS 94	Overshoes poultry Farm Y, Belgium	1999		PT 21			
4. KS 104	Eggshell, poultry Farm Z, Belgium	1999		PT 4			
5. 02-10562	Human, Belgium	2002		PT 21			
6. 07-06092	Human, Belgium	2007		PT 6			
7. FODSE 5	Layer Farm A, Belgium	2008		PT 8			
8. FODSE 130	Layer Farm B, Belgium	2008		PT 35			
9. MB 2045	Cheese, food outbreak S, Belgium	2001		PT 21			
10. MB 2046	Mayonnaise, food outbreak S, Belgium	2001		PT 21			
11. MB 2047	Smoked salmon, food outbreak S, Belgium	2001		PT 21			
12. MB 2048	Human, food outbreak S, Belgium	2001		PT 21			
<b>Panel 2</b>							
MB 2499	Lizard, Belgium	2002	A	PT 6a	Xba-10	Not-9	Sfi-3
SA07 1377	Layer farm, Belgium	2007	B	PT 4	Xba-1	Not-1	Sfi-6
FODSE 13	Layer Farm A, HH 2 laying round 1, Belgium	2008	B	PT 8	Xba-5	Not-3	Sfi-16
FODSE 85	Layer Farm A, HH 1 laying round 2, Belgium	2008	B	PT 23	Xba-5	Not-3	Sfi-16
04-10630	Human, Belgium	2004	C	RDNC 69	Xba-1	Not-2	Sfi-13
06-02195	Human, Belgium	2006	C	PT 1	Xba-1	Not-1	Sfi-16
MB 1456	Egg, Denmark	1999	C	PT 8	Xba-7	Not-3	Sfi-16
FODSE 189	Layer Farm B, HH 1 laying round 2, Belgium	2009	D	PT 1b	Xba-1	Not-6	Sfi-16
FODSE 229	Layer Farm C, HH laying round 1, Belgium	2009	D	PT 23	Xba-5	Not-3	Sfi-10
FODSE 210	Layer Farm C, HH laying round 1, Belgium	2009	D	PT 23	Xba-5	Not-7	Sfi-8
04-01032	Human, Belgium	2004	E	PT 4	Xba-1	Not-1	Sfi-16
MB 1175	Egg, Slovakia	1997	E	PT 8	Xba-5	Not-3	Sfi-16
05-05050	Human, Belgium	2005	E	PT 9a	Xba-9	Not-8	Sfi-1
SA02 478	Layer farm, Belgium	2002	F	PT 7	Xba-1	Not-1	Sfi-8
05-01202	Human, Belgium	2005	G	PT 6	Xba-1	Not-1	Sfi-12
MB 1535	Deer, Belgium	1999	H	RDNC/P20	Xba-9	Not-8	Sfi-2
02-01276	Human, Belgium	2002	I	PT 4	Xba-1	Not-1	Sfi-6
FODSE 157	Layer Farm B, HH 1 laying round 1, Belgium	2008	J	PT 4b	Xba-1	Not-6	Sfi-16
FODSE 26	Layer Farm A, ECA laying round 1, Belgium	2008	K	PT 7a	Xba-5	Not-7	Sfi-16
07-01032	Human, Belgium	2007	L	PT 4	Xba-1	Not-1	Sfi-16
03-08402	Human, Belgium	2003	L	PT 6a	Xba-2	Not-1	Sfi-16
07-00351	Human, Belgium	2007	L	PT 21	Xba-8	Not-4	Sfi-16
SA05 1205	Layer farm, Belgium	2005	L	PT 35	Xba-3	Not-4	Sfi-8
SA06 1660	Layer farm, Belgium	2006	M	PT 6a	Xba-3	Not-4	Sfi-9
04-06044	Human, Belgium	2004	M	PT 8	Xba-5	Not-3	Sfi-4
MB 2591	Pigeon, Belgium	2001	M	PT 4	Xba-1	Not-1	Sfi-16
03-04715	Human, Belgium	2003	M	PT 14b	Xba-1	Not-1	Sfi-16
FODSE 169	Layer Farm B, HH 1 laying round 1, Belgium	2008	M	RDNC 52	Xba-1	Not-6	Sfi-16
KS 104	Eggshell, poultry Farm Z, Belgium	1999	M	PT 4	Xba-1	Not-1	Sfi-6
SA07 794	Layer farm, Belgium	2007	M	PT 1	Xba-1	Not-1	Sfi-6
07-02806	Human, Belgium	2007	M	PT 6	Xba-3	Not-1	Sfi-16
MB 1355	Pastry, Belgium	1999	M	PT 4	Xba-4	Not-1	Sfi-14
MB 1717	Nerve node of pig, Belgium	2001	M	PT 4	Xba-4	Not-1	Sfi-16
SA02 596	Layer farm, Belgium	2002	N	PT 21	Xba-3	Not-4	Sfi-12
06-03044	Human, Belgium	2006	N	PT 8	Xba-5	Not-3	Sfi-16
SA00 575	Layer farm, Belgium	2000	O	PT 6a	Xba-1	Not-1	Sfi-12
FODSE 321	Layer Frm H, ECA laying round 1, Belgium	2009	O	PT 21c	Xba-3	Not-4	Sfi-12
MB 1842	Dairy environment, Belgium	2001	O	PT 4	Xba-1	Not-1	Sfi-15
MB 2045	Cheese, food outbreak A, Belgium	2001	O	PT 21	Xba-1	Not-1	Sfi-16
MB 1221	Tiramisu, Belgium	1998	O	PT 6	Xba-1	Not-1	Sfi-16
MB 1425	Egg, the Netherlands	1999	O	PT 1	Xba-1	Not-1	Sfi-16
FODSE 288	Layer Farm H, HH 2 laying round 1, Belgium	2009	O	PT 1b	Xba-3	Not-4	Sfi-12
MB 2602	Rabbit, Belgium	2000	O	NT	Xba-3	Not-4	Sfi-15
FODSE 317	Layer Frm H, ECA laying round 1, Belgium	2009	O	PT 21c	Xba-3	Not-4	Sfi-16
MB 2609	Bird, Belgium	2000	O	PT 21	Xba-3	Not-4	Sfi-6
KS 94	Overshoes poultry Frm Y, Belgium	1999	O	PT 21	Xba-3	Not-4	Sfi-6
02-00941	Human, Belgium	2002	O	PT 1	Xba-3	Not-4	Sfi-6

(continued)

TABLE 1. (CONTINUED)

Strain ID	Origin	Year	MLVA		PFGE type		
			type	PT	XbaI	NotI	SfiI
FODSE 258	Layer Frm C, HH laying round 1, Belgium	2009	O	PT 28	Xba-5	Not-3	Sfi-12
06-02542	Human, Belgium	2006	O	PT 28	Xba-5	Not-3	Sfi-16
MB 2588	Sludge, Belgium	2002	O	PT 17	Xba-1	Not-1	Sfi-16
SA00 367	Layer farm, Belgium	2000	O	PT 14b	Xba-1	Not-5	Sfi-16
SA06 407	Layer farm, Belgium	2006	O	PT 34	Xba-1	Not-1	Sfi-7
SA05 306	Layer farm, Belgium	2005	O	PT 4a	Xba-1	Not-1	Sfi-8
MB 1418	Egg, Austria	1995	O	PT 21	Xba-3	Not-4	Sfi-16
02-09574	Human, Belgium	2002	O	PT 21	Xba-3	Not-4	Sfi-16
SA03 1406	Layer farm, Belgium	2003	O	PT 6	Xba-3	Not-4	Sfi-8
SA03 2252	Layer farm, Belgium	2003	O	PT 8	Xba-5	Not-3	Sfi-11
03-08145	Human, Belgium	2003	O	PT 8	Xba-6	Not-3	Sfi-16
05-02959	Human, Belgium	2005	P	PT 34	Xba-8	Not-3	Sfi-5
ATCC 13076 <sup>T</sup>			F	RDNC961	Xba-5	Not-3	Sfi-16

The isolates' corresponding origin, year of isolation, and phage type are noted. Isolates are grouped according to the results obtained by MLVA and PFGE (panel 2).

MLVA, multiple-locus variable number tandem-repeat assay; PFGE, pulsed-field gel electrophoresis; PT, phage typing; RDNC, reacted but did not conform with any standard phage pattern; HH, henhouse; ECA, egg collecting area.

tolerance of 0.7–1.7%) and the unweighted-pair group method using arithmetic averages algorithm (UPGMA).

#### Delineation of MLVA types and pulsotypes

The VNTR code was defined in the following order: ENTR13—SE5—SE7b—SE9—SENTR1—SENTR6. An MLVA type was assigned based on a difference in repeat numbers of at least one repeat in one VNTR locus. MLVA types were indicated by capital letters. For each PFGE restriction enzyme, a corresponding pulsotype was assigned based on the difference in presence, absence, or clear shift of at least one band in the PFGE fingerprint (Gatto *et al.*, 2006). A pulsotype was indicated by the name of the restriction enzyme followed by a number (e.g., Xba-1).

#### Calculation of discriminatory power and concordance

The discriminatory index (DI) was calculated as described by Hunter and Gaston (Hunter, 1990; Hunter and Gaston, 1988) on the collection of 47 epidemiologically unrelated isolates and the ATCC 13076 strain. In addition, Wallace's coefficient (*W*) was determined together with the proposed Wallace 95% confidence interval (CI) and Wallace's coefficient

under independence (*W<sub>i</sub>*) (Carrico *et al.*, 2006; Pinto *et al.*, 2008). The *W* coefficient indicates the probability that two isolates classified as the same type by one method will also be classified as the same type when using the other method (Rasschaert *et al.*, 2009). If the *W* value is not significantly different from the *W<sub>i</sub>* value, one can conclude that such congruence of classification could arise by chance.

## Results

#### Analysis of the 60 SE isolates using MLVA

All tested SE isolates (Table 1, panel 2) were typeable using the optimized MLVA, except MB 2499, where only two primer pairs (ENTR13, SE9) generated a band. Based on the given VNTR codes, two main clusters and one separate isolate (MB 2499) were generated. In total, 16 allele combinations or MLVA types were found among the 60 SE isolates tested (Table 1 and Fig. 1). VNTR loci SE5 and SENTR6 showed the highest variation.


MLVA was able to discriminate among isolates from different layer farms and between isolates from the same layer farm, except within Farm H. MLVA profiles with their respective pulsotype (*XbaI*, *NotI*, and *SfiI*) and PT type are shown in Figure 1.

TABLE 2. PRIMERS SELECTED FOR USE IN THE OPTIMIZED MULTIPLE-LOCUS VARIABLE NUMBER TANDEM-REPEAT ASSAY

Locus	Primers	Primer sequence (5'-3')	PCR mix	Reference
<i>ushA</i>	SE9-F	PET-CGTAGCCAATCAGATTCATCCC	1	Cho <i>et al.</i> , 2007
	SE9-R	GCGTTTGAAACGGGGTGTGGCGCTG		
<i>yohM</i>	SE5-F	PET-CGGGAAACCACCATCAC	2	Cho <i>et al.</i> , 2007
	SE5-R	CAGGCCGAACAGCAGGAT		
<i>ygbF</i>	SE7b-F	FAM-GATAATGCTGCCGTGGTAA	1	Malorny <i>et al.</i> , 2008
	SE7b-R	ACTGCGTTGGTTTCTTTTCT		
Non-coding	SENTR6-F	FAM-ATGGACGGAGGCGATAGAC	1	Malorny <i>et al.</i> , 2008
	SENTR6-R	AGCTTACAATTTGCGTATTTCG		
<i>tolA</i>	SENTR1-F	VIC-GCAACAGCAGCAGCAACAG	2	Malorny <i>et al.</i> , 2008
	SENTR1-R	CCGAGCTGAGATCGCCAAG		
Non-coding	ENTR13-F	VIC-TATGAACCAATGGCAACGAGAC	1	Beranek <i>et al.</i> , 2009
	ENTR13-R	CGTGGCAAGGAACAGTAGAGG		


PCR, polymerase chain reaction.





(\*) VNTR code -2.0 was given (by the Bionumerics software programme) when no feasible band was detected

FIG. 1. Dendrogram and repeat numbers of each variable number tandem repeat (VNTR) locus for multiple-locus variable number tandem-repeat assay (MLVA) performed on 60 *Salmonella* Enteritidis (SE) isolates. The similarities between the VNTR codes were calculated using categorical values and the fingerprints were grouped according to their similarities using the unweighted-pair group method using arithmetic averages algorithm (UPGMA) algorithm.



**FIG. 2.** Dendrogram and fingerprints for pulsed-field gel electrophoresis (PFGE) using restriction enzyme *XbaI* with 60 *Salmonella* Enteritidis (SE) isolates. The similarities between the fingerprints were calculated using the Dice coefficient (optimization 1.0% and position tolerance 1.5%), and the fingerprints were grouped according to their Dice similarities using the unweighted-pair group method using arithmetic averages algorithm (UPGMA) algorithm. *XbaI* pulstypes are given with their respective *NotI* and *SfiI* pulstypes and multiple-locus variable number tandem-repeat assay (MLVA) type.

*Analysis of the 60 SE isolates using PFGE*

All isolates (Table 1, panel 2) were typeable by PFGE using *XbaI*, *SfiI*, and *NotI* restriction analysis. A cut-off value of 97% for *XbaI* and *SfiI* and 96% for *NotI* for delineation of the dif-

ferent pulstypes was determined, according to the criteria for the delineation of pulstypes as described above.

Ten *XbaI* (Table 1 and Fig. 2) and nine *NotI* (Table 1 and Fig. 3) pulstypes were determined within the isolates. Using *SfiI* (Table 2 and Fig. 4), 16 pulstypes were distinguished. For





FIG. 3. Dendrogram and fingerprints for pulsed-field gel electrophoresis (PFGE) using restriction enzyme *NotI* with 60 *Salmonella* Enteritidis (SE) isolates. The similarities between the fingerprints were calculated using the Dice coefficient (optimization 1.0% and position tolerance 0.7%), and the fingerprints were grouped according to their Dice similarities using the unweighted-pair group method using arithmetic averages algorithm (UPGMA) algorithm. *NotI* pulsotypes are given with their respective *XbaI* and *SfiI* pulsotype and multiple-locus variable number tandem-repeat assay (MLVA) type.

each method, the MB 2499 isolate formed a separate pulso-type from the other SE isolates.

Using PFGE *XbaI* or *NotI*, it was not possible to discriminate between isolates from Farms A and C, whereas PFGE using *SfiI* found the same pulsotypes on Farms A and B. PFGE using

*NotI* was restricted to discriminate only within isolates recovered from Farm A or from Farm C. PFGE using *SfiI* could discriminate within isolates within Farm C or Farm H. PFGE using *XbaI* could not discriminate among any isolates within the same farm.



**FIG. 4.** Dendrogram and fingerprints for pulsed-field gel electrophoresis (PFGE) using restriction enzyme *SfiI* with 60 *Salmonella* Enteritidis (SE) isolates. The similarities between the fingerprints were calculated using the Dice coefficient (optimization 1.0% and position tolerance 1.7%), and the fingerprints were grouped according to their Dice similarities using the unweighted-pair group method using arithmetic averages algorithm (UPGMA) algorithm. *SfiI* pulsotypes are given with their respective *XbaI* and *NotI* pulsotype and multiple-locus variable number tandem-repeat assay (MLVA) type.

*Discriminatory power and Wallace coefficient*

The discriminatory index (DI) of each method was determined separately and combined with PT, as calculated for the 48 epidemiologically unrelated SE isolates and the ATCC

13076 strain (layer farm isolates FODSE were not included; Table 3). For each method considered separately, the discriminatory power of PFGE using *SfiI*, *XbaI* or *NotI* was lower (DI=0.77, 0.75, and 0.69, respectively) compared to MLVA (DI=0.80). Combining PT with MLVA (DI=0.98) or PFGE



TABLE 3. DISCRIMINATORY POWER OF THE VARIOUS METHODS (INDIVIDUALLY AND IN COMBINATION WITH PT) EVALUATED ON 48 EPIDEMIOLOGICALLY UNRELATED SE ISOLATES

Method/combination	No. of types	No. of unique isolates	No. of clustered isolates	Cluster size	DI
PFGE <i>NotI</i>	7	3	45	2–23	0.69
PFGE <i>XbaI</i>	10	4	44	2–21	0.75
PFGE <i>SfiI</i>	15	10	38	2–22	0.77
PFGE (all)	28	20	28	2–9	0.95
MLVA	13	6	42	2–19	0.80
PT+PFGE <i>NotI</i>	25	19	29	2–9	0.94
PT+PFGE <i>XbaI</i>	29	21	27	2–7	0.96
PT+PFGE <i>SfiI</i>	32	23	25	2–4	0.98
PT+PFGE (all)	40	34	14	2–3	0.99
PT+MLVA	38	33	15	2–5	0.98
Total	46	44	4	2	1.00*

\*Exact value is 0.998, because there were two times two clustered isolates, which could not be distinguished using either of the typing methods.

PT, phage typing; SE, *Salmonella* Enteritidis; DI, discriminatory index; PFGE, pulsed-field gel electrophoresis.

using *SfiI* (DI=0.98) resulted in more discriminatory power than combining PT with PFGE using restriction enzyme *XbaI* or *NotI* (DI=0.96 and 0.94, respectively).

The congruence between typing methods, expressed by the Wallace coefficient (*W*), is shown in Table 4. When comparing PT with another typing method, the highest correlation was found between the information provided by PT and PFGE using *NotI* in both directions. When comparing MLVA with another typing method, the highest correlation was observed with PFGE using *XbaI* or *NotI*. A high bidirectional correspondence between PFGE was seen when using *XbaI* and *NotI*. However, for this data set, *W* values were very low and most of the calculated 95% confidence intervals (CI) for *W* included the respective Wallace coefficient under independence (*W<sub>i</sub>*). This indicates that the congruence of classification could have arisen by chance.

## Discussion

Several MLVA typing schemes for the characterization of SE have been described (Beranek *et al.*, 2009; Boxrud *et al.*, 2007; Cho *et al.*, 2007, 2008; Malorny *et al.*, 2008; Ross and Heuzenroeder, 2009). However, the use of different loci in each protocol and different primers for the same loci makes it difficult to select the most suitable MLVA scheme. In addition, the different conditions used for running and analysing PCR greatly hinder interlaboratory comparison of the results of the test (Hopkins *et al.*, 2011), which was encountered by our laboratory staff. We therefore evaluated primer pairs from

existing MLVA systems for their typeability and discriminatory power and developed an optimized MLVA capillary electrophoresis protocol for the characterization of SE isolates using a new primer combination. Typeability and discriminatory power of this six-locus MLVA were compared with PFGE using restriction enzymes *XbaI*, *NotI*, or *SfiI* on a diverse collection of SE isolates. In this way, we determined the most suitable genotyping method to use in addition to PT. For the different typing methods, we also determined *W* to analyze correspondence among the classifications of the typing methods. Epidemiologically unrelated (SE isolates with different origins collected over several years) as well as SE isolates sampled on the same layer farm were used to define a suitable subtyping method or a polyphasic approach (combination of typing methods). This enabled us to evaluate their practical use (i.e., a sufficiently high discriminatory power) for the following epidemiological purposes: (i) to distinguish among epidemiologically unrelated SE isolates over several years, (ii) to compare SE isolates originating from layer farms and from human origin, and (iii) to describe contamination routes on SE contaminated layer farms. Results of this study showed that the optimized MLVA method had higher discriminatory power in comparison to PFGE performed with a single restriction enzyme (*XbaI*, *SfiI*, or *NotI*). Only a combination of these three enzymes in PFGE had a considerably higher discriminatory power than MLVA. However, the combination of MLVA with PT had a discriminatory power comparable to combining PT with PFGE using all three enzymes. For any given typing method, *W* provides an estimate

TABLE 4. VALUES OF *W<sub>i</sub>* AND *W* WITH CORRESPONDING 95% CI FOR THE TYPING METHODS BETWEEN BRACKETS

	PT	MLVA	PFGE <i>XbaI</i>	PFGE <i>NotI</i>	PFGE <i>SfiI</i>
<i>W<sub>i</sub></i>	0.082	0.196	0.251	0.306	0.232
PT		0.207 (0.055–0.358)	0.478 (0.296–0.660)	0.793 (0.686–0.901)	0.283 (0.141–0.425)
MLVA	0.086 (0.002–0.169)		0.303 (0.188–0.418)	0.339 (0.224–0.455)	0.240 (0.094–0.386)
PFGE <i>XbaI</i>	0.155 (0.056–0.255)	0.237 (0.107–0.367)		0.823 (0.650–0.996)	0.233 (0.103–0.364)
PFGE <i>NotI</i>	0.212 (0.097–0.327)	0.217 (0.115–0.320)	0.675 (0.488–0.863)		0.278 (0.138–0.419)
PFGE <i>SfiI</i>	0.099 (0.038–0.160)	0.202 (0.082–0.323)	0.252 (0.111–0.392)	0.366 (0.223–0.509)	

*W*, Wallace's coefficient; *W<sub>i</sub>*, Wallace's coefficient under independence; PT, phage typing; MLVA, multiple-locus variable number tandem-repeat assay; PFGE, pulsed-field gel electrophoresis.

of how much new information is obtained from another typing method. Results indicate that partitions defined either by PT, MLVA, or PFGE using *Xba*I or *Sfi*I could have been best predicted by PFGE using *Not*I and vice versa. A combination of methodologies likely provides additional information; however, due to the limited number of isolates tested, the estimated *W* value was very low and in most cases not significantly different from *W<sub>i</sub>*. No reliable information could be obtained on the directional agreement between the typing methods tested (Pinto *et al.*, 2008; Severiano *et al.*, 2011).

All methods showed good discrimination between isolates from different layer farms. However, only the optimized MLVA, PFGE using restriction enzyme *Sfi*I, and PFGE using all three restriction enzymes provided high resolution for SE isolates from the same layer farm.

The optimized MLVA showed good epidemiological concordance because the isolates from a single-strain outbreak were assigned to identical types (data not shown), which was also confirmed by Boxrud *et al.* (2007). MLVA typing, albeit with a different combination of primers, has been shown to provide enhanced resolution and good reproducibility for characterizing SE (Boxrud *et al.*, 2007; Cho *et al.*, 2007). Cho *et al.* (2007) also found that MLVA (although with a different combination of primers than ours) had a higher discriminatory power than PFGE combined with PT as tested on various SE isolates from human and non-human sources. Cho *et al.* (2010) showed that MLVA (with a composition of seven primers) in combination with PT can be used for effective characterization of SE isolates collected from sporadic human clinical cases. Although they found an association of MLVA-based clusters with phage types using human clinical isolates, this was not confirmed by the present study in which a more diverse and smaller collection of isolates was used.

In conclusion, this optimized MLVA method provides good discriminatory power for characterizing SE isolates. The actual isolate diversity observed by PT could not be obtained by the use of MLVA. A combination of PT and MLVA seems to be providing a higher discriminatory power, as literature and the results obtained from the present study would indicate. In terms of discriminating between SE isolates of different origin and studying contamination routes on a particular layer farm, MLVA, PFGE using *Sfi*I, and PFGE combining all three restriction enzymes can all be used. Nevertheless, MLVA has several advantages over PFGE. MLVA has good discriminatory capacity and has a high throughput because it is a PCR- and capillary-electrophoresis-based technique. These characteristics make MLVA less labor-intensive than PFGE, because the data generated are easier to analyze and interpret. MLVA is thus appropriate for epidemiological studies with a large collection of strains.

### Acknowledgments

We thank Rik Lenaerts, Séverine Danschutter, Pieter Siau, and Sabine Van Glabeke for excellent technical assistance, and Miriam Levenson for the English language editing of this manuscript. This research was funded by the Belgian Federal Public Service for Health, Food Chain Safety and Environment (RF 6195).

### Disclosure Statement

No competing financial interests exist.

### References

- Beranek A, Mikula C, Rabold P, Arnhold D, Berghold C, Lederer I, Allerberger F, Kornschöber C. Multiple-locus variable-number tandem repeat analysis for subtyping of *Salmonella enterica* subsp. *enterica* serovar Enteritidis. *Int J Med Microbiol* 2009;299:43–51.
- Boxrud D, Pederson-Gulrud K, Wotton J, Medus C, Lyszkwicz E, Besser J, Bartkus JM. Comparison of multiple-locus variable-number tandem repeat analysis, pulsed-field gel electrophoresis, and phage typing for subtype analysis of *Salmonella enterica* serotype Enteritidis. *J Clin Microbiol* 2007;45:536–543.
- Brown DJ, Baggesen DL, Platt DJ, Olsen JE. Phage type conversion in *Salmonella enterica* serotype Enteritidis caused by the introduction of a resistance plasmid of incompatibility group X (IncX). *Epidemiol Infect* 1999;122:19–22.
- Carrico JA, Silva-Costa C, Melo-Cristino J, Pinto FR, de Lencastre H, Almeida JS, Ramirez M. Illustration of a common framework for relating multiple typing methods by application to macrolide-resistant *Streptococcus pyogenes*. *J Clin Microbiol* 2006;44:2524–2532.
- Chart H, Row B, Threlfall EJ, Ward LR. Conversion of *Salmonella enteritidis* phage type 4 to phage type 7 involves loss of lipopolysaccharide with concomitant loss of virulence. *FEMS Microbiol Lett* 1989;51:37–40.
- Cho S, Boxrud DJ, Bartkus JM, Whittam TS, Saeed M. Multiple-locus variable-number tandem repeat analysis of *Salmonella* Enteritidis isolates from human and non-human sources using a single multiplex PCR. *FEMS Microbiol Lett* 2007;275:16–23.
- Cho S, Whittam TS, Boxrud DJ, Bartkus JM, Saeed AM. Allele distribution and genetic diversity of VNTR loci in *Salmonella enterica* serotype Enteritidis isolates from different sources. *BMC Microbiol* 2008;8:146.
- Cho S, Whittam TS, Boxrud DJ, Bartkus JM, Rankin SC, Wilkins MJ, Somsel P, Downes FP, Musser KA, Root TP, Warnick LD, Wiedmann M, Saeed AM. Use of multiple-locus variable number tandem repeat analysis and phage typing for subtyping of *Salmonella* Enteritidis from sporadic human cases in the United States. *J Appl Microbiol* 2010;108:859–867.
- De Lappe N, Doran G, O'Connor J, O'Hare C, Cormican M. Characterization of bacteriophages used in the *Salmonella enterica* serovar Enteritidis phage-typing scheme. *J Med Microbiol* 2009;58:86–93.
- [EFSA] European Food Safety Authority. Scientific opinion on a quantitative estimation of the public health impact of setting a new target for the reduction of *Salmonella* in laying hens. *EFSA J* 2010;8:1546 [86 pp.]. Available at: [www.efsa.europa.eu/en/efsajournal/doc/1546.pdf](http://www.efsa.europa.eu/en/efsajournal/doc/1546.pdf), accessed August 20, 2012.
- Foley SL, Zhao SH, Walker RD. Comparison of molecular typing methods for the differentiation of *Salmonella* foodborne pathogens. *Foodborne Pathog Dis* 2007;4:253–276.
- Foxman B, Zhang L, Koopman JS, Manning SD, Marrs CF. Choosing an appropriate bacterial typing technique for epidemiologic studies. *Epidemiol Perspect Innov* 2005;2:10.
- Gantois I, Ducatelle R, Pasmans F, Haesebrouck F, Gast R, Humphrey TJ, Van Immerseel F. Mechanisms of egg contamination by *Salmonella* Enteritidis. *FEMS Microbiol Rev* 2009;33:718–738.
- Gatto AJ, Peters TM, Green J, Fisher IS, Gill ON, O'Brien SJ, Maguire C, Berghold C, Lederer I, Gerner-Smidt P, Torpdahl M, Siitonen A, Lukinmaa S, Tschape H, Prager R, Luzzi I, Dionisi AM, van der Zwaluw WK, Heck M, Coia J, Brown D, Usera M, Echeita A, Threlfall EJ. Distribution of molecular subtypes within *Salmonella enterica* serotype Enteritidis phage type 4 and *S. Typhimurium* definitive phage type 104 in nine

- European countries, 2000–2004: Results of an international multi-centre study. *Epidemiol Infect* 2006;134:729–736.
- Hopkins KL, Peters TM, de Pinna E, Wain J. Standardisation of multilocus variable-number tandem-repeat analysis (MLVA) for subtyping of *Salmonella enterica* serovar Enteritidis. *Euro Surveill* 2011;16: pii: 19942.
- Hunter PR, Gaston MA. Numerical index of the discriminatory ability of typing systems—An application of Simpsons Index of diversity. *J Clin Microbiol* 1988;26:2465–2466.
- Hunter PR. Reproducibility and indices of discriminatory power of microbial typing methods. *J Clin Microbiol* 1990;28:1903–1905.
- Kang ZW, Jung JH, Kim SH, Lee BK, Lee DY, Kim YJ, Lee JY, Won HK, Kim EH, Hahn TW. Genotypic and phenotypic diversity of *Salmonella* Enteritidis isolated from chickens and humans in Korea. *J Vet Med Sci* 2009;71:1433–1438.
- Kruy SL, van Cuyck H, Koeck JL. Multilocus variable number tandem repeat analysis for *Salmonella enterica* subspecies. *Eur J Clin Microbiol Infect Dis* 2011;30:465–473.
- Laconcha I, Baggesen DL, Rementeria A, Garaizar J. Genotypic characterisation by PFGE of *Salmonella enterica* serotype Enteritidis phage types 1, 4, 6, and 8 isolated from animal and human sources in three European countries. *Vet Microbiol* 2000;75:155–165.
- Lapuz R, Tani H, Sasai K, Shirota K, Katoh H, Baba E. An epidemiological analysis of *Salmonella* Enteritidis contamination in a rat-infested chicken layer farm, an egg processing facility, and liquid egg samples by pulsed-field gel electrophoresis. *J Vet Med Sci* 2007;69:649–652.
- Lukinmaa S, Schildt R, Rinttila T, Siitonen A. *Salmonella* Enteritidis phage types 1 and 4: Pheno- and genotypic epidemiology of recent outbreaks in Finland. *J Clin Microbiol* 1999;37:2176–2182.
- Majtanova L, Majtan J, Majtan V. Trends in phage types of *Salmonella enterica* serovars Enteritidis and Typhimurium isolated in Slovakia from 1995 to 2009. *Diagn Microbiol Infect Dis* 2011;69:454–456.
- Malorny B, Junker E, Helmuth R. Multi-locus variable-number tandem repeat analysis for outbreak studies of *Salmonella enterica* serotype Enteritidis. *BMC Microbiol* 2008;8:84.
- Much P, Pichler J, Kasper S, Lassnig H, Kornschöber C, Buchner A, König C, Allerberger F. A foodborne outbreak of *Salmonella* Enteritidis phage type 6 in Austria, 2008. *Wien Klin Wochenschr* 2009;121:132–136.
- Pang JC, Chiu TH, Chiou CS, Schroeter A, Guerra B, Helmuth R, Tsen HY. Pulsed-field gel electrophoresis, plasmid profiles and phage types for the human isolates of *Salmonella enterica* serovar Enteritidis obtained over 13 years in Taiwan. *J Appl Microbiol* 2005;99:1472–1483.
- Pinto FR, Melo-Cristino J, Ramirez M. A confidence interval for the Wallace coefficient of concordance and its application to microbial typing methods. *PLoS One* 2008;3:e3696.
- Rasschaert G, Vanderhaeghen W, Dewaele I, Janez N, Huijsdens X, Butaye P, Heyndrickx M. Comparison of fingerprinting methods for typing methicillin-resistant *Staphylococcus aureus* sequence type 398. *J Clin Microbiol* 2009;47:3313–3322.
- Rivoal K, Protais J, Queguiner S, Boscher E, Chidaine B, Rose V, Gautier M, Baron F, Grosset N, Ermel G, Salvat G. Use of pulsed-field gel electrophoresis to characterize the heterogeneity and clonality of *Salmonella* serotype Enteritidis, Typhimurium and Infantis isolates obtained from whole liquid eggs. *Int J Food Microbiol* 2009;129:180–186.
- Ross IL, Heuzenroeder MW. A comparison of two PCR-based typing methods with pulsed-field gel electrophoresis in *Salmonella enterica* serovar Enteritidis. *Int J Med Microbiol* 2009;299:410–420.
- Saeed AM, Walk ST, Arshad M, Whittam TS. Clonal structure and variation in virulence of *Salmonella* Enteritidis isolated from mice, chickens, and humans. *J AOAC Int* 2006;89:504–511.
- Severiano A, Pinto FR, Ramirez M, Carrico JA. Adjusted Wallace coefficient as a measure of congruence between typing methods. *J Clin Microbiol* 2011;49:3997–4000.
- Tankouo-Sandjong B, Kinde H, Wallace I. Development of a sequence typing scheme for differentiation of *Salmonella* Enteritidis strains. *FEMS Microbiol Lett* 2012;331:165–175.
- Threlfall EJ, Rowe B, Ward LR. Subdivision of *Salmonella* Enteritidis phage types by plasmid profile typing. *Epidemiol Infect* 1989;102:459–465.
- Van Belkum A. Tracing isolates of bacterial species by multi-locus variable number of tandem repeat analysis (MLVA). *FEMS Immunol Med Microbiol* 2007;49:22–27.
- Ward LR, Desa JDH, Rowe B. A phage-typing scheme for *Salmonella* Enteritidis. *Epidemiol Infect* 1987;99:291–294.

Address correspondence to:

G. Rasschaert, Ph.D.

Technology and Food Science Unit

Institute for Agricultural and Fisheries Research (ILVO)

Brusselsesteenweg 370

9090 Melle, Belgium

E-mail: geertrui.rasschaert@ilvo.vlaanderen.be