Lymph drainage from the ovine tonsils: an anatomical study of the tonsillar lymph vessels

C. Casteleyn^{1*}, P. Cornillie¹, C. Van Ginneken², P. Simoens¹, S. Van Cruchten², K. Vandewelde¹, W. Van den Broeck¹

¹Department of Morphology, Faculty of Veterinary Medicine, Ghent University,
Salisburylaan 133, 9820 Merelbeke, Belgium

²Laboratory of Applied Veterinary Morphology, Department of Veterinary Sciences, Faculty
of Pharmaceutical, Biomedical and Veterinary Sciences, University of Antwerp,
Universiteitsplein 1, 2610 Wilrijk, Belgium

*Corresponding author with present address: Laboratory of Applied Veterinary Morphology,
Department of Veterinary Sciences, Faculty of Pharmaceutical, Biomedical and Veterinary
Sciences, University of Antwerp, Wilrijk, Belgium. Tel.: +32 3-265-28-21; fax: +32 3-265-
24-33; e-mail: Christophe.Casteleyn@ua.ac.be (C. Casteleyn)

With 6 figures

Summary

Although the tonsils of sheep have gained much attention during the last decade, only few data are available on their lymph vessel architecture. Tonsillar lymph vessels are immunologically important as they form the efferent routes for locally activated immune cells to reach the draining lymph nodes. To gain insight into the tonsillar lymph drainage in the sheep, Indian ink and a casting polymer were injected into the interstitium of the five tonsils present in the heads of slaughtered sheep. This enabled to determine the draining lymph node and to examine the microscopic organization of lymph vessels using light and scanning electron microscopy. No lymph vessels were observed within the tonsillar lymphoid follicles. The corrosion casts demonstrated that the lymphoid follicles are surrounded by numerous sacculated lymph sinuses that drain into a dense interfollicular lymph vessel network. From here the lymph flows into single small lymph vessels that in turn drain into larger lymph vessels extending towards the medial retropharyngeal lymph node. The presented results can be valuable for immunological studies, e.g. during oral or intranasal vaccine development.

Keywords: Sheep; Tonsils; Lymph drainage; Lymph vessels; Corrosion cast

Introduction

Immune responses against ingested and inhaled foreign antigens are initiated in the tonsils since these mucosa-associated lymphoid tissues (MALT) are the body's most proximal line of defence (Casteleyn et al., 2007; Brandtzaeg, 2011; Maunsell et al., 2012). Besides maintaining pharyngeal immunity, locally activated antigen-specific T and B lymphocytes can migrate from the tonsils towards distant MALT sites such as the Peyer's patches (Brandtzaeg and Pabst, 2004; Kuper, 2006). To this purpose, the lymphocytes leave the tonsils via efferent lymph vessels, subsequently drain into the blood circulation and finally enter immunological effector sites by diapedesis through high endothelial venules (HEVs) (Indrasingh et al., 2002; Cesta, 2006; Liebler-Tenorio and Pabst, 2006). As such, lymph vessels and HEVs form the structural basis of the integrated mucosal immune system since they provide the routes for lymphocyte circulation between different MALT sites (Belz and Heath, 1995; Girard and Springer, 1995; Rothkötter et al., 1995; Belz, 1998).

During the past decade the tonsils of sheep have received particular interest within the scope of prion diseases (Schreuder et al., 1998; Van Keulen et al., 2008). To aid the unravelling of the pathogenesis of transmissible spongiform encephalopathies (TSEs), morphological and immunological data on the five ovine tonsils, including the paired palatine tonsil (tonsilla (t.) palatina) located in the oropharynx, the paired paraepiglottic tonsil (t. paraepiglottica) in the laryngopharynx, and the tonsil of the soft palate (t. veli palatina), the pharyngeal tonsil (t. pharyngea) and the paired tubal tonsils (t. tubaria) in the nasopharynx (Casteleyn et al., 2007, 2011a; Breugelmans et al., 2011a), were gathered. The exact anatomical localizations, histological characteristics, volumes and surface areas, distribution of lymphoid cell populations, and innervation of the ovine tonsils have yet been described (Russo et al., 2009; Casteleyn et al., 2007, 2008a, 2010a; Breugelmans et al., 2011a, 2011b, 2011c; Toppets et al., 2012).

Remarkably, only few data are available on the lymphatic system of the ovine tonsils. Indeed, many conventional anatomical handbooks do not mention the lymphatic drainage from the tonsils (Martin and Schauder, 1938; Grau-Karlsbad, 1974; Saar and Getty, 1975; Vollmerhaus, 1976; König et al., 2002; Gille, 2008). Moreover, when literature data are available, they are inconsistent or only general in nature, without mentioning species-specific characteristics. For example, according to Berg (1988) the lymph from the tonsils of the common domestic mammals is conveyed to the mandibular lymph centre, while Barone (1996) states that all tonsillar lymph flows directly to the retropharyngeal lymph centre.

Detailed, species-specific knowledge of the tonsillar lymphatic system can be valuable for immunological studies, e.g. during oral or intranasal vaccine development investigating the generation of antigen-specific lymphocytes (Casteleyn et al., 2013). The usefulness of such mucosal vaccines is illustrated by the successful eradication of human polio by oral vaccination (Murphy et al., 2008). Veterinary examples of potent intranasal vaccination include the vaccines against kennel cough in the dog, bovine and feline rhinotracheitis, *Streptococcus equi* infections in horses, and infectious bronchitis and Newcastle disease in poultry (Tizard, 2013). Besides the induction of mucosal immunity at the sites of potential infection, immune cells activated in the tonsils by mucosal vaccines use the efferent tonsillar lymph vessels to migrate towards the draining lymph node where serum IgG antibodies, providing systemic immunity, are secreted (Kuper et al., 1992; Zuercher, 2003).

In the framework of assessing the effectiveness of a newly developed mucosal vaccine, it is important to identify the lymph nodes draining the ovine tonsils. In line with this, the intratonsillar distribution of the lymph vessels should be investigated to gain insight in the emigration of activated lymphocytes into the tonsillar lymphatic system. In contrast to the well described distribution of lymph vessels within lymph nodes (Heath et al., 1986; Heath and Spalding, 1987; Azzali, 2003), descriptions of the lymphatic systems present in MALT

85 sites are restricted to the human and canine palatine tonsils, the Peyer's patches of the rabbit,
86 guinea pig, mouse, sheep and pig, and the vermiform appendix of the rabbit (Ohtani and
87 Murakami, 1990; Lowden and Heath, 1992, 1994; Belz and Heath, 1995; Regoli et al., 1995;
88 Fujisaka et al., 1996; Azzali, 1998; Azzali et al., 2002; Azzali, 2003).

89 Since data on the lymphatic system of the ovine tonsils are largely lacking and ambiguous,
90 the aims of the present study were to identify the draining lymph node(s) and to describe the
91 microstructure of the tonsillar lymphatic system in the sheep. The research methods included
92 interstitial injection of Indian ink and casting polymer into the five ovine tonsils, combined
93 with light and scanning electron microscopy (SEM).

95 **Materials and Methods**

96 *Lymph drainage from the tonsils*

97 The heads of five healthy one-year-old Texel sheep were collected at a local abattoir. Using a
98 band saw, they were longitudinally sectioned 1 cm lateral to the median plane in order not to
99 transect the pharyngeal tonsil located at the caudal end of the pharyngeal septum (Casteleyn et
100 al., 2007, 2008a). Indian ink (Pelikan black 17 Fount India, Pelikan Benelux N.V., Groot-
101 Bijgaarden, Belgium) was interstitially injected into the centre of each tonsil (ten palatine,
102 paraepiglottic and tubal tonsils, and five pharyngeal tonsils and tonsils of the soft palate) via a
103 26 gauge (G) needle inserted just underneath the tonsillar epithelium (Castenholz, 1984, 1986;
104 Lowden and Heath, 1992, 1993; Casteleyn et al., 2007, 2008b). A volume of 0.5 ml ink was
105 injected into the largest ovine tonsil, i.e. the pharyngeal tonsil (Casteleyn et al., 2007). The
106 second largest tonsil, being the palatine tonsil, was injected with 0.3 ml ink, while all other,
107 small, tonsils received 0.1 ml ink (Casteleyn et al., 2007). Subsequently, the heads were
108 dissected to demonstrate black stained lymph vessels emerging from the injected tonsils and
109 coursing towards either the parotid, mandibular or retropharyngeal lymph nodes. These lymph

nodes were examined both macroscopically and histologically using a light microscope (Olympus BX 61, Olympus Belgium, Aartselaar, Belgium) for the appearance of black staining and the presence of carbon particles, respectively (Casteleyn et al., 2008b). For histological analysis, the lymph nodes were fixated in 3.5% neutral buffered formaldehyde for two days at room temperature. They were then routinely processed to a series of 10 histological sections, cut at fixed intervals throughout the entire lymph nodes that were stained with haematoxylin and eosin (HE).

Light microscopy on the tonsils after interstitial injection of Indian ink

The interstitially injected tonsils were excised and fixated in 3.5% neutral buffered formaldehyde for two days at room temperature. The fixated samples were routinely processed to a series of 10 HE-stained histological sections, cut at fixed intervals throughout the entire tonsils. The presence and distribution of carbon particles were examined by light microscopy (Olympus BX 61).

SEM on corrosion casts of the tonsillar lymphatic system

The heads of an additional number of five healthy one-year-old Texel sheep were collected at a local abattoir. Each tonsil was interstitially injected via a 26 G needle inserted just underneath the tonsillar epithelium with, according to the size of the tonsil as described above, 0.1 ml to 0.5 ml Mercor II[®] (methyl methacrylate-based casting medium, Ladd Research, Williston, USA). After polymerization, which took approximately 15 min, the casted tonsils were excised and macerated overnight in 25% KOH (Carl Roth GmbH, Karlsruhe, Germany). The obtained lymphatic casts were gently rinsed with distilled water, air dried, coated with platinum and examined with SEM (JEOL JSM 5600 LV, Jeol, Zaventem, Belgium).

135

136 **Results**

137 *Lymph drainage from the tonsils*

138 Indian ink that was injected superficially into each of the five ovine tonsils drained directly to
139 the medial retropharyngeal lymph node in all five examined animals. The afferent lymph
140 vessels to this lymph node were easily distinguishable by their black colour (Fig. 1A). At the
141 surface of the lymph node they divided into primary branches, giving off secondary
142 ramifications that in turn divided into tertiary or terminal branches. The latter terminated in
143 the subcapsular sinus of the lymph node (Fig. 1B).

144

145 *Light microscopy on the tonsils after interstitial injection of Indian ink*

146 After interstitial injection of Indian ink into the tonsils, carbon particles were abundantly
147 present in the lymph vessels located in between neighbouring lymphoid follicles (Fig. 2A). As
148 such, an interfollicular lymph vessel network surrounding the lymphoid follicles could be
149 observed. Only limited amounts of carbon particles were seen within the lymphoid follicles
150 (Fig. 2B). These carbon particles seemed not to be enclosed by endothelium-lined vessels, but
151 appeared as free particles lying in the interstitial spaces between the densely packed
152 lymphocytes composing the lymphoid follicles. The interfollicular lymph vessel network
153 drained into larger lymph vessels that were present in the connective tissue underneath the
154 scattered lymphoid follicles located in the tonsil of the soft palate and in the paraepiglottic and
155 tubal tonsils. Larger lymph vessels also coursed through the connective tissue core of the
156 pharyngeal tonsil, and through the connective tissue septa of the palatine tonsil (Fig. 2A).
157 Blood vessels, containing erythrocytes, contrasted well with the lymph vessels by their lack of
158 carbon particles (Fig. 2B).

159

SEM on corrosion casts of tonsillar lymph vessels

At low magnification round impressions were visible at the surfaces of the casts from the palatine, paraepiglottic and pharyngeal tonsils (Figs. 3-5). The diameters of these impressions that are located just underneath the tonsillar epithelium amounted to approximately 500 μm . These data combined with the histological images presented in Fig. 2 demonstrate that the round impressions represent the locations of lymphoid follicles in unmacerated tonsils. No indications for the presence of intraepithelial lymph vessels were noticed on the corrosion casts. At higher magnification sacculated lymph sinuses were observed at the bottom and the sides of the lymphoid follicles. In between these follicles a finely-meshed interfollicular lymph vessel network, characterized by numerous cellular impressions, was present (Fig. 5). Some distinct lymph vessels extended through the interfollicular regions and connected to larger efferent lymph vessels that emerged from the tonsils. Deeply indented, V-shaped constrictions were visible at intervals of approximately 500 μm in larger lymph vessels travelling through the connective tissue cores and septa of tonsils and in those running in the subepithelial connective tissue towards the draining lymph node (Figs. 5 and 6).

Since the tubal tonsils and the tonsils of the soft palate only contain scattered lymphoid follicles, impressions could not be recognized on the surfaces of their casts (Fig. 6) (Casteleyn et al., 2007). The interstitial spaces on the nasopharyngeal side of the soft palate and the lateral nasopharyngeal mucosa were recognized on the casts by their numerous cellular impressions. The interstitium drained into small lymph vessels that merged into larger lymph vessels. These vessels extended in rostrocaudal direction towards the medial retropharyngeal lymph nodes and showed deeply indented, often V-shaped constrictions (Fig. 6).

Discussion

The present study describes the lymph drainage from the t. palatina, t. paraepiglottica, t. pharyngea, t. tubaria and t. veli palatini. Although an additional, lingual tonsil has been described in the sheep (Casteleyn et al., 2007, 2011b), this tonsil was not included since it only consists of small non-organized aggregations of lymphoid cells mainly located in the centres of the vallate papillae at the root of the tongue (Casteleyn et al., 2007; Breugelmans et al., 2011a).

The data presented here are morphological in nature, but aim to contribute to the knowledge of the mucosal immune system as well. In order to correspond to the commonly used immunological terminology, the official anatomical terms defining a well-circumscribed aggregation of lymphoid cells, i.e. lymph nodule, lymphonodulus or nodulus lymphaticus, were replaced by the immunological term lymphoid follicle (Casteleyn et al., 2011a).

During the interstitial injections with Indian ink in the centres of the tonsils it was macroscopically observed that the injected volumes were sufficient to allow spreading of the ink towards the periphery of the tonsils. The lymph, in our experiments represented by ink, drained directly from each tonsil to the medial retropharyngeal lymph node. Since all experiments were performed on slaughtered animals, this draining was due to the manual pressure by which the liquids were injected. It is plausible that swallowing enhances lymph drainage from the ovine tonsils in vivo. In this respect, the V-shaped indentations that were observed on the casted larger lymph vessels, that are suggestive of lymphatic valves (Castenholz, 1986), are meaningful. The application of manual pressure during the injections of ink might have forced carbon particles into the interstitium of lymphoid follicles, as suggested by Fig. 2B. The high cellular density of lymphoid follicles could explain the presence of only limited amounts of intrafollicular carbon particles. These particles could also be present in small intrafollicular lymph vessels, but this was not demonstrated on the corrosion casts. In addition, other authors also state that no lymph vessels are present within

lymphoid follicles (Ohtani and Murakami, 1990; Lowden and Heath, 1992, 1994; Belz and Heath, 1995; Regoli et al., 1995; Fujisaka et al., 1996; Azzali, 1998; Azzali et al., 2002; Azzali, 2003).

For the preparation of the corrosion casts, Mercor[®] was used since its very low viscosity is highly advantageous for interstitial injection and casting of capillary structures (Hodde, 1981; Murakami et al., 1984). Unfortunately, Mercor[®] casts are very brittle and thus have to be handled with care. Another disadvantage is that Mercor[®] is approximately twice as expensive as Batson's[®] (Polysciences, Warrington, USA). The latter casting polymer is also based on methyl methacrylate, but has a much higher viscosity than Mercor[®]. It is therefore commonly used to cast larger blood vessels and other lumen containing structures such as the avian air sacs (Konerding, 1991; Casteleyn et al., 2010b; Stefanov et al., 2013).

The distribution of the injected Mercor[®] was similar to that of the Indian ink. Its low viscosity also enabled the polymer to course towards the medial retropharyngeal lymph nodes. The scanning electron microscopic knowledge obtained in the present study concerning the lymph vessel organization of the ovine tonsils correspond with the data presented by other authors investigating the lymph vessel organization of the human and canine palatine tonsils, the Peyer's patches of the rabbit, guinea pig, mouse, sheep and pig, and the rabbit vermiform appendix (Ohtani and Murakami, 1990; Lowden and Heath, 1992, 1994; Belz and Heath, 1995; Regoli et al., 1995; Fujisaka et al., 1996; Azzali, 1998; Azzali et al., 2002; Azzali, 2003).

The parotid and mandibular lymph nodes did not receive Indian ink. Our observation therefore contrasts with the statement of Berg (1988) that all tonsillar lymph flows to the mandibular lymph centre, but corresponds with the data presented by Barone (1996). These lymph nodes therefore seem to play no role in the induction of immunity after antigen uptake at tonsillar epithelia, e.g. by M cell-like cells (Casteleyn et al., 2010a, 2013). Since tonsils

lack afferent lymph vessels (Mair et al., 1987; Chen et al., 1989; Indrasingh et al., 2002), it is not surprising that lymph vessels connecting different tonsils were not observed. Thus, no lymphatic connection between the different tonsillar structures of Waldeyer's ring exists (Von Waldeyer-Hartz, 1884).

It should be noted that the results presented here have been obtained in Texel sheep. Since this ovine breed is reared for meat production, tissues from Texel sheep are easily collected at the slaughterhouse. This breed is therefore very valuable for ex vivo experiments. To the best of our knowledge, no anatomical studies investigating the lymph drainage from the tonsils in other ovine breeds have yet been performed. As a result, no statement on the existence of breed related anatomical variations can be made. However, it is not plausible that the general pattern of lymph drainage from the tonsils, i.e. collection of interstitial fluid into small lymph vessels that convey into larger lymph vessels flowing towards the medial retropharyngeal lymph node, differs between various ovine breeds.

Immunocompetent cells deriving from those tonsils that are considered as the main inductive sites for mucosal immunity in the sheep, i.e. the palatine and pharyngeal tonsils (Breugelmans et al., 2011b) as well as lymphocytes activated in the medial retropharyngeal lymph node by drained antigens all emigrate together with secreted systemic antibodies produced in the draining lymph node (Kuper et al., 1992; Zuercher, 2003) via efferent lymph vessels towards the ipsilateral lateral retropharyngeal lymph nodes (Schummer et al., 1981; Barone, 1996). From here, the lymph flows in the ipsilateral tracheal trunks that extend towards the deep caudal cervical lymph nodes located at the ventral side of the trachea near the median plane (Barone, 1996). Finally, the lymph from these lymph nodes is conveyed in the common duct of the two tracheal trunks which discharge by a variable pattern in the cranialmost segment of the cranial vena cava at the thoracic inlet (Schummer et al., 1981; Barone, 1996). These anatomical structures thus provide the route for antigen-specific

lymphocytes and antibodies to reach the systemic circulation, enabling them to travel to and protect distant MALT sites.

Conclusions

The tonsillar lymphatic system in the sheep is well-developed. It is composed of interfollicular lymph vessel networks draining into single small lymph vessels that join to form larger efferent lymph vessels containing valves. All efferent lymph vessels from each of the five ovine tonsils drain to the retropharyngeal lymph centre. The other cranial lymph centres, i.e. the parotid and mandibular lymph centres, play no role in the induction of immunity against harmful antigens sampled at the tonsillar surfaces.

Acknowledgements

The authors are grateful to the slaughterhouse Isla-Meat (Lennik, Belgium) for supplying the sheep heads. The technical assistance of P. Vervaet and L. Standaert is kindly acknowledged.

References

- Azzali, G., 1998: Three-dimensional and ultrastructural aspects of the lymphatic vascularization of the vermiform appendix. *J. Submicrosc. Cytol. Pathol.* **30**, 545-553.
- Azzali, G., 2003: Structure, lymphatic vascularization and lymphocyte migration in mucosa-associated lymphoid tissue. *Immunol. Rev.* **195**, 178-189.
- Azzali, G., M. Vitale, and M. L. Arcari, 2002: Ultrastructure of absorbing peripheral lymphatic vessel (ALPA) in Guinea pig Peyer's patches. *Microvasc. Res.* **64**, 289-301.
- Barone, R., 1996: *Anatomie Comparée des Mammifères Domestiques*, Tome Cinquième: Angiologie (R. Barone ed). Paris: Éditions Vigot.

283 Belz, G. T., 1998: An unusual structure of venules in tonsils of the soft palate of young pigs.
 284 J. Anat. **192**, 131-135.

285 Belz, G. T., and T. J Heath, 1995: Lymphatic drainage from the tonsil of the soft palate in
 286 pigs. J. Anat. **187**, 491-495.

287 Berg, R., 1988: Kopf, Caput. In: Angewandte und topographische Anatomie der Haustiere, 3.
 288 Auflage (R. Berg ed). Jena: VEB Gustav Fischer Verlag.

289 Brandtzaeg, P., 2011: Immune functions of nasopharyngeal lymphoid tissue. Adv.
 290 Otorhinolaryngol. **72**, 20-24.

291 Brandtzaeg, P., and R. Pabst, 2004: Let's go mucosal: Communication on slippery ground.
 292 Trends Immunol. **25**, 570-577.

293 Breugelmans, S., C. Casteleyn, P. Simoens, and W. Van den Broeck, 2011a: Distribution of
 294 the lingual lymphoid tissue in domestic ruminants. Anat. Histol. Embryol. **40**, 426-432.

295 Breugelmans, S., W. De Spiegelaere, C. Casteleyn, P. Simoens, and W. Van den Broeck,
 296 2011b: Differences between the ovine tonsils based on an immunohistochemical
 297 quantification of the lymphocyte subpopulations. Comp. Immunol. Microbiol. Infect. Dis.
 298 **34**, 217-225.

299 Breugelmans, S., W. Van den Broeck, K. Demeyere, E. Meyer, and P. Simoens, 2011c:
 300 Immunoassay of lymphocyte subsets in ovine palatine tonsils. Acta Histochem. **113**, 416-
 301 422.

302 Casteleyn C., D. François, P. Simoens, and W. Van den Broeck, 2010b. The avian air sacs:
 303 Visualization in the chicken by means of the corrosion casting technique. Vlaams Diergen.
 304 Tijds. **79**, 429-435.

305 Casteleyn, C. R., S. Breugelmans, P. Simoens, and W. Van den Broeck, 2008b:
 306 Morphological and immunological characteristics of the bovine temporal lymph node and
 307 hemal node. Vet. Immunol. Immunopathol. **126**, 339-350.

308 Casteleyn, C. R., W. Van den Broeck, and P. Simoens, 2007: Histological characteristics and
 309 stereological volume assessment of the ovine tonsils. *Vet. Immunol. Immunopathol.* **120**,
 310 124-135.

311 Casteleyn, C., M. Cornelissen, P. Simoens, and W. Van den Broeck, 2010a: Ultramicroscopic
 312 examination of the ovine tonsillar epithelia. *Anat. Rec.* **293**, 879-889.

313 Casteleyn, C., P. Cornillie, P. Simoens, and W. Van den Broeck, 2008a: Stereological
 314 assessment of the epithelial surface area of the ovine palatine and pharyngeal tonsils. *Anat.*
 315 *Histol. Embryol.* **37**, 366-368.

316 Casteleyn, C., P. Simoens, and W. Van den Broeck, 2011a: Terminology of the tonsils. *Anat.*
 317 *Histol. Embryol.* **40**, 204-209.

318 Casteleyn, C., S. Breugelmans, P. Simoens, and W. Van den Broeck, 2011b: The tonsils
 319 revisited: Review of the anatomical localization and histological characteristics of the
 320 tonsils of domestic and laboratory animals. *Clin. Dev. Immunol.* (open access journal; doi:
 321 10.1155/2011/472460).

322 Casteleyn, C., W. Van den Broeck, A. Gebert, B. Tambuyzer, S. Van Cruchten, and C. Van
 323 Ginneken, 2013: M cell specific markers in man and domestic animals: Valuable tools in
 324 vaccine development. *Comp. Immunol. Microbiol. Infect. Dis.* **36**, 353-364.

325 Castenholz, A., 1984: Morphological characteristics of initial lymphatics in the tongue as
 326 shown by scanning electron microscopy. *Scan. Electron Microsc.* **1984/III**, 1343-1352.

327 Castenholz, A., 1986: Corrosion cast technique applied in lymphatic pathways. *Scan. Electron*
 328 *Microsc.* **1986/II**, 599-605.

329 Cesta, M. F., 2006: Normal structure, function, and histology of mucosa-associated lymphoid
 330 tissue. *Toxicol. Pathol.* **34**, 599-608.

331 Chen, W., M. R. Alley, and B. W. Manktelow, 1989: Respiratory tract-associated lymphoid
 332 tissue in conventionally raised sheep. *J. Comp. Pathol.* **101**, 327-340.

333 Fujisaka, M., O. Ohtani, and Y. Watanabe, 1996: Distribution of lymphatics in human
 334 palatine tonsils: A study by enzyme-histochemistry and scanning electron microscopy on
 335 lymphatic corrosion casts. *Arch. Histol. Cytol.* **59**, 273-280.

336 Gille, U., 2008: Herz, Kreislauf- und Abwehrsystem, Angiologia. In: *Anatomie für die*
 337 *Tiermedizin*, 2. Auflage (F.-V. Salomon, H. Geyer and U. Gille eds). Stuttgart: Enke
 338 Verlag.

339 Girard, J.-P., and T. A. Springer, 1995: High endothelial venules (HEV's): specialized
 340 endothelium for lymphocyte migration. *Immunol. Today* **16**, 449-457.

341 Grau-Karlsbad, H., 1974: Das Lymphgefäßsystem. In: *Ellenberger/Baum: Handbuch der*
 342 *vergleichenden Anatomie der Haustiere*, 18. Auflage (O. Zietzschmann, E. Ackerknecht
 343 and H. Grau eds). Berlin: Springer-Verlag.

344 Heath, T. J., and H. J. Spalding, 1987: Pathways of lymph flow to and from the medulla of
 345 lymph nodes in sheep. *J. Anat.* **155**, 177-188.

346 Heath, T. J., R. L. Kerlin, and H. J. Spalding, 1986: Afferent pathways of lymph flow within
 347 the popliteal node in sheep. *J. Anat.* **149**, 65-75.

348 Hodde, K.C., 1981: Cephalic vascular patterns in the rat: A scanning electron microscopic
 349 (SEM) study of casts. PhD dissertation, University of Amsterdam.

350 Indrasingh, I., G. Chandi, and S. Vettivel, 2002: Route of lymphocyte migration through the
 351 high endothelial venule (HEV) in human palatine tonsil. *Ann. Anat.* **184**, 77-84.

352 Konerding, M. A., 1991: Scanning electron microscopy of corrosion casting in medicine.
 353 *Scanning Microsc.* **5**, 851-865.

354 König, H. E., J. Sautet, and H.-G. Liebich, 2002: Verdauungsapparat (Apparatus digestorius).
 355 In: *Anatomie der Haussäugetiere – Lehrbuch und Farbatlas für Studium und Praxis*, Band
 356 *II Organe, Kreislauf- und Nervensystem* (H. E. König and H.-G. Liebig eds). Stuttgart:
 357 Schattauer.

358 Kuper, C. F., P. J. Koornstra, D. M. H. Hamelers, J. Biewenga, B. J. Spit, A. M. Duijvestijn,
 359 P. J. C. van Breda Vriesman, and T. Sminia, 1992: The role of nasopharyngeal lymphoid
 360 tissue. *Immunol. Today* **13**, 219-224.
 361 Kuper, C.F., 2006: Histopathology of mucosa-associated lymphoid tissue. *Toxicol. Pathol.* **34**,
 362 609-615.
 363 Liebler-Tenorio, E., and R. Pabst, 2006: MALT structure and function in farm animals. *Vet.*
 364 *Res.* **37**, 257-280.
 365 Lowden, S., and T. Heath, 1992: Lymph pathways associated with Peyer's patches in sheep. *J.*
 366 *Anat.* **181**, 209-217.
 367 Lowden, S., and T. Heath, 1993: Lymphatic drainage from the distal small intestine in sheep.
 368 *J. Anat.* **183**, 13-20.
 369 Lowden, S., and T. Heath, 1994: Ileal Peyer's patches in pigs: intercellular and lymphatic
 370 pathways. *Anat. Rec.* **239**, 297-305.
 371 Mair, T. S., E. H. Batten, C. R. Stokes, and F. J. Bourne, 1987: The histological features of
 372 the immune system of the equine respiratory tract. *J. Comp. Pathol.* **97**, 575-586.
 373 Martin, P., and W. Schauder, 1938: Lymphgefäßsystem des Rindes. In: *Lehrbuch der*
 374 *Anatomie der Haustiere, III. Band, III. Teil: Harn- und Geschlechtsorgane, Blut- und*
 375 *Lymphgefäßsystem, Nervensystem, Haut- und Sinnesorgane der Hauswiederkäuer* (P.
 376 Martin and W. Schauder eds). Stuttgart: Verlag von Schickhardt & Ebner.
 377 Maunsell, F., M. B. Brown, J. Powe, J. Ivey, M. Woolard, W. Love, and J. W. Simecka, 2012:
 378 Oral inoculation of young dairy calves with *Mycoplasma bovis* results in colonization of
 379 tonsils, development of otitis media and local immunity. *PLoS One* **7**, e44523.
 380 Murakami, T., T. Itoshima, K. Hitomi, A. Ohtsuka, and A. L. Jones, 1984: A monomeric
 381 methyl and hydroxypropyl methacrylate injection medium and its utility in casting blood

382 capillaries and liver bile canaliculi for scanning electron microscopy. Arch. Histol. Jpn.
383 **47**, 223-37.

384 Murphy, K., P. Travers, and M. Walport, 2008: Manipulation of the immune response. In:
385 Janeway's Immunobiology, 7th edn. (K. Murphy, P. Travers and M. Walport eds).
386 London: Garland Science.

387 Ohtani, O., and T. Murakami, 1990: Organization of the lymphatic vessels and their
388 relationships to the blood vessels in the rabbit Peyer's patches. Arch. Histol. Cytol. **53**,
389 155-164.

390 Regoli, M., C. Borghes, E. Bertelli, and L. Comparini, 1995: Arrangement of the small
391 intestine lymphatic network in the Peyer's patches of the mouse. A light and transmission
392 electron microscopic study. Ann. Anat. **177**, 229-235.

393 Rothkötter, H. J., C. Hriesi, and R. Pabst, 1995: More newly formed T than B lymphocytes
394 leave the intestinal mucosa via lymphatics. Eur. J. Immunol. **25**, 866-869.

395 Russo, D., C. Mongardi Fantaguzzi, G. Di Guardo, P. Clavenzani, G. Lalatta Costerbosa, C.
396 Ligios, and R. Chiocchetti, 2009: Characterization of sheep (*Ovis aries*) palatine tonsil
397 innervation. Neuroscience **161**, 813-826.

398 Saar, L. I., and R. Getty, 1975: Ruminant lymphatic system: part II Ovine. In: Sisson and
399 Grossman's: The Anatomy of the Domestic Animals, 5th edn. (R. Getty ed). London: W.
400 B. Saunders Company.

401 Schreuder, B. E. C., L. J. M. van Keulen, M. E. W. Vromans, J. P. M. Langeveld, and M. A.
402 Smits, 1998: Tonsillar biopsy and PrP^{Sc} detection in the preclinical diagnosis of scrapie.
403 Vet. Rec. **142**, 564-568.

404 Schummer, A., H. Wilkens, B. Vollmerhaus, and K.-H. Habermehl, 1981: The Circulatory
405 System, the Skin, and the Cutaneous Organs of the Domestic Mammals. In: The Anatomy

of the Domestic Animals (R. Nickel, A. Schummer and E. Seiferle eds). Berlin: Verlag Paul Parey.

Stefanov, M., J. D. Kim, M. H. Nam, and Soh K. S., 2013: New approach of corrosion casting using direct injection of mercox into the parenchyma of different organs. *Anat. Rec. (Hoboken)* **296**, 724-725.

Tizard, I. R., 2013: The use of vaccines. In: *Veterinary Immunology*, 9th edn. (I. R. Tizard ed). St. Louis: Elsevier.

Toppets, V., J. Piret, N. Kirschvink, F. Lantier, I. Lantier, P. Berthon, G. Daube, L. Massart, L. Grobet, and N. Antoine, 2012: Neuroimmune connections in ovine pharyngeal tonsil: potential site for prion neuroinvasion. *Cell Tissue Res.* **348**, 167-176.

van Keulen, L. J. M., A. Bossers, and F. Van Zijderveld, 2008: TSE pathogenesis in cattle and sheep. *Vet. Res.* **39**, 24-35.

Vollmerhaus, B., 1976: Lymphatisches System. In: *Lehrbuch der Anatomie der Haustiere*, Band III Kreislaufsystem, Haut und Hautorgane (R. Nickel, A. Schummer and E. Seiferle eds). Berlin: Verlag Paul Parey.

von Waldeyer-Hartz, W., 1884: Ueber den lymphatischen Apparat des Pharynx. *Dtsch. Med. Wochenschr.* 10, 313.

Zuercher, A. W., 2003: Upper respiratory tract immunity. *Viral Immunol.* **16**, 279-289.

Figure legends

Fig. 1. Paramedian section through the ovine head showing a medial view of the right side of the nasopharynx. After Indian ink was interstitially injected into the tubal tonsil (1), black carbon particles spread through lymph vessels (arrows) that extended towards the medial retropharyngeal lymph node (2). Some anatomical landmarks, such as the soft palate (3) and the epiglottis (4) are additionally indicated. B: Medial retropharyngeal lymph node receiving

carbon particles via three afferent lymph vessels (alv) giving off primary branches (pb) that subsequently divide into secondary branches (sb) which finally give off terminal branches (tb) that enter the subcapsular sinus (ss).

Fig. 2. Histological section through the ovine pharyngeal tonsil after interstitial injection of Indian ink. A: Carbon particles are visible in the interfollicular lymph vessel network (iln) which surrounds the lymphoid follicles (F). Larger lymph vessels (ILV) are present in the connective tissue core of the tonsil. B: Higher magnification showing the abundant presence of carbon particles in the interfollicular lymph vessel network (iln) and lymph sinuses surrounding the lymphoid follicles (F). The interfollicular blood vessels (arrows) are not loaded with carbon particles. Notice the intrafollicular presence of carbon particles that are not enclosed by endothelium-lined vessels (arrowheads).

Fig. 3. SEM views of the lymph vessel architecture of the ovine pharyngeal tonsil. A: Spaces previously occupied by lymphoid follicles (F) are surrounded by numerous sacculated lymph sinuses (sl) that drain into interfollicular lymph vessel networks (iln). Small lymph vessels (sLV) emerge from this network. B: Higher magnification of the interfollicular lymph vessel network (iln) showing the imprints of lymphoid cells (arrows) present in the interfollicular areas. C: The small lymph vessels (sLV) that emerge from the interfollicular lymph vessel network (iln) surrounding the lymphoid follicles (F) drain into large lymph vessels that travel through the connective tissue core of the tonsil. D: A larger lymph vessel (ILV) characterized by the presence of many, often V-shaped, constrictions (arrows) corresponding to lymph vessel valves courses through interstitial spaces (i).

Fig. 4. SEM images of the lymph vessel architecture of the ovine palatine tonsil. A: The locations of macerated lymphoid follicles (F) can be recognized as round impressions. Each lymphoid follicle is surrounded by the interfollicular lymph vessel network (iln). Some lymph vessels (LV) are additionally indicated. B: Higher magnification of the impression of the lymphoid follicle encircled in A. Sacculated lymph sinuses (sl) surrounding the lymphoid follicle are prominent. They drain into the interfollicular lymph vessel network (iln).

Fig. 5. SEM view of the lymph vessel architecture of the ovine paraepiglottic tonsil. Sacculated lymph sinuses (sl) surrounding the lymphoid follicles (F) drain into the interfollicular lymph vessel network (iln) which is connected by small calibre lymph vessels (sLV) to larger lymph vessels (ILV).

Fig. 6. SEM images of the lymph vessel architecture of the ovine tonsil of the soft palate. A: No imprints of lymphoid follicles can be seen in this diffusely organized tonsil. The subepithelial interstitial spaces (i) drain into small lymph vessels (sLV), which in turn flow into large lymph vessels (ILV) that travel through the nasopharyngeal mucosa. B: Higher magnification of the subepithelial interstitial spaces (i) that drain into a small lymph vessel (sLV).

473 **Figures**

474
475 Fig. 1
476

477
478 Fig. 2
479

Fig. 3

Fig. 4

Fig. 5

Fig. 6