

Preoperative chemoradiation versus radiation alone for stage II and III resectable rectal cancer (Review)

Ceelen WP, Van Nieuwenhove Y, Fierens K



**THE COCHRANE
COLLABORATION®**

This is a reprint of a Cochrane review, prepared and maintained by The Cochrane Collaboration and published in *The Cochrane Library* 2009, Issue 1

<http://www.thecochranelibrary.com>



Preoperative chemoradiation versus radiation alone for stage II and III resectable rectal cancer (Review)
Copyright © 2009 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
BACKGROUND	2
OBJECTIVES	3
METHODS	3
RESULTS	4
DISCUSSION	6
AUTHORS' CONCLUSIONS	7
ACKNOWLEDGEMENTS	8
REFERENCES	8
CHARACTERISTICS OF STUDIES	10
DATA AND ANALYSES	15
Analysis 1.1. Comparison 1 radiotherapy vs radiochemotherapy, Outcome 1 Overall Survival at 5y.	15
Analysis 1.2. Comparison 1 radiotherapy vs radiochemotherapy, Outcome 2 Disease free survival at 5 y.	16
Analysis 1.3. Comparison 1 radiotherapy vs radiochemotherapy, Outcome 3 Mortality 30 d.	16
Analysis 1.4. Comparison 1 radiotherapy vs radiochemotherapy, Outcome 4 Postop morbidity.	17
Analysis 1.5. Comparison 1 radiotherapy vs radiochemotherapy, Outcome 5 Grade III - IV toxicity.	17
Analysis 1.6. Comparison 1 radiotherapy vs radiochemotherapy, Outcome 6 Sphincter preservation.	18
Analysis 1.7. Comparison 1 radiotherapy vs radiochemotherapy, Outcome 7 pCR.	18
Analysis 1.8. Comparison 1 radiotherapy vs radiochemotherapy, Outcome 8 Anastomotic leak.	19
Analysis 1.9. Comparison 1 radiotherapy vs radiochemotherapy, Outcome 9 Local Recurrence at 5y.	19
WHAT'S NEW	19
HISTORY	20
CONTRIBUTIONS OF AUTHORS	20
DECLARATIONS OF INTEREST	20
INDEX TERMS	20

[Intervention Review]

Preoperative chemoradiation versus radiation alone for stage II and III resectable rectal cancer

Wim P Ceelen¹, Yves Van Nieuwenhove², Kjell Fierens³

¹Dept. of Surgery 2K12 1C, Ghent University Hospital, Ghent, Belgium. ²Department of Experimental Surgery, Free University of Brussels, Brussels, Belgium. ³Dept. of Surgery 2K12 1c, Ghent University Hospital, Ghent, Belgium

Contact address: Wim P Ceelen, Dept. of Surgery 2K12 1C, Ghent University Hospital, De Pintelaan 185, Ghent, B-9000, Belgium. wim.ceelen@ugent.be.

Editorial group: Cochrane Colorectal Cancer Group.

Publication status and date: New, published in Issue 1, 2009.

Review content assessed as up-to-date: 1 September 2008.

Citation: Ceelen WP, Van Nieuwenhove Y, Fierens K. Preoperative chemoradiation versus radiation alone for stage II and III resectable rectal cancer. *Cochrane Database of Systematic Reviews* 2009, Issue 1. Art. No.: CD006041. DOI: 10.1002/14651858.CD006041.pub2.

Copyright © 2009 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

Preoperative radiotherapy (RT) decreases local recurrence rate and improves survival in stage II and III rectal cancer patients. The combination of chemotherapy with RT has a sound radiobiological rationale, and phase II trials of combined chemoradiation (CRT) have shown promising activity in rectal cancer.

Objectives

To compare preoperative RT with preoperative CRT in patients with resectable stage II and III rectal cancer.

Search strategy

We searched the Cochrane Central Register of Controlled Trials, Web of Science, Embase.com, and Pubmed from 1975 until June 2007. A manual search was performed of *Ann Surg*, *Arch Surg*, *Cancer*, *J Clin Oncol*, *Int J Radiat Oncol Biol Phys* and the proceedings of ASTRO, ECCO and ASCO from 1990 until June 2007.

Selection criteria

Relevant studies randomized resectable stage II or III rectal cancer patients to at least one arm of preoperative RT alone or at least one arm of preoperative CRT.

Data collection and analysis

Primary outcome parameters included overall survival (OS) at 5 years and local recurrence (LR) rate at 5 years. Secondary outcome parameters included disease free survival (DFS) at 5 years, metastasis rate, pathological complete response rate, clinical response rate, sphincter preservation rate, acute toxicity, postoperative mortality and morbidity, and anastomotic leak rate. Outcome parameters were summarized using the Odds Ratio (OR) and associated 95% confidence interval (CI) using the fixed effects model.

Main results

Four trials were identified and included in the meta-analysis. The addition of chemotherapy to preoperative RT significantly increased grade III and IV acute toxicity (OR 1.68-10, P = 0.002) while no differences were observed in postoperative morbidity or mortality. Compared to preoperative RT alone, preoperative CRT significantly increased the rate of complete pathological response (OR 2.52-

Preoperative chemoradiation versus radiation alone for stage II and III resectable rectal cancer (Review)

Copyright © 2009 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

1

5.27, $P < 0.001$) although this did not translate into a higher sphincter preservation rate (OR 0.92-1.31, $P = 0.29$). The incidence of local recurrence at five years was significantly lower in the CRT group compared to RT alone (OR 0.39-0.72, $P < 0.001$). No statistically significant differences were observed in DFS (OR 0.92-1.34, $P = 0.27$) or OS (OR 0.79-1.14, $P = 0.58$) at five years.

Authors' conclusions

Compared to preoperative RT alone, preoperative CRT enhances pathological response and improves local control in resectable stage II and III rectal cancer, but does not benefit disease free or overall survival. The effects of preoperative CRT on functional outcome and quality of life are incompletely understood and should be addressed in future trials.

PLAIN LANGUAGE SUMMARY

Preoperative chemoradiation versus radiation alone for stage II and III resectable rectal cancer

In rectal cancer patients, local control is improved by administering radiotherapy (RT) before surgery. Recently, studies have combined preoperative RT with chemotherapy aiming to further improve local control. This review compared preoperative RT with preoperative chemoradiation (CRT) in patients with stage II and III rectal cancer.

Based on the combined analysis of four randomized studies, we found that compared to preoperative RT alone, preoperative CRT enhances pathological response and improves local control in resectable stage II and III rectal cancer, but does not benefit disease free or overall survival. Combined CRT enhances treatment related acute toxicity, but does not affect postoperative complication rate. The effects of preoperative CRT on functional outcome and quality of life are incompletely understood and should be addressed in future trials.

BACKGROUND

The incidence of fatal cases of colorectal cancer in Europe exceeds 200,000 per year. Due to the specific anatomy and biology of rectal cancer, surgery alone historically has been associated with local recurrence in up to one in four patients. Locally recurrent disease is usually incurable, causes important morbidity and suffering and gives rise to systemic metastases. In the last few decades, improvements in surgical technique have dramatically lowered the incidence of locally recurrent disease. Careful pathological studies have clearly demonstrated that the major cause of local recurrence is the persistence of tumor foci within the mesorectum (Quirke 1986; Quirke 2003). Intact removal of the entire mesorectum (total mesorectal excision or TME) in cancers of the mid or lower third of the rectum was pioneered by Heald and has resulted in local recurrence rates lower than 5-10% (Heald 1982; Heald 1998; Enker 1999). The importance of complete removal of the surrounding mesorectum necessitates precise preoperative evaluation of the circumferential resection margin using imaging. Recently, magnetic resonance imaging (MRI) using a phased array coil has emerged as the imaging modality of choice in the preoperative evaluation of locally advanced rectal cancer (Beets-Tan 2003; Beets-

Tan 2005; Brown 2004; Daniels 2005; Brown 2006).

Parallel to improvements in surgical technique, adjuvant therapy regimens have been tested in clinical trials in an effort to reduce local recurrence rates. Neoadjuvant radiotherapy (RT) has been shown to significantly decrease local recurrence rate and improve survival provided a biologically equivalent dose (BED) of at least 30 Gy is administered (Gray 2001). The advantages of preoperative over postoperative RT include enhanced effectiveness in well oxygenated tissue, downstaging of advanced tumors and better treatment compliance (Glimelius 2002). This theoretical superiority of the preoperative approach over postoperative adjuvant therapy has been confirmed in the recent German rectal cancer trial (Sauer 2004). The effect of preoperative RT on local recurrence rate is consistent even when optimal surgical technique (TME) is implemented. This was demonstrated by the results of the Dutch rectal cancer trial which randomized rectal cancer patients to undergo either RT followed by TME or TME alone in the setting of a national surgical training programme (Kapiteijn 2001). Compared to TME alone, 5x5 Gray (Gy) of RT followed by TME resulted in a significantly lower local recurrence rate, although no

improvement in overall survival (OS) was noted.

Although preoperative RT results in a significant pathological response in a minority of patients, significant downsizing is rarely achieved using short schedule RT regimens. In order to improve tumor response, preoperative RT has been combined with chemotherapeutic regimens. There is a strong radiobiological rationale to combine RT with chemotherapy. Combined chemoradiation (CRT) for rectal cancer was introduced in the adjuvant setting and subsequently in irresectable disease, where significant downsizing and downstaging was observed in many patients (Minsky 1993; Minsky 1997). The argument for preoperative CRT in resectable disease is based primarily on possible downsizing and downstaging of tumors close to the circumferential resection margin or the sphincter apparatus, thereby enhancing both R0 resection and the sphincter preservation rate. The paramount importance of performing the resection with a negative CRM was shown in several clinical studies (Nagtegaal 2002). Secondly, the addition of chemotherapy could eliminate microscopic systemic disease present at the time of surgery. Possible concerns of preoperative CRT include an increase of both local and systemic toxicity and overtreatment of inaccurately staged patients (Ammann 2003). Several phase I and II studies using preoperative CRT have shown a promising tumor response with acceptable toxicity (Rodel 2003; Osti 2004). A limited number of prospective randomized trials comparing preoperative RT alone with preoperative CRT in resectable rectal cancer are published or ongoing.

OBJECTIVES

To compare preoperative RT with preoperative CRT in patients with resectable stage II or III rectal cancer.

METHODS

Criteria for considering studies for this review

Types of studies

Randomized controlled trials (RCT's) which randomize patients before surgery with curative intent to one of at least two schedules of preoperative therapy including RT and CRT.

Types of participants

Patients with clinical stage II or III resectable rectal cancer undergoing preoperative RT or CRT followed by surgery.

Types of interventions

Preoperative RT or CRT using fractionated external radiotherapy followed by surgery with curative intent (resectable rectal cancer). The surgical procedure must consist of rectal amputation or sphincter preserving anterior resection using an open or laparoscopic approach; local excisions are excluded.

Types of outcome measures

Primary

-local recurrence rate at 5 years

Secondary

-overall survival at 5 years

-disease free survival at 5 years

-systemic metastasis rate

-pathological complete response rate

-clinical response rate

-sphincter preservation rate

- postoperative mortality within 30 days

- postoperative morbidity

- anastomotic leak rate

Search methods for identification of studies

See: Colorectal Cancer Group methods used in reviews.

We searched the following electronic databases

-Cochrane Central Register of Controlled Trials

-ISI Web of Science (Science Citation Index, Current Contents) from 1975 until June 2007

-Embase.com

-Pubmed

Electronic database searches were performed with MeSH terms and free text terms:

-MeSH: \Rectal Neoplasms"[MeSH] AND

Radiotherapy"[MeSH] AND \Drug Therapy"[MeSH]

-Free text terms:

rectal, rectum, cancer, adenocarcinoma, neoplasm, radiotherapy, irradiation, chemotherapy, chemoradiation, radiochemotherapy, combined modality, multimodal, preoperative, neoadjuvant
Manual search/abstract search

- Journals from 1990: Ann Surg, Arch Surg, Cancer, J Clin Oncol, Int J Radiat Oncol Biol Phys

- Proceedings from ASTRO, ECCO and ASCO

No language constraints were applied.

Data collection and analysis

All three reviewers (WPC, KE, YVN) obtained the full text of all relevant studies and were assessed for methodological quality according to the Cochrane Collaboration's tool for assessing the risk of bias (Higgins 2008). Methodological details relevant for

potential bias included sequence generation, allocation concealment, blinding, incomplete outcome data, and selective outcome reporting. Disagreement was resolved by consensus.

Data were extracted by one reviewer (KF) on custom designed forms and entered in a computer database for transfer and statistical analysis in the Review Manager software. The data extracted included first author, year of publication, source, method of preoperative therapy and surgery, method of randomization, number of patients included, randomized, and analyzed, and outcome parameters as listed above. Data accuracy was verified by the senior author (WPC).

RT dose was converted to the biologically equivalent dose (BED) using the linear quadratic equivalent formula (Dale 2005): $BED = nd(1+1/(\alpha/\beta)-(\gamma/\alpha)(T-T_k)$, with n = number of fractions, d = dose per fraction, α/β = the linear quadratic quotient (set at 10 Gy), γ/α = repair rate (set at 0.6 Gy/d), and T_k = the initial time delay in days (set at 7). Differences between categorical outcome parameters were quantified using the odds ratio (OR) and corresponding 95% confidence interval (95CI). Summary statistics were calculated using the Mantel-Haenszel methods. Heterogeneity analysis was performed using the Q test, with significance accepted when $P < 0.1$. When present, heterogeneity will be addressed as recommended by the Cochrane Collaboration Handbook (Higgins 2008).

RESULTS

Description of studies

See: [Characteristics of included studies](#).

Of a total of 17925 studies resulting from the primary search, 324 papers were selected for full review. In all, 320 papers were discarded (Table 1). Four randomized trials were identified comparing preoperative RT with preoperative CRT in resectable stage II or III rectal cancer (Bosset 2006; Boulis-Wassif 1984; Bujko 2006; Gerard 2006).

Table 1. excluded studies

Type	N
Non randomized trials	144
Adjuvant therapy trials	28
Trials not including at least one chemotherapy arm combined with radiotherapy	71

Table 1. excluded studies (Continued)

Trials not including radiotherapy	27
Trials using local or no resection	18
Trials including other tumor types	25
Trials not including stage II/III cancer	7
Total	320

From November 1972 through April 1976, Boulis-Wassif et al. recruited 247 patients with histologically proven localized adenocarcinoma of the rectum and no clinical or surgical evidence of distant metastases. All patients in both groups received preoperative RT by two parallel opposing diamond and chimney fields. All patients received a total dose of 34.5 Gy in 15 fractions of 2.3 Gy each over a total treatment time of 18 days (BED = 35.8 Gy). In the preoperative CRT group, intravenous 5-FU injection (375 mg/m²) was administered during the first 4 days of irradiation. Surgery usually followed within 2 weeks after the last irradiation. Two patients died before surgery. Assessed outcomes included ease of surgery, type of operation, radical resectability rate, histopathological response, postoperative mortality, postoperative period of hospitalization, local control of the disease, distant metastases, disease free survival, and median survival. Follow-up was available up to 7 years.

From April 1999 until February 2002, Bujko et al. included 316 patients with resectable T3-T4 rectal carcinoma without sphincter infiltration and with a lesion accessible to digital rectal examination. Patients were randomised to either preoperative 5 x 5 Gy short-term RT (BED = 37.5 Gy) with subsequent total mesorectal excision (TME) performed within 7 days or to CRT to a total dose of 50.4 Gy (1.8 Gy per fraction during 25 days; BED = 42.2 Gy) concomitantly with two courses of bolus 5-fluorouracil (325 mg/m²) and leucovorin during weeks 1 and 5 of RT. Chemoradiation was followed by TME after 4-6 weeks. Three patients did not undergo surgery. Assessed outcomes were acute postirradiation toxicity, sphincter preservation rate, postoperative mortality, pathology, overall survival, disease free survival, local recurrence rate, distant metastases, late toxicity and incidence of permanent stoma. Median follow up was 48 months.

Between April 1993 and November 2003, Gerard et al. recruited 762 patients with a histologically confirmed, previously untreated rectal adenocarcinoma accessible to digital rectal examination (T3

or resectable T4 tumor with no evidence of distant metastases). Patients were allocated to two treatment arms: preoperative RT vs. preoperative CRT, both followed by surgery. RT was delivered with photons from a linear accelerator in a three- or four-field box technique. The dose per fraction was 1.8 Gy and all fields were treated each day with five fractions per week. The total dose was 45 Gy in 25 fractions during 5 weeks (BED = 42.2 Gy). Concurrent chemotherapy (CT) consisted of bolus 5-fluorouracil (350 mg/m²) and leucovorin administered during week 1 and 5 of RT. Surgery was planned between 3 and 10 weeks after the end of the preoperative RT (+/- CT). TME was recommended. Assessed outcomes were surgical procedures and postoperative complications, pathology, overall survival, progression free survival and local recurrence. Median follow-up was 81 months.

Between April 1993 and March 2003, Bosset et al. recruited 1011 patients with a T3 or resectable T4 M0 adenocarcinoma of the rectum within 15 cm from the anal margin and without previous treatment for this disease (EORTC 22921 trial). Patients were allocated to four treatment arms: preoperative RT, preoperative CRT, preoperative RT plus postoperative CT and preoperative CRT plus postoperative CT. RT consisted of 45 Gy delivered to the posterior pelvis in 25 fractions of 1.8 Gy over a period of 5 weeks (BED = 42.2 Gy). The target volume of RT was not a classical pelvic volume but was limited to the main field of tumor spread and to the perirectal nodes. Preoperative chemotherapy was delivered in two 5-day courses of 5-fluorouracil (350 mg/m²) with leucovorin during the first and fifth weeks of RT. Surgery was scheduled to take place 3 to 10 weeks after treatment. TME was recommended beginning in 1999. Assessed outcomes were toxicity of the preoperative treatment, performed surgical procedures, rate of postoperative complications, pathology, late side effects, overall survival, disease free survival, local and distant recurrence rate. Median follow-up of 5.4 years.

Risk of bias in included studies

Randomisation was adequately performed in all four studies using communication with a central office. Three studies based randomisation on the minimization method (Bujko 2006; Gerard 2006; Bosset 2006). In the fourth study, the randomisation method is not specified (Boulis-Wassif 1984). None of the studies were described as double blind or used blinded outcome assessment. Description of withdrawals and dropouts was given in all four studies. There were no imbalances between treatment arms in the number of patients that did not undergo the complete trial procedure. Three studies were performed on an intention-to-treat basis (Bujko 2006; Gerard 2006; Bosset 2006); no imbalances were identified between treatment arms.

Effects of interventions

The main primary outcome parameter was the local recurrence rate at five years, which was reported in three studies (Bosset 2006; Boulis-Wassif 1984; Gerard 2006). In the RT group, 122 of 740 patients (16.5%) developed a local recurrence while in the CRT group this event was observed in 71 out of 754 patients (9.4%). This difference was statistically significant (OR 0.53, 95%CI 0.39-0.72, $P < 0.001$). No statistically significant heterogeneity among studies was present ($P = 0.12$). Survival data at 5 years were available in three studies (Bosset 2006; Boulis-Wassif 1984; Gerard 2006). In the CRT group, 644 of 1007 patients (63.9%) were alive at 5 years while in the RT group 647 of 993 patients (65.2%) survived 5 years. This difference did not reach statistical significance (OR 0.95, 95%CI 0.79-1.14, $P = 0.58$). No heterogeneity was present ($P = 0.15$).

The results of the analysis of the secondary outcome parameters were as follows. Disease free survival at 5 years, available in the studies of Bosset 2006 and Gerard 2006, was 507/881 (57.5%) in the CRT group and 479/872 (54.9%) in the RT group. This difference was not statistically significant (OR 1.11, 95%CI 0.92-1.34, $P = 0.27$). Significant heterogeneity did not occur ($P = 0.64$). Grade III or IV treatment related toxicity developed in 151 of 1015 patients (14.9%) treated with CRT while in patients treated with RT alone, this occurred in 52 of 1017 patients (5.1%). This difference was statistically significant (OR 4.1, 95%CI 1.68-10, $P = 0.002$). There was, however, significant heterogeneity ($P = 0.005$) which remained when the data were reanalysed using the random effects assumption. Among patients who underwent surgery, sphincter preservation was possible in 551 of 1111 patients (49.6%) in the CRT group and in 527 of 1108 patients (47.6%) in the RT group; this difference failed to reach statistical significance (OR 1.1, 95%CI 0.92-1.31, $P = 0.29$). No heterogeneity was observed ($P = 0.48$). Postoperative 30 day mortality was observed in 31 of 1122 (2.8%) patients in the CRT group and in 21 of 1117 (1.9%) patients in the RT group. This difference did not reach statistical significance (OR 1.48, 95%CI 0.84-2.6, $P = 0.17$); no heterogeneity was detected ($P = 0.6$). Similarly, no

significant differences were observed in postoperative morbidity or anastomotic leak rate (OR 0.68-1.03, $P = 0.1$ and OR 0.57-1.85, $P = 0.93$ respectively). Pathological complete response (ie, ypT0N0) of the resected specimen was observed in 129 of 1096 patients (11.8%) in the CRT group and in 39 of 1105 patients (3.5%) in the RT group. This difference was statistically significant (OR 3.65, 95%CI 2.52-5.27, $P < 0.001$) while significant heterogeneity for this parameter was not observed ($P = 0.15$).

Because of the limited number of included studies, no sensitivity analysis was performed.

DISCUSSION

Preoperative RT has been shown to reduce local recurrence rates and marginally improve survival over surgery alone provided a BED > 30 Gy is delivered to the target region (Gray 2001). The current review addresses the question whether the addition of chemotherapy to preoperative RT further enhances pathological and clinical outcome parameters. Four randomized trials were identified comparing preoperative CRT with preoperative RT alone in resectable, locally advanced rectal cancer. Although there was considerable variation in radiotherapy dose and fractionation, all four studies have used a BED > 30 Gy. In three trials (Gerard 2006; Boulis-Wassif 1984; Bosset 2006), RT regimens were identical in both groups. In the Polish study (Bujko 2006), RT dose and fractionation and time interval until surgery were different in both groups (5x5 Gy followed by immediate surgery versus 50.4 Gy followed by surgery after a waiting period of 4-6 weeks). In this study, therefore, it remains unclear whether the observed differences in tumor response between both arms are attributable to the addition of chemotherapy or to a different RT schedule and a different waiting period until surgery. Since, moreover, actuarial local recurrence data at five years are not available in this study, it was left out from the meta-analysis of local recurrence at five years. This analysis demonstrates a significant reduction in local recurrence rate with the addition of chemotherapy (OR 0.39-0.72, $P < 0.001$). Importantly, the cumulative incidence rates of local recurrence in the RT group of the studies of Bosset 2006 and Gerard 2006 (17%) and in both groups of the study of Boulis-Wassif 1984 (15%) seem high compared to the 5.5% local recurrence rate at five years achieved by the Dutch rectal cancer trial using 5x5 Gy preoperative RT followed by surgery (van den Brink 2004). Differences in stage distribution and variation in surgical technique might explain this observation. Indeed, during the Dutch rectal cancer trial a formal surgical training and quality control program was implemented in order to guarantee optimal surgery (TME). The study of Boulis-Wassif 1984 predated the introduction of TME surgery (inclusion period 1972-1976), whereas in the studies of Bosset 2006 and Gerard 2006, TME surgery was 'recommended' without any formal surgical training or quality control.

Although in the study of [Boulis-Wassif 1984](#) a marginally significant five year survival benefit was associated with CRT, the combined analysis failed to demonstrate a significant difference in either overall or disease free survival at five years (OR 0.79-1.14, $P = 0.58$ and OR 0.92-1.34, $P = 0.27$, respectively). One of the hypotheses formulated to explain the observed lack of survival benefit found in many pre- or postoperative adjuvant therapy trials in rectal cancer is the existence of early, subclinical systemic disease present at diagnosis. This hypothesis is supported by the finding that the rate of distant metastatic disease in all four trials is consistently around 30%, without any difference between CRT and RT groups, indicating a future role of more effective systemic therapy to eradicate micrometastatic disease from the onset of therapy. Others have argued that the follow up time of the included trials is too short to observe a survival benefit, or that the incidence of local recurrence is too low to influence survival ([Gerard 2006](#)).

Grade III and IV acute treatment related toxicity was more pronounced in the CRT group in the three studies reporting this parameter ([Bosset 2006](#); [Bujko 2006](#); [Gerard 2006](#)), with an overall OR of 1.68-10 and a P value of 0.002. However, chemotherapy related toxicity was generally acceptable as evidenced by the high compliance rates in the studies mentioned (82%, 78.1%, and 69% respectively). In resected patients, no differences were observed in 30 day mortality, postoperative morbidity, or anastomotic leak rate. The results concerning anastomotic leakage should be interpreted with caution, since the exceedingly low leakage rate in the study of [Bosset](#) compared to currently accepted and published leakage rates following anterior resection suggests underreporting of this specific complication.

Postoperative quality of life (QoL) is an important, though often underreported aspect of cancer trials. From the Swedish and Dutch rectal cancer trials, it is known that preoperative 5x5 Gy followed by surgery significantly worsens functional outcome in terms of bowel, sexual, and bladder function compared to surgery alone ([Holm 1996](#); [Dahlberg 1998](#); [Peeters 2005](#)). A number of phase II trials have suggested that preoperative CRT followed by surgery does not adversely affect functional outcome ([Feliu 2002](#); [Bosset 2000](#)). The scarce available data in the four included studies did not allow to perform a meta-analysis of QoL related parameters. However, preliminary functional outcome data of the EORTC 22921 study (published as abstract only) demonstrated a significantly worse anorectal function in CRT patients compared to RT alone ([Mercier 2005](#)). Interestingly, in the final results paper of this study the incidence of 'late side effects' including fecal incontinence did not seem to differ between the four treatment arms ([Bosset 2006](#)).

The results of the meta-analysis confirm the enhanced cytotoxic efficacy of combined RT with 5-fluorouracil based chemotherapy. The incidence of a complete pathological response (pCR, ypT0N0) was 129 of 1096 (11.8%) patients in the CRT group and 39 of 1105 (3.5%) patients in the RT group; this difference

was statistically highly significant (OR 3.65, 95%CI 2.52-5.27, $P < 0.001$) while no heterogeneity was observed between the four studies. The results of the EORTC study moreover confirmed the difference in radioresponsiveness of the tumor in the bowel wall compared to that of mesorectal lymph nodes, as evidenced by nodal involvement in up to 12% of ypT0 patients ([Bosset 2004](#)). Although in two studies ([Bosset 2006](#); [Boulis-Wassif 1984](#)) a trend towards increased sphincter preservation was observed in the CRT group, the overall results failed to demonstrate an increase in sphincter preserving surgery following CRT (OR 0.92-1.31, $P = 0.29$), notwithstanding the downsizing and downstaging effect often noted with the combined therapy. This finding may be related to reluctance of the colorectal surgeon to alter a preoperative assessment of the need to perform a rectal amputation, since reversal of this decision would possibly imply performing an anastomosis in previously macroscopically invaded tissue. Moreover, in at least two studies it was specifically advised not to change a preoperative decision to perform a rectal amputation even after a significant downsizing. Data from the German rectal cancer trial, however, suggested that a change in operative strategy (ie, perform sphincter preserving surgery when a significant clinical response is observed) may be safely performed. Longer follow up will be needed to confirm the safety of this approach.

AUTHORS' CONCLUSIONS

Implications for practice

Compared to preoperative RT alone, preoperative CRT enhances tumor response and improves local recurrence rates. Therefore, the addition of CT should be considered when the tumor is located close (< 2 mm) to the circumferential resection margin or the sphincter apparatus. The addition of chemotherapy causes a moderate increase in acute toxicity, although postoperative complications including anastomotic leakage are not enhanced.

At this moment, it is unclear from the available data whether the addition of chemotherapy to preoperative RT influences sphincter preservation. Patients should be informed about the possible functional and QoL related aspects of preoperative therapy.

Implications for research

1. Since the improvement of local control obtained with CRT did not translate into a better overall or disease free survival and up to one third of all patients develop distant spread, priority should be given to trials addressing early subclinical systemic spread;
2. Trials are needed that specifically address the oncological safety of performing sphincter preserving surgery (including intersphincteric resection and colo-anal anastomosis) in patients deemed to require amputation before the start of CRT and in whom a significant clinical response is observed;

3. Preoperative therapy trials in rectal cancer should include formal evaluation of functional and QoL related aspects of therapy.

ACKNOWLEDGEMENTS

None

REFERENCES

References to studies included in this review

Bosset 2006 {published data only}

Bosset JF, Collette L, Calais G, Mineur L, Maingon P, Radosevic-Jelic L, Daban A, Bardet E, Beny A, Ollier JC, EORTC Radiotherapy Group Trial 22921. Chemotherapy with Preoperative Radiotherapy in Rectal Cancer. *The New England Journal of Medicine* 2006;**355**:1114–23.

Boulis-Wassif 1984 {published data only}

S. Boulis-Wassif, A. Gerard, J. Loygue, D. Camelot, M. Buyse, N. Duez. Final Results of a Randomized Trial on the Treatment of Rectal Cancer with Preoperative Radiotherapy Alone or in Combination With 5-Fluorouracil, Followed by radical surgery. *Cancer* 1984;**53**:1811–1818.

Bujko 2006 {published data only}

Bujko K, Nowacki MP, Nasierowska-Guttmejer A, Michalski W, Bebenek M, Kryj M. Long-term results of a randomized trial comparing preoperative short-course radiotherapy with preoperative conventionally fractionated chemoradiation for rectal cancer. *British Journal of Surgery* 2006;**93**:1215–1223.

Gerard 2006 {published data only}

Jean-Pierre Gérard, Thierry Conroy, Franck Bonnetain, Olivier Bouché, Olivier Chapet, Marie-Thérèse Closon-Dejardin, Michel Untereiner, Bernard Leduc, Eric Francois, Jean Maurel, Jean-Francois Seitz, Bruno Buecher, Rémy Mackiewicz, Michel Ducreux, Laurent Bedenne. Preoperative Radiotherapy With or Without Concurrent Fluorouracil and Leucovorin in T3-T4 Rectal Cancers: Results of FFC0 9203. *J Clin Oncol* 2006;**24**:4620–4625.

Additional references

Ammann 2003

Ammann K, Kirchmayr W, Klaus A, Muhlmann G, Kafka R, Oberwalder M, De Vries A, Ofner D, Weiss H. Impact of neoadjuvant chemoradiation on anal sphincter function in patients with carcinoma of the midrectum and low rectum. *Archives of Surgery* 2003;**138**(3):257–261.

Beets-Tan 2003

Beets-Tan RGH, Beets GL. Rectal cancer: how accurate can imaging predict the T stage and the circumferential resection margin?. *International Journal of Colorectal Disease* 2003;**18**(5): 385–391.

Beets-Tan 2005

Beets-Tan RGH, Lettinga T, Beets GL. Pre-operative imaging of rectal cancer and its impact on surgical performance and treatment outcome. *Ejso* 2005;**31**(6):681–688.

Bosset 2000

Bosset JF, Magnin V, Maingon P, Manton G, Pelissier EP, Mercier M, Chaillard G, Horiot JC. Preoperative radiochemotherapy in rectal cancer: Long-term results of a phase II trial. *International Journal of Radiation Oncology Biology Physics* 2000;**46**(2):323–327.

Bosset 2004

Bosset JF, Calais G, Daban A, Berger C, Radosevic-Jelic L, Maingon P, Bardet E, Pierart M, Briffaux A. Preoperative chemoradiotherapy versus preoperative radiotherapy in rectal cancer patients: assessment of acute toxicity and treatment compliance - Report of the 22921 randomised trial conducted by the EORTC Radiotherapy Group. *European Journal of Cancer* 2004;**40**(2):219–224.

Brown 2004

Brown G, Kirkham A, Williams GT, Bourne M, Radcliffe AG, Sayman J, Newell R, Sinnatamby C, Heald RJ. High-resolution MRI of the anatomy important in total mesorectal excision of the rectum. *American Journal of Roentgenology* 2004;**182**(2):431–439. [: 0361–803X]

Brown 2006

Brown G, Daniels IR, Heald RJ, Quirke P, Blomqvist L, Sebag-Montefiore D, Moran BJ, Holm T, Strassbourg J, Peppercorn PD, Fisher SE, Mason B. Diagnostic accuracy of preoperative magnetic resonance imaging in predicting curative resection of rectal cancer: prospective observational study. *British Medical Journal* 2006;**333**(7572):779–782. [: 0959–8146]

Dahlberg 1998

Dahlberg M, Glimelius B, Graf W, Pahlman L. Preoperative irradiation affects functional results after surgery for rectal cancer - Results from a randomized study. *Diseases of the Colon & Rectum* 1998;**41**(5):543–549.

Dale 2005

Dale R, Carabe-Fernandez A. The radiobiology of conventional radiotherapy and its application to radionuclide therapy. *CANCER BIOTHERAPY AND RADIOPHARMACEUTICALS* 2005;**20**(1):47–51.

Daniels 2005

Daniels IR. High spatial resolution MRI predicts turnout spread in patients with rectal cancer - Results from the MERCURY study. *Diseases of the Colon & Rectum* 2005;**48**(3):688–689.

Enker 1999

Enker WE, Merchant N, Cohen AM, Lanouette NM, Swallow C, Guillem J, Paty P, Minsky B, Weyrauch K, Quan SHQ. Safety and efficacy of low anterior resection for rectal cancer - 681 consecutive cases from a specialty service. *Annals of Surgery* 1999;**230**(4):544–552.

Feliu 2002

Feliu J, Calvillo J, Escribano A, de Castro J, Sanchez ME, Mata A, Espinosa E, Grande AG, Mateo A, Baron MG. Neoadjuvant therapy of rectal carcinoma with UFT-leucovorin plus radiotherapy. *Annals of Oncology* 2002;**13**(5):730–736.

Glimelius 2002

Glimelius B. Radiotherapy in rectal cancer. *Br Med Bull* 2002;**64**:141–57.

Gray 2001

Gray R, Hills R, Stowe R, Clarke M, Peto R, Buyse M, Piedbois P. Adjuvant radiotherapy for rectal cancer: a systematic overview of 8507 patients from 22 randomised trials. *Lancet* 2001;**358**(9290):1291–1304.

Heald 1982

Heald RJ, Husband EM, Ryall RDH. The Mesorectum in Rectal-Cancer Surgery - the Clue to Pelvic Recurrence. *British Journal of Surgery* 1982;**69**(10):613–616. [0007–1323]

Heald 1998

Heald RJ, Moran BJ, Ryall RDH, Sexton R, MacFarlane JK. Rectal cancer - The Basingstoke experience of total mesorectal excision, 1978-1997. *Archives of Surgery* 1998;**133**(8):894–898.

Higgins 2008

Higgins JPT, Green S (editors). Cochrane Handbook for Systematic Reviews of Interventions Version 5.0.0 [updated February 2008]. The Cochrane Collaboration. Available from www.cochrane-handbook.org. 2008.

Holm 1996

Holm T, Singnomklo T, Rutqvist LE, Cedermark B. Adjuvant preoperative radiotherapy in patients with rectal carcinoma - Adverse effects during long term follow-up of two randomized trials. *Cancer* 1996;**78**(5):968–976.

Kapiteijn 2001

Kapiteijn E, Marijnen CAM, Nagtegaal ID, Putter H, Steup WH, Wiggers T, Rutten HJT, Pahlman L, Glimelius B, van Krieken J, Leer JWH, van de Velde CJH. Preoperative radiotherapy combined with total mesorectal excision for resectable rectal cancer. *New England Journal of Medicine* 2001;**345**(9):638–646.

Mercier 2005

Mercier M, Pasquet P, Puyraveau M, Bosset J, Calais G, Mineur L, Maingon P, Daban A, Bardet E, Collette L. Evaluation of the sphincter function and quality of life in French patients with rectal cancer who entered the EORTC 22921 study. *Eur J Cancer* 2005; **Supplements 3, No.2**:171.

Minsky 1993

Minsky BD, Cohen AM, Kemeny N, Enker WE, Kelsen DP, Schwartz G, Saltz L, Dougherty J, Frankel J, Wiseberg J. Preoperative Combined 5-Fu, Low-Dose Leucovorin, and Sequential Radiation-Therapy for Unresectable Rectal-Cancer. *International Journal of Radiation Oncology Biology Physics* 1993;**25**(5):821–827.

Minsky 1997

Minsky BD, Cohen AM, Enker WE, Saltz L, Guillem JG, Paty PB, Kelsen DP, Kemeny N, Ilson D, Bass J, Conti J. Preoperative 5-FU, low-dose leucovorin, and radiation therapy for locally advanced and unresectable rectal cancer. *International Journal of Radiation Oncology Biology Physics* 1997;**37**(2):289–295.

Nagtegaal 2002

Nagtegaal ID, Marijnen CAA, Kranenbarg EK, van de Velde CJH, van Krieken J. Circumferential margin involvement is still an important predictor of local recurrence in rectal carcinoma - Not one millimeter but two millimeters is the limit. *American Journal of Surgical Pathology* 2002;**26**(3):350–357.

Osti 2004

Osti MF, Valeriani M, Masoni L, Tombolini V, Enrici RM. Neoadjuvant chemoradiation for locally advanced carcinoma of the rectum. *Tumori* 2004;**90**(3):303–309.

Peeters 2005

Peeters K, van de Velde CJH, Leer JWH, Martijn H, Junggeburst JMC, Kranenbarg EK, Steup WH, Wiggers T, Rutten HJ, Marijnen CAM. Late side effects of short-course preoperative radiotherapy combined with total mesorectal excision for rectal cancer: Increased bowel dysfunction in irradiated patients - A dutch colorectal cancer group study. *Journal of Clinical Oncology* 2005;**23**(25):6199–6206.

Quirke 1986

Quirke P, Dixon MF, Durley P, Williams NS. Local Recurrence of Rectal Adenocarcinoma Due to Inadequate Surgical Resection - Histopathological Study of Lateral Tumor Spread and Surgical Excision. *Lancet* 1986;**2**(8514):996–999. [0140–6736]

Quirke 2003

Quirke P. Training and quality assurance for rectal cancer: 20 years of data is enough. *Lancet Oncology* 2003;**4**(11):695–702.

Rodel 2003

Rodel C, Grabenbauer GG, Papadopoulos T, Hohenberger W, Schmoll HJ, Sauer R. Phase I/II trial of capecitabine, oxaliplatin, and radiation for rectal cancer. *Journal of Clinical Oncology* 2003;**21**(16):3098–3104.

Sauer 2004

Sauer R, Becker H, Hohenberger W, Rodel C, Wittekind C, Fietkau R, Martus P, Tschmelitsch J, Hager E, Hess CF, Karstens JH, Liersch T, Schmidberger H, Raab R. Preoperative versus postoperative chemoradiotherapy for rectal cancer. *New England Journal of Medicine* 2004;**351**(17):1731–1740.

van den Brink 2004

van den Brink M, Stiggelbout AM, van den Hout WB, Kievit J, Kranenbarg EK, Marijnen CAM, Nagtegaal ID, Rutten HJT, Wiggers T, van de Velde CJH. Clinical nature and prognosis of locally recurrent rectal cancer after total mesorectal excision with or

without preoperative radiotherapy. *Journal of Clinical Oncology*
2004;**22**(19):3958–3964. [: 0732–183X]

* *Indicates the major publication for the study*

CHARACTERISTICS OF STUDIES

Characteristics of included studies *[ordered by study ID]*

Bosset 2006

Methods	<ol style="list-style-type: none"> 1. Randomization method: telephone to central office (assumed) 2. Abdominal imaging: CT 3. Chest imaging: CXR 4. 4 arm study: Arm 1 preop RT + S; Arm 2 preop XRT + concurrent 5FU LV + S; Arm 3 preop RT + S + post op 5FU LV ; Arm 4 preop RTCT+ S + postop 5FU LV 5. Total randomized 1011
Participants	<ol style="list-style-type: none"> 1. Rectal Cancer 2. Location: within 15 cm from anal verge 3. Resectability: locally resectable 4. T3 or resectable T4 (defined by clinical criteria or endoscopic ultrasound) 5. WHO PS 0-1 6. </=80yr
Interventions	<ol style="list-style-type: none"> 1. Surgery: AP/anterior resection or Hartman with TME 2. RT : 45 Gy in 25fr. 3. RT volume: 5cm above and below tumor and perirectal nodes below S2-3. If tumor above 10cm, include only 3 cm above tumor. If tumor in low rectum, S2-3 to perineum. Posteriorly to include entire sacrum with 3cm beyond macroscopic extension 4. RT-S: within 3-10 weeks of completing neoadjuvant therapy 5. 3 or 4 field 6. Chemotherapy: 5FU 325mg/m2/d; Leucovorin 20mg/m2/day Dy1-5 & 28-32 for arms 2 and 4, and postoperative for arms 3 and 4
Outcomes	<ol style="list-style-type: none"> 1. Duration of FU: 5.4 yrs 2. Perioperative mortality: CRTS 2.4 % (12/506) RTS 1.2% (6/505) 3. Mets (liver) @ lap: Y 4. Curative resection: not stated 5. Overall resection: 94.5 % 6. Compliance to radiotherapy: CRTS 483/506 (95.5%) RTS 495/505 (98.0%) 7. OAS: Y 8. CSS? 9. Tox post RT: Y 10. Acute tox post S: Y 11. Late tox post S: Y 12. LR: Y 13. QOL: N 14. Proportion sfincter sparing: CRTS 267/506 (52.8%) RTS 255/505 (50.5%) 15. Proportion downstaging: Y
Notes	
<i>Risk of bias</i>	

Bosset 2006 (Continued)

Item	Authors' judgement	Description
Allocation concealment?	Yes	A - Adequate

Boulis-Wassif 1984

Methods	<ol style="list-style-type: none"> 1. Randomization method: not stated. conducted by cooperative group. Likely via central office 2. Abdominal imaging: Not stated 3. Chest imaging: Not stated 4. Study arm (Preop chemoradiotherapy arm) : 171 randomized, 45 excluded. 5. Control arm (Preop radiotherapy arm): 168 randomized , 47 excluded
Participants	<ol style="list-style-type: none"> 1. Rectal Cancer 2. Location: below within 15cm anal verge 3. Resectability: fit for surgery
Interventions	<ol style="list-style-type: none"> 1. Surgery: AP/anterior resection 2. RT : 3450 cGy in 15fr. (for both arms) 3. BED: 35.2Gy1 4. RT volume: "chimney and diamond fields" paraaortics and pelvis. 5. RT-S: within 2 wk 6. 2 field 7. Cointervention: none 8. 2 arms, control (Radiotherapy followed by surgery), Study (Chemoradiotherapy followed by surgery) 9. Chemotherapy: 5FU 10mg/kg/d day 1-4
Outcomes	<ol style="list-style-type: none"> 1. Duration of FU: mean 5.2yrs 2. Perioperative mortality: CRTS 19/126 RTS 11/121 3. Mets @ lap: CRTS 13/126 RTS 15/121 4. Curative resection: Not stated 5. Overall resection: CRTS 121, RTS 124 6. Compliance to radiotherapy: not given 7. OAS: Y 8. CSS: N 9. Tox post RT: not given 10: Acute tox post S: not given No complication not given 11. Late tox post S: not given 12: LR: N 13. QoL:N
Notes	

Risk of bias

Item	Authors' judgement	Description
------	--------------------	-------------

Boulis-Wassif 1984 (Continued)

Allocation concealment?	Yes	A - Adequate
-------------------------	-----	--------------

Bujko 2006

Methods	<ol style="list-style-type: none"> 1. Randomization method: telephone to central office 2. Abdominal imaging: ultrasound or CT 3. Chest imaging: CXR 4. XRT + S arm : (short XRT) 155 randomized, 0 excluded. 5. Arm B: (Long XRT+CT): 157 randomized , 0 excluded 	
Participants	<ol style="list-style-type: none"> 1. Rectal Cancer 2. Location: inferior edge palpable of digital exam 3. Resectability: locally resectable 4. T3 or resectable T4 5. not involving sphincter 	
Interventions	<ol style="list-style-type: none"> 1. Surgery: AP/anterior resection or Hartman with TME 2. RT : XRT +S arm: 2500cGy cGy in 5fr. ; Arm B: 50.4Gy in 28 fr with concomitant CT weeks 1 & 5 3. BED: Arm A 38.7Gy10, Arm B 40.9Gy10 4. RT volume: Not stated 5. RT-S: XRT+S within 7 days; Arm B: within 4-6 weeks 6. 3 or 4 field 7. Arm B chemotherapy: 5FU 325mg/m2/d; Leucovorin 20mg/m2/day Dy1-5 & 28-32 	
Outcomes	<ol style="list-style-type: none"> 1. Duration of FU: not stated 2. Perioperative mortality: XRT+S 0/155 Arm B 0/157 3. Mets @ lap: not stated 4. Curative resection: not stated 5. Overall resection: XRT+S 145/155 Arm B 147/157 6. Compliance to radiotherapy: XRT+S 152/155 Arm B 141/157 7. OAS: N 8. CSS: N 9. Tox post RT: no complications XRT+S 118/155 Arm B 24/157 Any complications XRT+S 37/155 Arm B 133/157 Gd 3-4 XRT + S 5/155 Arm B 26/157 Gd 5 (Death) XRT +S 0/155 Arm B 2/157 10. Acute tox post S: Not stated 11. Late tox post S: not given 12. LR: N 13. QoL: N 14. Proportion sphincter sparing 15. Proportion downstaged (by T stage, N stage, Tumor size) 	
Notes		
Risk of bias		
Item	Authors' judgement	Description

Bujko 2006 (Continued)

Allocation concealment?	Yes	A - Adequate
-------------------------	-----	--------------

Gerard 2006

Methods	1. Randomization method: not stated 2. Abdominal imaging: liver sonography - CT scan 3. Chest imaging: CXR 4. Study arm: CRT 375 5. Control arm: RT 367	
Participants	1. Rectal Cancer 2. Location: accessible by digital examination 3. Resectability: locally resectable	
Interventions	2 arms: preop XRT vs preop CRT 1. Surgery: TME recommended 2. RT 45Gy in 25 fr for both arms 3. BED: 32.5Gy10 4. RT volume: NA 5. RT-S: NA 6. NA 7. Cointervention: postoperative CT (5FU FA) x 4 cycles	
Outcomes	1. Duration of FU: 81m 2. Perioperative mortality (60 days): 2% for both arms 3. Mets @ lap: not stated 4. Curative resection: not stated 5. Overall resection: not stated 6. Compliance to radiotherapy: not stated 7. OAS: Y 8. CSS: Y 9. Tox (gr 3-4) post RT: Preop RT arm: 10/367 CRT arm: 55/375 10: Acute tox post S: not stated 11. Late tox post S: Y	
Notes		
<i>Risk of bias</i>		
Item	Authors' judgement	Description
Allocation concealment?	Yes	A - Adequate

DATA AND ANALYSES

Comparison 1. radiotherapy vs radiochemotherapy


Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Overall Survival at 5y	3	2000	Odds Ratio (M-H, Fixed, 95% CI)	0.95 [0.79, 1.14]
2 Disease free survival at 5 y	2	1753	Odds Ratio (M-H, Fixed, 95% CI)	1.11 [0.92, 1.34]
3 Mortality 30 d	4	2239	Odds Ratio (M-H, Fixed, 95% CI)	1.48 [0.84, 2.60]
4 Postop morbidity	3	1994	Odds Ratio (M-H, Fixed, 95% CI)	0.84 [0.68, 1.03]
5 Grade III - IV toxicity	3	2032	Odds Ratio (M-H, Random, 95% CI)	4.10 [1.68, 10.00]
6 Sphincter preservation	4	2219	Odds Ratio (M-H, Fixed, 95% CI)	1.10 [0.92, 1.31]
7 pCR	4	2201	Odds Ratio (M-H, Fixed, 95% CI)	3.65 [2.52, 5.27]
8 Anastomotic leak	3	1068	Odds Ratio (M-H, Fixed, 95% CI)	1.03 [0.57, 1.85]
9 Local Recurrence at 5y	3	1494	Odds Ratio (M-H, Fixed, 95% CI)	0.53 [0.39, 0.72]

Analysis 1.1. Comparison 1 radiotherapy vs radiochemotherapy, Outcome 1 Overall Survival at 5y.

Review: Preoperative chemoradiation versus radiation alone for stage II and III resectable rectal cancer

Comparison: 1 radiotherapy vs radiochemotherapy

Outcome: 1 Overall Survival at 5y




Analysis 1.2. Comparison 1 radiotherapy vs radiochemotherapy, Outcome 2 Disease free survival at 5 y.

Review: Preoperative chemoradiation versus radiation alone for stage II and III resectable rectal cancer

Comparison: 1 radiotherapy vs radiochemotherapy

Outcome: 2 Disease free survival at 5 y




Analysis 1.3. Comparison 1 radiotherapy vs radiochemotherapy, Outcome 3 Mortality 30 d.

Review: Preoperative chemoradiation versus radiation alone for stage II and III resectable rectal cancer

Comparison: 1 radiotherapy vs radiochemotherapy

Outcome: 3 Mortality 30 d




Analysis I.4. Comparison I radiotherapy vs radiochemotherapy, Outcome 4 Postop morbidity.

Review: Preoperative chemoradiation versus radiation alone for stage II and III resectable rectal cancer

Comparison: I radiotherapy vs radiochemotherapy

Outcome: 4 Postop morbidity




Analysis I.5. Comparison I radiotherapy vs radiochemotherapy, Outcome 5 Grade III - IV toxicity.

Review: Preoperative chemoradiation versus radiation alone for stage II and III resectable rectal cancer

Comparison: I radiotherapy vs radiochemotherapy

Outcome: 5 Grade III - IV toxicity




Analysis I.6. Comparison I radiotherapy vs radiochemotherapy, Outcome 6 Sphincter preservation.

Review: Preoperative chemoradiation versus radiation alone for stage II and III resectable rectal cancer

Comparison: I radiotherapy vs radiochemotherapy

Outcome: 6 Sphincter preservation




Analysis I.7. Comparison I radiotherapy vs radiochemotherapy, Outcome 7 pCR.

Review: Preoperative chemoradiation versus radiation alone for stage II and III resectable rectal cancer

Comparison: I radiotherapy vs radiochemotherapy

Outcome: 7 pCR




Analysis I.8. Comparison I radiotherapy vs radiochemotherapy, Outcome 8 Anastomotic leak.

Review: Preoperative chemoradiation versus radiation alone for stage II and III resectable rectal cancer

Comparison: I radiotherapy vs radiochemotherapy

Outcome: 8 Anastomotic leak




Analysis I.9. Comparison I radiotherapy vs radiochemotherapy, Outcome 9 Local Recurrence at 5y.

Review: Preoperative chemoradiation versus radiation alone for stage II and III resectable rectal cancer

Comparison: I radiotherapy vs radiochemotherapy

Outcome: 9 Local Recurrence at 5y



WHAT'S NEW

Last assessed as up-to-date: 1 September 2008.

27 May 2008	Amended	Converted to new review format.
-------------	---------	---------------------------------

HISTORY

Protocol first published: Issue 2, 2006

Review first published: Issue 1, 2009

26 February 2007	New citation required and conclusions have changed	Substantive amendment
------------------	--	-----------------------

CONTRIBUTIONS OF AUTHORS

Drafted the protocol: WPC

Developed the search strategy: WPC

Searched for trials: WPC and YVN

Trials selection for inclusion: WPC and YVN

Data extraction: KF

Data into revman analyses: KF

Interpretation of analyses: WPC, YVN and KF

DECLARATIONS OF INTEREST

None.

INDEX TERMS

Medical Subject Headings (MeSH)

*Neoplasm Recurrence, Local [prevention & control]; Randomized Controlled Trials as Topic; Rectal Neoplasms [*drug therapy; pathology; *radiotherapy; surgery]

MeSH check words

Humans