Journal Pre-proof

MENTAL HEALTH DURING THE COVID-19 PANDEMIC AND BEYOND: THE IMPORTANCE OF THE VAGUS NERVE FOR BIOPSYCHOSOCIAL RESILIENCE

Josefien Dedoncker, Marie-Anne Vanderhasselt, Cristina Ottaviani,
George M. Slavich

PII: S0149-7634(21)00064-6
DOI: <https://doi.org/10.1016/j.neubiorev.2021.02.010>
Reference: NBR 4062

To appear in: *Neuroscience and Biobehavioral Reviews*

Received Date: 5 December 2020
Revised Date: 27 January 2021
Accepted Date: 4 February 2021

Please cite this article as: Dedoncker J, Vanderhasselt M-Anne, Ottaviani C, Slavich GM, MENTAL HEALTH DURING THE COVID-19 PANDEMIC AND BEYOND: THE IMPORTANCE OF THE VAGUS NERVE FOR BIOPSYCHOSOCIAL RESILIENCE, *Neuroscience and Biobehavioral Reviews* (2021), doi: <https://doi.org/10.1016/j.neubiorev.2021.02.010>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier.

MENTAL HEALTH DURING THE COVID-19 PANDEMIC AND BEYOND: THE IMPORTANCE OF THE VAGUS NERVE FOR BIOPSYCHOSOCIAL RESILIENCE

Running Head: VAGUS NERVE, MENTAL HEALTH, & BIOPSYCHOSOCIAL RESILIENCE

Josefien DEDONCKER^{a,b*}, Marie-Anne VANDERHASSELT^{a,b,c}, Cristina OTTAVIANI^{d,e}, George M. SLAVICH^f

^a Department of Head and Skin - Psychiatry and Medical Psychology, Ghent University Hospital, Ghent, Belgium

^b Ghent Experimental Psychiatry (GHEP) lab, Ghent, Belgium

^c Department of Experimental Clinical and Health Psychology, Ghent University, Ghent, Belgium

^d Department of Psychology, Sapienza University of Rome, Rome, Italy

^e Neuroimaging Laboratory, IRCCS Santa Lucia Foundation, Rome, Italy

^f Cousins Center for Psychoneuroimmunology and Department of Psychiatry and Biobehavioral Sciences, University of California, Los Angeles, CA, USA

*Corresponding Author

Josefien Dedoncker, PhD. Department of Head and Skin - Psychiatry and Medical Psychology, Ghent University Hospital, C. Heymanslaan 10 – 9000 Ghent, Belgium. Tel: +32 (0)9 332 5894.

E-mail: Josefien.Ddr@gmail.com

Highlights

- The COVID-19 pandemic has led to a notable increase in mental health problems
- Both viral and psychosocial stressors can trigger changes in biological processes that affect immunity and inflammation
- Vagus nerve function is one pathway that is central to health-relevant mind-body interactions
- Lifestyle changes and non-invasive tVNS are two strategies for improving vagus nerve function
- Improved vagus nerve function may in turn help promote biopsychosocial resilience

Abstract

The COVID-19 pandemic has led to widespread increases in mental health problems, including anxiety and depression. The development of these and other psychiatric disorders may be related to changes in immune, endocrine, autonomic/physiological, cognitive, and affective processes induced by a SARS-CoV-2 infection. Interestingly, many of these same changes can be triggered by psychosocial stressors such as social isolation and rejection, which have become increasingly common due to public policies aimed at reducing the spread of SARS-CoV-2. The present review aims to shed light on these issues by describing how viral infections and stress affect mental health. First, we describe the multi-level mechanisms linking viral infection and life stress exposure with risk for psychopathology. Then, we summarize how resilience can be enhanced by targeting vagus nerve function by, for example, applying transcutaneous vagus nerve stimulation and targeting lifestyle factors, such as exercise. With these biopsychosocial insights in mind, researchers and healthcare professionals will be better equipped to reduce risk for psychopathology and increase resilience during this challenging pandemic period and beyond.

Keywords: COVID-19; Coronavirus; Lifestyle Interventions; Psychiatric Disorders; Social Stress; Transcutaneous Vagus Nerve Stimulation

1. Introduction

The effects of the recent Coronavirus Disease 2019 (COVID-19) pandemic on mental health are becoming increasingly evident. Recent research has suggested that one in five COVID-19 patients present with a (new-onset) mental health issue (Kong et al., 2020; Varatharaj et al., 2020). Moreover, prior studies in SARS-CoV-1 patients showed that psychopathology can develop even when evaluated years after infection (Lam et al., 2009). More broadly, research has indicated that exposure to infection and inflammation significantly increase the risk of developing psychopathology later in life (Benros et al., 2011, 2013; Khandaker et al., 2014; Rosenblat et al., 2014), in part by triggering aberrant mind-body interactions (Rosenblat et al., 2014). At the heart of these mind-body interactions is the autonomic nervous system (ANS) and, more specifically, the vagus nerve, which carries most of the parasympathetic nervous system (PNS) fibers and acts as a major bidirectional communication pathway between the central nervous system (CNS) and the body.

One factor that strongly affects the activity of the vagus nerve is stress, and during the COVID-19 pandemic, people have been exposed to a variety of different psychosocial stressors that could exert vagus nerve-altering effects. In addition to notable increases in work-, financial-, and housing-related stressors (Nelson et al., 2020; Nicola et al., 2020), individuals have also experienced increases in social isolation (e.g., due to social distancing protocols) and rejection (e.g., due to disagreements involving disease prevention efforts, which have been polarizing and political). Similar to viral infections, social isolation and rejection have been shown to upregulate inflammatory activity, which can in turn lead to aberrant mind-body interactions that negatively impact mental health (Furman et al., 2019; Slavich, 2016, 2020a). Consequently, some researchers have hypothesized that we may witness a 'wave' of people developing mental health issues in the near future that is caused by the heightened risk of viral infection combined with high levels of psychosocial stress (Raony et al., 2020).

The goal of the present review is to describe how viral and social stressors can impact the body to affect mental health. We begin by reviewing the literature describing the impact of a SARS-CoV infection (i.e., SARS-CoV-1 and SARS-CoV-2) on mind-body interactions and the development of mental health issues. Second, we discuss the relation between perceived or actual social exclusion and mental distress, with the aim of providing mechanistic insights into how mind-body interactions may play an important role in influencing mental health. Lastly, based on these mechanistic insights, we highlight several interventions that can target aberrant mind-body interactions by enhancing vagal nerve function, with the hope that such interventions may help enhance resilience, prevent or reduce risk for mental health problems, and lessen the high levels of personal suffering and societal disease burden attributable to mental illness.

2. SARS-CoV-2 Infection, Maladaptive Changes in Mind-Body Interactions, and Mental Health

Over the past few decades, research has shown that exposure to infection and inflammation can significantly increase the risk of developing psychopathology, even during later life (Benros et al., 2011, 2013; Khandaker et al., 2014; Rosenblatt et al., 2014; Slavich & Irwin, 2014). This effect has also been observed in the COVID-19 pandemic. In a recent UK-wide surveillance study, it was shown that one out of five (hospitalized) COVID-19 patients presented with an altered mental status predominantly elicited (in 92% of cases) by a new-onset neuropsychiatric disorder, such as psychosis, neurocognitive dementia-like syndrome, or an affective disorder (Varatharaj et al., 2020). This effect has been replicated by another study, which indicated that being diagnosed with COVID-19 doubled the risk of being newly diagnosed with a psychiatric disorder (Taquet et al., 2020). Other reports have also revealed an increased prevalence of mood and anxiety disorders, as well as symptoms of posttraumatic stress disorder (PTSD) in acute phase COVID-19 patients, especially when they are older, female, and/or have less self-reported social support (Bo et al., 2020; Kong et al., 2020). These findings

are consistent with the described acute phase as well as longer phase mental health outcomes of SARS-CoV-1 (for a review, see Raony et al., 2020). Of interest, according to a recent review by Lam (2020), SARS-CoV-1 survivors have been most often diagnosed with PTSD (55%) or depression (39%), even when evaluated up to 4 years after the infection.

One mechanism by which exposure to an infection may lead to mental health problems is infection-induced maladaptive changes in mind-body interactions involving immune, endocrine, autonomic/physiological, cognitive, and affective functioning (see Figure 1, path A). First, viral-initiated increases in peripheral inflammation – characterized by enhanced pro-inflammatory cytokine [e.g., Interleukin-6 (IL-6) and Tumor Necrosis Factor- α (TNF- α)] signaling – can trigger activation of the hypothalamic-pituitary-adrenal (HPA) axis. More specifically, via the afferent fibers of the vagus nerve, which is the main nerve of the autonomic PNS, information regarding the presence of peripheral cytokines is conveyed to nuclei in the brainstem such as the nucleus tractus solitarius (NTS) and the hypothalamus, where it can trigger the release of corticotrophin-releasing hormone (CRH) and increase HPA axis activity. HPA axis activation, in turn, is characterized by enhanced cortisol release which can trigger a generalized physiological and psychological stress response (Raony et al., 2020; Rosenblat et al., 2014). In addition, this signal is propagated to brain regions implicated in physiological, cognitive, and emotional regulation, including the amygdala, insula, anterior cingulate cortex (ACC), and medial prefrontal cortex (mPFC) (Desforges et al., 2020; Pavlov & Tracey, 2012).

-- Insert Figure 1 Here --

To overcome spreading infection, transient cytokine responses and HPA axis activity are beneficial given the immunosuppressive effects of cortisol that trigger reduced cytokine signaling in immune cells (Wohleb & Godbout, 2013). However, when an infection is more intense or longer lasting and cytokine levels are more chronically enhanced, this can cause hyperactivation of the HPA axis. In this case, glucocorticoid receptors on immune cells become

less sensitive to the immunosuppressive effects of cortisol, leading to sustained cytokine signaling and inflammation, and a built-up of cortisol that has been called HPA axis overdrive (Pace & Miller, 2009). With respect to health, it has been hypothesized that sustained cytokine signaling in turn negatively impacts neurotransmitter metabolism (e.g., serotonin) and neuroplasticity, which have been implicated in the pathophysiology of psychopathology (Besedovsky & Del Ray, 2011; Rosenblat et al., 2014; Slavich & Irwin, 2014; Stapelberg et al., 2019; Wohleb & Godbout, 2013). Furthermore, peripheral inflammation may facilitate the development of reward hyposensitivity and anhedonia (Nusslock & Miller, 2016) due to inflammation-induced changes in dopamine signaling in the fronto-striatal reward circuit (for a review, see Miller et al., 2013).

It is through these biological interactions that SARS-CoV-2 infection-induced inflammation may lead to maladaptive mind-body interactions and thereby negatively impact mental health. In this regard, importantly, increasingly high serum levels of IL-6 and TNF- α have indeed been found depending on SARS-CoV-2 infection severity in COVID-19 patients (see Raony et al., 2020).

3. Psychosocial Stress, Maladaptive Changes in Mind-Body Interactions, and Mental Health

Interestingly, similarly to an infection, exposure to psychosocial stressors, including interpersonal loss and social rejection, has been shown to trigger changes in mind-body interactions that can have adverse health effects (Slavich et al., 2010a). As shown in Figure 1, path B, these changes include increased HPA axis activity and reactivity, activation of the innate immune system, and the release of pro-inflammatory cytokines that can lead to the development of systemic chronic inflammation (Haroon et al., 2012; Slavich, 2020b; Steptoe et al., 2007) even when assessed years later (Michopoulos et al., 2017; Tursich et al., 2014). These dynamics are of particular relevance to the present situation, given that during the

COVID-19 pandemic, many people have been repeatedly exposed to a variety of psychosocial stressors, including health, housing, financial, work, and interpersonal stressors (Nelson et al., 2020; Nicola et al., 2020).

Social isolation and rejection are two types of stress that are particularly relevant in this regard, especially given data showing that loneliness and sustained social exclusion have been key consequences of precautions designed to slow the spread of COVID-19 (Miller, 2020). In addition to being strong predictors of poor mental health (Slavich et al., 2009, 2014), these stressors are prevalent among individuals who have been living alone and those who have been prevented from meeting with family and friends (Mental Health Foundation, 2020; Pancani et al., 2020). Feelings of being socially criticized and/or rejected have also increased due to societal disagreement and polarization regarding government-installed health precautions (Van Bavel et al., 2020). Finally, many individuals have experienced a heightened sense of threat while outside, which has the benefit of helping to avoid contact with other people who could infect the individual with SARS-CoV-2 but has the downside of promoting neurocognitive appraisals of threat that upregulate inflammatory biology (Slavich et al., 2010b; Van Bavel et al., 2020).

This central role for social isolation and rejection in affecting health is consistent with Social Safety Theory (Slavich, 2020a), which posits that social isolation and rejection strongly upregulate inflammation, and with Social Baseline Theory, which argues that the social network is the primary source of safety (Coan, 2010). Indeed, loneliness is one factor that has been associated with generalized unsafety and the release of prolonged physiological activation (Brosschot et al., 2017). All in all, experiencing psychosocial stressors such as social isolation and rejection has been shown to negatively impact mental health (Benros et al., 2013; Rosenblat et al., 2014; Slavich, 2016), and this has been particularly relevant over the past year given that the COVID-19 pandemic has led to notable increases in the base rates of these stressors.

What is also interesting is that these social stressors have been shown to be particularly strongly related to changes in mind-body interactions that can affect health. As alluded to above, for example, social exclusion has been related to increased inflammatory activity, as indexed by IL-6, TNF- α , and C-reactive protein (CRP) (Slavich et al., 2010b; Smith et al., 2020). In fact, interpersonal loss and social exclusion stressors appear to be the strongest psychosocial triggers of inflammation (Slavich et al., 2010a). Social exclusion also has been related to (hyper) HPA axis activation and has been associated with a slower cortisol recovery after stressful experiences (Dickerson & Kemeny, 2004).

Theoretically, prolonged exposure to cortisol and pro-inflammatory cytokines as a result of chronic social exclusion may influence glucocorticoid receptor functioning on immune cells, which can cause glucocorticoid insensitivity and a reduced sensitivity of immune cells to the immunosuppressive influence of cortisol, leading to sustained inflammation (see above). Consistent with this possibility, social exclusion has been associated with abnormal glucocorticoid levels (Serra et al., 2005; Wohleb et al., 2015). In addition, exposure to social exclusion has been related to increased noradrenaline signaling and strong cardiovascular reactivity, which is indicative of autonomic sympathetic nervous system (SNS) activation rather than PNS activation (Slavich et al., 2010a), and to an increased risk for developing psychopathology (Masten et al., 2011; Monroe et al., 2014; Slavich, 2016). This may particularly true when the social isolation or exclusion is prolonged, as has been the case during the COVID-19 pandemic.

The mind-body interactions reviewed above that are triggered by psychosocial stress (i.e., in the absence of an infection) have also been directly implicated in the development of mental health problems such as mood and anxiety disorders, which may occur in part through the aforementioned impact of cytokine signaling on neurotransmitter activity or metabolism (Felger & Lotrich, 2013; Schiepers et al., 2005; Stapelberg et al., 2019). Consistent with this

possibility, meta-analyses have shown that levels of IL-6 and TNF- α are associated with the severity of several disorders including depression, PTSD, psychosis, and sleep disorders (Passos et al., 2015; Raony et al., 2020). This may help explain why recent lockdowns that aimed at slowing the transmission of SARS-CoV-2 have been related to an increased prevalence of depression, anxiety, PTSD, and other mental disorders (Brooks et al., 2020). Given the largescale and fast spread of SARS-CoV-2 and the major life stressors caused by the resulting public health precautions, the mental healthcare system can therefore be burdened even if only a fraction of COVID-19 survivors develops psychopathology. It is therefore crucial to understand the mechanisms underlying these increases in mental health problems during and in the aftermath of the COVID-19 pandemic, with this work ideally helping to inform the development of interventions that could reduce the likelihood of, or help prevent the development of, mental health problems.

4. The Vagus Nerve – At the Crossroad of Mind-Body Interactions

One key pathway linking exposure to both viruses (e.g., SARS-CoV-2 infection) and psychosocial stressors (e.g., social isolation and rejection) with mental health is the ANS. This system is centrally involved in the exchange of health-relevant information from the periphery to the brain and vice versa; additionally, it coordinates the body's reactivity to and recovery from stressors. As illustrated in Figure 1, the vagus nerve is particularly important in this regard, as it contains most of the PNS fibers.

The vagus nerve is the longest nerve of the body and is composed of approximately 80% afferent fibers that convey peripheral bodily information (e.g., cytokine levels) to the brain. The afferent vagal fibers enter the brain bilaterally and primarily target the NTS in the brainstem, the hypothalamus, and brain areas that have been implicated in the regulation of physiology, cognition, and affect (e.g., amygdala, ACC, mPFC; Pavlov & Tracey, 2012; Thayer & Sternberg, 2010). When pro-inflammatory cytokines are present due to an infection, for example, they can

activate afferent vagal neurons and trigger increased HPA axis activity and cortisol release (i.e., by triggering the release of CRH in the hypothalamus), leading to a systemic stress response (see above). Importantly, from the NTS, efferent vagal neurons also project back to cells in different organ systems throughout the body, including the heart, lungs, and gastrointestinal system. These efferent vagal neurons release acetylcholine, which counteracts pro-inflammatory cytokine (e.g., TNF- α) signaling and is called the 'cholinergic anti-inflammatory pathway' (CAP) (Burger et al., 2020; Kaniusas et al., 2019; Thayer, 2009). Increased *afferent* vagal activity can thus (indirectly via activating NTS neurons) impact the immune response by increasing *efferent* PNS influence (Tracey, 2009; Wohleb & Godbout, 2013).

The PNS also strongly interacts with the other component of the ANS, the SNS. During acute infection or psychosocial stress, SNS dominance is adaptive because it triggers a noradrenaline release at target tissues, which helps to mobilize energy resources by increasing heart rate and systolic and diastolic blood pressure (i.e., the fight-or-flight response; Epel et al., 2018). However, more chronic SNS activation, which has been found in COVID-19 patients (Porzionato et al., 2020), has also been related to increased cytokine production and the HPA axis overdrive described above (Tracey, 2009). In this regard, increased *efferent* vagal activity puts a 'break' on the continued activation of the SNS. The vagus nerve is, therefore, apart from immunomodulation, also implicated in autonomic control over cardiac activity (Kaniusas et al., 2019).

This is of particular importance given that by measuring the variability in the time interval between successive heartbeats—called, *heart rate variability* (HRV)—it is possible to infer the degree of vagal activity. Increased efferent vagal activity triggers neurons projecting to the sinoatrial node of the heart to release acetylcholine, thereby decreasing heart rate and increasing HRV (Burger et al., 2020). Particularly, the indexes root mean square of successive

differences (RMSSD) and high-frequency (HF) components can be used as indirect measures of (efferent) vagal function.

The ability to maintain high HRV in the occurrence of stressors has been suggested to be a flexible, adaptive physiological response to perceived stress (i.e., high vagal control). Furthermore, individuals with high resting-state HRV (i.e., high vagal tone) have been shown to recover faster in their acute immune, endocrine, and cardiovascular responses to stress (Kaniusas et al., 2019; Weber et al., 2010). In contrast, low HRV has been related to elevated inflammatory markers (Sloan et al., 2007), which can indicate a deficiency in the CAP. Moreover, individuals presenting with low (as compared to high) resting-state HRV have been found to experience more psychosocial stress, which might be exacerbated by different levels of social support (Lischke et al., 2018). Lastly, low HRV has been related to aberrant activity in a neural network that includes prefrontal and (para-)limbic brain regions that are involved in emotional and cognitive functioning, and that have been implicated in subjective, behavioral, and endocrine reactions to psychosocial stressors (Thayer et al., 2009, 2012). These bidirectional pathways between the brain and visceral organs via the ANS may also explain why sustained experiences of social isolation and exclusion negatively affect mental and physical health.

In sum, autonomic vagus nerve function plays a central role in governing mind-body interactions that have immune, endocrine, and neurocognitive and affective effects. Adaptive vagus nerve functioning is therefore considered crucial for the ability to inhibit ongoing physiological and psychosocial stress responses and is thus essential for enhancing physical and mental resilience.

5. Mental Health Interventions for Promoting Resilience

The above-described influence of both viral infections and psychosocial stressors on mind-body interactions provides one framework for understanding how dramatic increases in anxiety, mood, and other disorders are likely to occur during and in the aftermath of the COVID-19 pandemic. Although effective mental health interventions are available and have been shown to reduce health-damaging inflammation (e.g., Shields et al., 2020), cost-effectiveness models (based on epidemiological and socio-economic data) suggest that even in the unlikely event of optimal treatment being delivered, only 35%-50% of the overall burden of psychiatric disorders, such as anxiety disorders and depression, would be alleviated (Andrews et al., 2004). Therefore, there is an urgent need to develop prevention and treatment options that are safe, tolerable, affordable, and rapidly deployable on a large scale. Based on the above-described mechanistic picture, we propose that such interventions could benefit from targeting the vagus nerve in order to reduce risk for mental and physical disorders and enhance resilience to physiological and psychological stress. Consistent with this possibility, it was recently proposed that the balance between the PNS and the SNS might be crucial for understanding how individuals respond to the threat associated with the COVID-19 pandemic (Porges, 2020).

5.1. Transcutaneous Vagus Nerve Stimulation (tVNS)

As illustrated in Figure 2A, one promising, cost-effective approach to increase vagal activity is to apply non-invasive transcutaneous vagus nerve stimulation (tVNS), which has the potential to reduce SNS dominance, decrease pro-inflammatory markers, increase resiliency to and recovery from physiological and psychosocial stress, and reduce the risk of mental health complaints. There are two main ways to administer tVNS. The first is to superficially apply stimulation to the neck with a specialized device (e.g., GammaCore) to target the cervical branch of the vagus nerve. The second way is to apply stimulation at the cymba concha of the ear. This tissue is innervated by the exclusively afferent auricular branch of the vagus nerve

(ABVN), which fibers terminate in the NTS and subsequently project to the hypothalamus and forebrain regions (Pavlov & Tracey, 2012). fMRI studies suggest that tVNS can indeed activate these 'classic' vagal pathways as tVNS administration has been associated with altered activity in the NTS, hypothalamus, amygdala, hippocampus, ACC, insula, and nucleus accumbens (Frangos et al., 2015; Kong et al., 2018; Kraus et al., 2007; Yakunina et al., 2018; see also Burger & Verkuil, 2018).

-- Insert Figure 2 Here --

It has been suggested that tVNS induces anti-inflammatory effects directly by targeting the HPA axis (Bonaz et al., 2017), and indirectly by increasing activity in NTS neurons and activating the CAP (Bonaz et al., 2017; Thayer & Sternberg, 2010). In this regard, studies have shown a decrease in both salivary cortisol levels (Warren et al., 2019) and pro-inflammatory cytokine (e.g., TNF- α , IL-8, and possibly IL-6) levels after applying tVNS (Bremner et al., 2020; Brock et al., 2017), with long-lasting effects (Lerman et al., 2016). There is also some evidence that tVNS applied either after exposure to a stressor or without such exposure may improve the balance between the PNS and SNS (Brock et al., 2017; Gurel et al., 2020), and thereby increase HRV (Clancy et al., 2014; however, see also Borges et al., 2019).

Furthermore, based on preclinical invasive VNS studies, tVNS could potentially modulate serotonin and noradrenaline release in the hippocampus, amygdala, and mPFC, and increase neurogenesis and neuroplasticity in the hippocampus (Grimonprez et al., 2015). Lastly, recent studies have shown that multiple sessions of tVNS can be successfully applied as an intervention for disorders such as major depression, with improvements being seen in depression and anxiety symptoms, sleep, hopelessness, and cognitive function (Fang et al., 2016; Liu et al., 2016; Rong et al., 2016). In these studies, the reduction in depression severity was associated with changes in resting-state functional connectivity (rsFC) in brain networks implicated in physiological, cognitive, and emotional regulation. After four weeks of tVNS,

depressed patients who reported more substantial symptom improvements also exhibited greater decreases in rsFC between the default mode network (DMN, including the mPFC and ACC) and the anterior insula and parahippocampal gyrus (Fang et al., 2016); they also exhibited greater increases in rsFC between (a) the orbitofrontal cortex and the precuneus (Fang et al., 2016); (b) the nucleus accumbens (NAcc) and the rostral ACC/mPFC (Wang et al., 2018); and (c) the amygdala and the left dorsolateral prefrontal cortex (DLPFC; Liu et al., 2016).

In the COVID-19 context, lower cytokine (e.g., IL-6) levels have been associated with better treatment efficacy and remission of SARS-CoV-2 (Liu et al., 2020). The few ongoing pre-registered clinical trials that have investigated the efficacy of tVNS have focused on the physical health effects (e.g., respiratory function, cytokine storm) of cervically applied stimulation in inpatients during the acute infection phase (see Tornero et al., 2020). As of now, however, there are no pre-registered studies investigating the potential of tVNS to improve mental health in COVID-19 survivors. Given the mechanistic overview provided above, however, tVNS may be a potentially useful therapeutic intervention for COVID-19 survivors who have developed comorbid psychopathology, especially given that it is tolerable and can be applied safely with limited side-effects, even if used repeatedly (Redgrave et al., 2018). In addition, tVNS devices are relatively inexpensive, small and mobile, and easy to apply. In fact, under medical supervision, patients can be taught to self-administer the treatment at home (Fang et al., 2016; Rong et al., 2016).

5.2. Lifestyle Interventions to Increase Vagal Function

Apart from applying tVNS, activity in afferent vagal fibers can be increased through behavioral lifestyle interventions (see Figure 2B). Substantial research has shown that lifestyle interventions such as physical exercise, and mind-body interventions such as slow breathing (i.e., low respiration rate, long exhalations) and meditation, can increase vagal nerve functioning (Gerritsen & Band, 2018). During moderate-to-vigorous physical exercise (e.g., running, cycling,

rowing, swimming), for example, SNS activity and (nor)adrenaline production increases, elevating heart rate, blood pressure, and breathing rate above basal levels. Concurrently, vagal control drops. However, based on the continuous interaction between the PNS and the SNS branches of the ANS, vagal control (as measured using HRV) will rise again following the exercise (Stanley et al., 2013; Michael et al., 2017). In line with these effects, regular physical exercise of moderate intensity has been shown to increase vagal tone (i.e., during rest) and vagal control (i.e., with regard to a stressor; reviewed in Lujan & DiCarlo, 2013). In contrast, a sedentary lifestyle has been associated with PNS/SNS-imbalance, as characterized by reduced vagal nerve function (e.g., reduced HRV) (Thayer & Lane, 2009). In sum, therefore, physical exercise has been found to enhance vagal nerve functioning.

Consistent with the notion that vagal nerve functioning is a critical, health-relevant process, interventions based on physical exercise have also been shown to influence immune, endocrine, and CNS function (for a review, see Hendrikse et al., 2017). Systematic reviews and meta-analyses in healthy individuals have shown that even though physical exercise triggers an acute inflammatory response (e.g., due to muscle contractions and damage), it induces both short- and long-term anti-inflammatory effects. More specifically, exercise has been shown to reduce pro-inflammatory cytokine (e.g., IL-6, TNF- α) and CRP levels (Metsios et al., 2020), which may depend on the intensity of the training and individuals' age (Cronin et al., 2017; Souza et al., 2020). Interestingly, it has been suggested that daily exercise triggers an activation of the CAP, thereby protecting against inflammation (Lujan & DiCarlo, 2013). Research has also shown that healthy individuals who are under sustained stress exhibit improved mental health (i.e., fewer depressive symptoms) when they exercise at a moderate intensity for six weeks, as compared to individuals who do not exercise or who exercise at high intensity. Importantly, the beneficial effects of moderate exercise on mental health appear to be accompanied by decreases in levels of the pro-inflammatory cytokine TNF- α (Paolucci et al., 2018).

In addition to having anti-inflammatory effects, research has indicated that prolonged physical exercise can improve HPA axis-related endocrine function. Acute exercise triggers HPA axis activation and cortisol secretion (among other factors), depending on the intensity and the duration of the exercise. This hypercortisolism lasts for up to 2 hours following the exercise and is needed for energy mobilization but it also has beneficial, anti-inflammatory effects that help inhibit the exercise-induced inflammatory responses in muscles. Interestingly, however, individuals who are physically active, and thus repeatedly exposed to the associated physiological stress responses, appear to be better protected against the damaging effects of chronically elevated cortisol levels (Duclos & Tabarin, 2010). Relatedly, physically fit individuals have been found to exhibit a blunted cortisol response to laboratory-induced psychosocial stress as compared to moderately fit and untrained individuals (Rimmele et al., 2009).

Finally, the effects of physical exercise on CNS functioning, including neuroplasticity, cognitive, and affective functioning, are well known. Physical exercise has been shown to positively influence the connectivity of neural networks implicated in vagal and cognitive function (Mandolesi et al., 2018). Relatedly, exercise has been associated with improved prefrontal cortical and ACC functioning, which has been implicated in cognitive as well as emotional function (Kandola et al., 2019). Interestingly, a recent pilot study in which depressed patients completed a 12-week exercise program demonstrated improvements in cardiorespiratory fitness that were associated with increased ACC volume (Gujral et al., 2019). Lastly, physical exercise has also been associated with improved cognitive (Fernandes et al., 2017) and affective function (e.g., less persistent states of stress, anxiety, and depression; Harvey et al., 2018; von Haaren et al., 2015). In sum, physical exercise appears to strengthen resilience at least in part by facilitating recovery from stress on a physiological, immunological, endocrine, and psychological level, with improved vagal nerve function being a potential key mechanism of

action. Hence, the importance of physical exercise during the COVID-19 pandemic and beyond should not be underestimated.

Other lifestyle interventions such as yoga-based mind-body interventions that include posture, slow breathing, and meditation have also been shown to improve vagal nerve function (e.g., increase HRV) and thereby restore potential imbalances between the PNS and SNS (Streeter et al., 2012; Sullivan et al., 2018; Wells et al., 2016). These mind-body interventions have been associated with improved endocrine responses (i.e., decreases in cortisol levels; Streeter et al., 2012; Wells et al., 2016), improved immune function (i.e., decreases in pro-inflammatory markers; Black & Slavich, 2016; Gautam et al., 2020; Sullivan et al., 2018), increased telomerase activity (Black & Slavich, 2016), and adaptive changes in brain regions that are involved in cognitive and affective regulation (e.g., amygdala, prefrontal cortex, hippocampus; Kinser et al., 2012; Streeter et al., 2012). This may help explain why such interventions have a positive effect on emotional well-being and stress resiliency. Indeed, yoga-based mind-body interventions have been shown to reduce self-reported stress, anxiety, and depression levels in both clinical and non-clinical populations (Gautam et al., 2020; Gerbarg et al., 2015; Streeter et al., 2012; Wells et al., 2016).

Importantly, several of these lifestyle interventions are low cost, low risk, easily taught, rapidly effective, and sustainable first-line interventions that can have both preventive and therapeutic effects. Although lifestyle interventions have been studied in the context of resilience, the impact of vagal functioning has not been considered a central psychophysiological mechanism of change. Therefore, we believe that a more thorough understanding of how these and other lifestyle interventions promote emotional health acutely and over time – and how they influence mind-body interactions – will help advance the search for effective strategies for improving well-being and motivating healthy behaviors.

6. Open Questions and Future Directions

Looking forward, there are several avenues of investigation that could help advance our understanding of vagal function, health, and resilience. First, our review highlights a need for more information regarding the extent to which psychopathological complaints in COVID-19 survivors are related to aberrant inflammatory (e.g., IL-6, TNF- α), endocrine (e.g., cortisol), CNS (e.g., cortical thickness, functional connectivity), and ANS (e.g., vagal) function. Research is also needed to examine if these associations differ by infection severity (e.g., asymptomatic, mild, moderate, severe), or actual or perceived psychosocial stress levels. Along these lines, we also need to better understand the extent to which psychosocial stress plays a role in initial psychopathological symptom presentation in the general (non-infected) population, given that COVID-19-related restrictions increase the social isolation that all individuals experience. To accomplish this goal, researchers could conduct prospective, longitudinal studies in healthy individuals to evaluate if experiencing social isolation during the pandemic is related to harmful immune-neuroendocrine dynamics and, in addition, whether these changes are related to the development of cognitive or affective complaints over time.

Second, the potential of tVNS and lifestyle interventions to improve mind-body interactions (i.e., immune-neuroendocrine function, autonomic function, cognitive and affective processing) and mental health should be further evaluated. Particularly important in this context is the goal to identify whether such interventions reduce individuals' stress levels and, if so, what neural and biological mechanisms underlie these intervention-related changes in stress (Slavich, 2015). Given that the brain is the central organ responsible for perceiving stress and regulating peripheral biological responses to stressors, integrating neuroimaging into this research would be especially fruitful as it would help elucidate neural mechanisms that may underlie the efficacy of these interventions. For example, some prior research has identified

links between the brain and peripheral inflammation (e.g., Muscatell et al., 2015, 2016; Slavich et al., 2010b), but overall, this literature is small.

Furthermore, it will be important to evaluate whether individuals who present with distinct vagal function profiles at baseline (i.e., vagal endophenotypes) respond differently to interventions that aim to specifically target the vagus nerve. Consistent with the National Institute of Mental Health Research Domain Criteria (RDoC) approach (Cuthbert & Insel, 2013), we suggest that investigating vagal function as a dimensional human trait to predict response to a lifestyle intervention or tVNS would be desirable. Possibly, individuals who present with efficient vagal function might show ceiling effects in response to these interventions, as there might be less room for improvement. In contrast, those with more aberrant vagal function may benefit most from these interventions (see Brock et al., 2016), because they are experiencing a pre-interventional ANS imbalance characterized by higher SNS dominance and reduced PNS engagement. Altogether, future research is needed to understand how these interventions work and when they should be recommended for increasing resilience toward viral and psychosocial stressors, and, in addition, whether this is particularly true for individuals presenting with certain vagal endophenotypes.

Lastly, consistent with diathesis-stress models of psychopathology (e.g., Monroe & Simons, 1991), some people may be more prone to develop mental health problems following stressor exposure than others owing to neural, genetic, or other vulnerabilities (e.g., Quinn et al., 2020; Seiler et al., 2020; Slavich et al., 2014). For instance, individuals may exhibit prefrontal-mediated cognitive biases in response to psychosocial stressors, including the tendency to direct attention toward negative information, difficulty disengaging from it, and/or negative biases in interpreting social cues. Such cognitive biases may increase the risk of developing psychopathology such as mood and anxiety disorders (De Raedt & Koster, 2010). Of relevance here is whether biased neural processing of social exclusion is related to

heightened stress reactivity. Experiencing social exclusion has been related to increased activity in the insula, rostral ACC, and mPFC – particularly in individuals who are more vulnerable to develop psychopathology – thus reflecting increased reactivity toward social exclusion (for a review, see Dedoncker, 2020). Increased insula and ACC activity during social exclusion has also been related to TNF- α responses to social stress (Slavich et al., 2010a). Furthermore, the failure to engage the prefrontal cortex to (cognitively) control the stress response has also been associated with decreased HRV – and relatedly, SNS/PNS imbalance (Thayer et al., 2009; Smith et al., 2017). Individuals with the lowest vagal tone may therefore also benefit from prefrontally-targeted neuromodulatory interventions with techniques other than tVNS, such as transcranial direct current stimulation (tDCS) and transcranial magnetic stimulation (TMS; Remue et al., 2017), which particularly target these cognitive control processes to improve emotion regulation (for a review, see Dedoncker et al., 2020).

7. Conclusion

In conclusion, aberrant mind-body interactions can emerge from both being exposed to viral challenges, such as an infection, as well as in response to experiencing social stressors, such as social isolation or rejection. Risk for both exposure types has been greatly elevated during the COVID-19 pandemic, thus having profound implications for mental health and well-being on a worldwide level. On a positive note, several safe and tolerable mental health interventions already exist that can be implemented to reduce psychiatric risk and improve wellbeing. However, more research is needed to identify the specific types of interventions that are most affordable and effective for improving human health during these challenging times.

8. Funding Sources

This work was supported by a Society in Science—Branco Weiss Fellowship, Brain & Behavior Research Foundation NARSAD Young Investigator Grant [grant number 23958], and the

National Institutes of Health [grant number K08 MH103443] to GMS; the Fonds Wetenschappelijk Onderzoek, Flanders research projects 'Rode Neuzen' [grant number G0F4617N]; the Bijzonder Onderzoeksfonds [grant numbers BOF16/GOA/017, BOFSTA2017002501]; and the Sapienza University of Rome [grant number RM11715C7F74C683]. These funders had no role in planning or writing this article, or the decision to submit the article for publication.

9. Conflict of Interest

None

References

- Andrews, G., Issakidis, C., Sanderson, K., Corry, J., & Lapsley, H. (2004). Utilising survey data to inform public policy: Comparison of the cost-effectiveness of treatment of ten mental disorders. *The British Journal of Psychiatry*, 184(6), 526-533. doi:10.1192/bjp.184.6.526
- Benros, M. E., Nielsen, P. R., Nordentoft, M., Eaton, W. W., Dalton, S. O., & Mortensen, P. B. (2011). Autoimmune diseases and severe infections as risk factors for schizophrenia: a 30-year population-based register study. *American Journal of Psychiatry*, 168(12), 1303-1310. doi: 10.1176/appi.ajp.2011.11030516
- Benros, M. E., Waltoft, B. L., Nordentoft, M., Østergaard, S. D., Eaton, W. W., Krogh, J., & Mortensen, P. B. (2013). Autoimmune diseases and severe infections as risk factors for mood disorders: a nationwide study. *JAMA Psychiatry*, 70(8), 812-820. doi: 10.1001/jamapsychiatry.2013.1111
- Besedovsky, H. O., & Del Rey, A. (2011). Central and peripheral cytokines mediate immune-brain connectivity. *Neurochemical Research*, 36(1), 1-6. doi: 10.1007/s11064-010-0252-x
- Black, D. S., & Slavich, G. M. (2016). Mindfulness meditation and the immune system: A systematic review of randomized controlled trials. *Annals of the New York Academy of Sciences*, 1373, 13-24. doi: 10.1111/nyas.12998
- Bo, H. X., Li, W., Yang, Y., Wang, Y., Zhang, Q., Cheung, T., ... & Xiang, Y. T. (2020). Posttraumatic stress symptoms and attitude toward crisis mental health services among clinically stable patients with COVID-19 in China. *Psychological Medicine*, 1-2. doi: 10.1017/S0033291720000999
- Bonaz, B., Sinniger, V., & Pellissier, S. (2017). The vagus nerve in the neuro-immune axis: implications in the pathology of the gastrointestinal tract. *Frontiers in Immunology*, 8, 1452. doi : 10.3389/fimmu.2017.01452
- Borges, U., Laborde, S., & Raab, M. (2019). Influence of transcutaneous vagus nerve stimulation on cardiac vagal activity: Not different from sham stimulation and no effect of stimulation intensity. *PloS One*, 14(10), e0223848. doi: 10.1371/journal.pone.0223848
- Bremner, J. D., Gurel, N. Z., Jiao, Y., Wittbrodt, M. T., Levantsevych, O. M., Huang, M., ... & Pearce, B. D. (2020). Transcutaneous vagal nerve stimulation blocks stress-induced activation of interleukin-6 and interferon- γ in posttraumatic stress disorder: A double-

- blind, randomized, sham-controlled trial. *Brain, Behavior, & Immunity-Health*, 9, 100138. doi: 10.1016/j.bbih.2020.100138
- Brock, C., Brock, B., Aziz, Q., Møller, H. J., Pfeiffer Jensen, M., Drewes, A. M., & Farmer, A. D. (2017). Transcutaneous cervical vagal nerve stimulation modulates cardiac vagal tone and tumor necrosis factor-alpha. *Neurogastroenterology & Motility*, 29(5), e12999. doi: 10.1111/nmo.12999
- Brooks, S. K., Webster, R. K., Smith, L. E., Woodland, L., Wessely, S., Greenberg, N., & Rubin, G. J. (2020). The psychological impact of quarantine and how to reduce it: Rapid review of the evidence. *The Lancet*, 395(10227), 912-920. doi: 10.1016/S0140-6736(20)30460-8
- Brosschot, J. F., Verkuil, B., & Thayer, J. F. (2017). Exposed to events that never happen: Generalized unsafety, the default stress response, and prolonged autonomic activity. *Neuroscience & Biobehavioral Reviews*, 74, 287-296. doi: 10.1016/j.neubiorev.2016.07.019.
- Burger, A. M., D'Agostini, M., Verkuil, B., & Van Diest, I. (2020). Moving beyond belief: A narrative review of potential biomarkers for transcutaneous vagus nerve stimulation. *Psychophysiology*, 57(6), e13571. doi: 10.1111/psyp.13571
- Burger, A. M., & Verkuil, B. (2018). Transcutaneous nerve stimulation via the tragus: are we really stimulating the vagus nerve?. *Brain Stimulation: Basic, Translational, and Clinical Research in Neuromodulation*, 11(4), 945-946. doi: 10.1016/j.brs.2018.03.018
- Clancy, J. A., Mary, D. A., Witte, K. K., Greenwood, J. P., Deuchars, S. A., & Deuchars, J. (2014). Non-invasive vagus nerve stimulation in healthy humans reduces sympathetic nerve activity. *Brain Stimulation*, 7(6), 871-877. doi: 10.1016/j.brs.2014.07.031
- Coan, J. A. (2010). Adult attachment and the brain. *Journal of Social and Personal Relationships*, 27(2), 210-217. doi: 10.1177/0265407509360900
- Cronin, O., Keohane, D. M., Molloy, M. G., & Shanahan, F. (2017). The effect of exercise interventions on inflammatory biomarkers in healthy, physically inactive subjects: A systematic review. *QJM: An International Journal of Medicine*, 110(10), 629-637. doi: 10.1093/qjmed/hcx091
- Cuthbert, B. N., & Insel, T. R. (2013). Toward the future of psychiatric diagnosis: the seven pillars of RDoC. *BMC Medicine*, 11(1), 126. doi: 10.1186/1741-7015-11-126.
- De Raedt, R., & Koster, E. H. (2010). Understanding vulnerability for depression from a cognitive neuroscience perspective: A reappraisal of attentional factors and a new

- conceptual framework. *Cognitive, Affective, & Behavioral Neuroscience*, 10(1), 50-70. doi: 10.3758/CABN.10.1.50
- Dedoncker, J. (2020). A neuroscientific perspective on being criticized: Individual differences in neurobiological sensitivity to criticism and the potential for modifying it through neuromodulation (Doctoral dissertation, Ghent University). <http://hdl.handle.net/1854/LU-8668829>
- Dedoncker, J., Baeken, C., De Raedt, R., & Vanderhasselt, M. A. (2020). Combined transcranial direct current stimulation and psychological interventions: state of the art and promising perspectives for clinical psychology. *Biological Psychology*, 158, 107991. doi: 10.1016/j.biopsycho.2020.107991
- Desforges, M., Le Coupanec, A., Dubeau, P., Bourgouin, A., Lajoie, L., Dubé, M., & Talbot, P. J. (2020). Human coronaviruses and other respiratory viruses: underestimated opportunistic pathogens of the central nervous system?. *Viruses*, 12(1), 14. doi: 10.3390/v12010014.
- Dickerson, S. S., & Kemeny, M. E. (2004). Acute stressors and cortisol responses: a theoretical integration and synthesis of laboratory research. *Psychological Bulletin*, 130(3), 355-391. doi: 10.1037/0033-2909.130.3.355
- Duclos, M., & Tabarin, A. (2011). Exercise, training, and the hypothalamo–pituitary–adrenal axis. In *Hormone use and abuse by athletes* (pp. 9-15). Springer, Boston, MA. doi: 10.1007/978-1-4419-7014-5_2
- Epel, E. S., Crosswell, A. D., Mayer, S. E., Prather, A. A., Slavich, G. M., Puterman, E., & Mendes, W. B. (2018). More than a feeling: A unified view of stress measurement for population science. *Frontiers in Neuroendocrinology*, 49, 146-169. doi: 10.1016/j.yfrne.2018.03.001
- Fang, J., Rong, P., Hong, Y., Fan, Y., Liu, J., Wang, H., ... & Liu, R. (2016). Transcutaneous vagus nerve stimulation modulates default mode network in major depressive disorder. *Biological Psychiatry*, 79(4), 266-273. doi: 10.1016/j.biopsych.2015.03.025
- Felger, J. C., & Lotrich, F. E. (2013). Inflammatory cytokines in depression: Neurobiological mechanisms and therapeutic implications. *Neuroscience*, 246, 199-229. doi: 10.1016/j.neuroscience.2013.04.060
- Fernandes, J., Arida, R. M., & Gomez-Pinilla, F. (2017). Physical exercise as an epigenetic modulator of brain plasticity and cognition. *Neuroscience & Biobehavioral Reviews*, 80, 443-456. doi: 10.1016/j.neubiorev.2017.06.012.

- Frangos, E., Ellrich, J., & Komisaruk, B. R. (2015). Non-invasive access to the vagus nerve central projections via electrical stimulation of the external ear: fMRI evidence in humans. *Brain stimulation*, 8(3), 624-636. doi: 10.1016/j.brs.2014.11.018.
- Furman, D., Campisi, J., Verdin, E., Carrera-Bastos, P., Targ, S., Franceschi, C., ..., & Slavich, G. M. (2019). Chronic inflammation in the etiology of disease across the life span. *Nature Medicine*, 25, 1822-1832. doi: 10.1038/s41591-019-0675-0
- Gautam, S., Kumar, M., Kumar, U., & Dada, R. (2020). Effect of an 8-week yoga-based lifestyle intervention on psycho-neuro-immune axis, disease activity, and perceived quality of life in rheumatoid arthritis patients: A randomized controlled trial. *Frontiers in Psychology*, 11, 2259. doi: 10.3389/fpsyg.2020.02259
- Gerbarg, P. L., Jacob, V. E., Stevens, L., Bosworth, B. P., Chabouni, F., DeFilippis, E. M., ... & Scherl, E. J. (2015). The effect of breathing, movement, and meditation on psychological and physical symptoms and inflammatory biomarkers in inflammatory bowel disease: A randomized controlled trial. *Inflammatory Bowel Diseases*, 21(12), 2886-2896. doi: 10.1097/MIB.0000000000000568
- Gerritsen, R. J., & Band, G. P. (2018). Breath of life: The respiratory vagal stimulation model of contemplative activity. *Frontiers in Human Neuroscience*, 12, 397. doi: 10.3389/fnhum.2018.00397
- Grimonprez, A., Raedt, R., Baeken, C., Boon, P., & Vonck, K. (2015). The antidepressant mechanism of action of vagus nerve stimulation: Evidence from preclinical studies. *Neuroscience & Biobehavioral Reviews*, 56, 26-34. doi: 10.1016/j.neubiorev.2015.06.019
- Gujral, S., Aizenstein, H., Reynolds III, C. F., Butters, M. A., Grove, G., Karp, J. F., & Erickson, K. I. (2019). Exercise for depression: a feasibility trial exploring neural mechanisms. *The American Journal of Geriatric Psychiatry*, 27(6), 611-616. doi: 10.1016/j.jagp.2019.01.012.
- Gurel, N. Z., Huang, M., Wittbrodt, M. T., Jung, H., Ladd, S. L., Shandhi, M. M. H., ... & Inan, O. T. (2020). Quantifying acute physiological biomarkers of transcutaneous cervical vagal nerve stimulation in the context of psychological stress. *Brain Stimulation*, 13(1), 47-59. doi: 10.1016/j.brs.2019.08.002.
- Haroon, E., Raison, C. L., & Miller, A. H. (2012). Psychoneuroimmunology meets neuropsychopharmacology: Translational implications of the impact of inflammation on behavior. *Neuropsychopharmacology*, 37(1), 137-162. doi: 10.1038/npp.2011.205.

- Harvey, S. B., Øverland, S., Hatch, S. L., Wessely, S., Mykletun, A., & Hotopf, M. (2018). Exercise and the prevention of depression: Results of the HUNT cohort study. *American Journal of Psychiatry*, 175(1), 28-36. doi: 10.1176/appi.ajp.2017.16111223.
- Hendrikse, J., Kandola, A., Coxon, J., Rogasch, N., & Yücel, M. (2017). Combining aerobic exercise and repetitive transcranial magnetic stimulation to improve brain function in health and disease. *Neuroscience & Biobehavioral Reviews*, 83, 11-20. doi: 10.1016/j.neubiorev.2017.09.023
- Kandola, A., Ashdown-Franks, G., Hendrikse, J., Sabiston, C. M., & Stubbs, B. (2019). Physical activity and depression: Towards understanding the antidepressant mechanisms of physical activity. *Neuroscience & Biobehavioral Reviews*, 107, 525-539. doi: 10.1016/j.neubiorev.2019.09.040
- Kaniusas, E., Kampusch, S., Tittgemeyer, M., Panetsos, F., Gines, R. F., Papa, M., ... & Samoudi, A. M. (2019). Current directions in the auricular vagus nerve stimulation I-a physiological perspective. *Frontiers in Neuroscience*, 13, 854. doi: 10.3389/fnins.2019.00854.
- Khandaker, G. M., Pearson, R. M., Zammit, S., Lewis, G., & Jones, P. B. (2014). Association of serum interleukin 6 and C-reactive protein in childhood with depression and psychosis in young adult life: A population-based longitudinal study. *JAMA Psychiatry*, 71(10), 1121-1128. doi: 10.1001/jamapsychiatry.2014.1332.
- Kinser, P. A., Goehler, L., & Taylor, A. G. (2012). How might yoga help depression? A neurobiological perspective. *Explore (NY)*, 8(2), 118-126. doi: 10.1016/j.explore.2011.12.005.
- Kong, J., Fang, J., Park, J., Li, S., & Rong, P. (2018). Treating depression with transcutaneous auricular vagus nerve stimulation: State of the art and future perspectives. *Frontiers in Psychiatry*, 9, 20. doi: 10.3389/fpsy.2018.00020.
- Kong, X., Zheng, K., Tang, M., Kong, F., Zhou, J., Diao, L., ... & Dong, Y. (2020). Prevalence and factors associated with depression and anxiety of hospitalized patients with COVID-19. *MedRxiv*. doi: 10.1101/2020.03.24.20043075
- Kraus, T., Hösl, K., Kiess, O., Schanze, A., Kornhuber, J., & Forster, C. (2007). BOLD fMRI deactivation of limbic and temporal brain structures and mood enhancing effect by transcutaneous vagus nerve stimulation. *Journal of Neural Transmission*, 114(11), 1485-1493. doi: 10.1007/s00702-007-0755-z.
- Lam, M. H. B., Wing, Y. K., Yu, M. W. M., Leung, C. M., Ma, R. C., Kong, A. P., ... & Lam, S. P. (2009). Mental morbidities and chronic fatigue in severe acute respiratory syndrome

- survivors: long-term follow-up. *Archives of Internal Medicine*, 169(22), 2142-2147. doi: 10.1001/archinternmed.2009.384.
- Lerman, I., Hauger, R., Sorkin, L., Proudfoot, J., Davis, B., Huang, A., ... & Baker, D. G. (2016). Noninvasive transcutaneous vagus nerve stimulation decreases whole blood culture-derived cytokines and chemokines: A randomized, blinded, healthy control pilot trial. *Neuromodulation: Technology at the Neural Interface*, 19(3), 283-290. doi: 10.1111/ner.12398.
- Lischke, A., Jacksteit, R., Mau-Moeller, A., Pahnke, R., Hamm, A. O., & Weippert, M. (2018). Heart rate variability is associated with psychosocial stress in distinct social domains. *Journal of psychosomatic research*, 106, 56-61. doi: 10.1016/j.jpsychores.2018.01.005
- Liu, J., Fang, J., Wang, Z., Rong, P., Hong, Y., Fan, Y., ... & Kong, J. (2016). Transcutaneous vagus nerve stimulation modulates amygdala functional connectivity in patients with depression. *Journal of Affective Disorders*, 205, 319-326. doi: 10.1016/j.jad.2016.08.003.
- Liu, T., Zhang, J., Yang, Y., Ma, H., Li, Z., Zhang, J., ... & Zhang, L. (2020). The role of interleukin-6 in monitoring severe case of coronavirus disease 2019. *EMBO Molecular Medicine*, 12(7), e12421. doi: 10.15252/emmm.202012421
- Lujan, H. L., & DiCarlo, S. E. (2013). Physical activity, by enhancing parasympathetic tone and activating the cholinergic anti-inflammatory pathway, is a therapeutic strategy to restrain chronic inflammation and prevent many chronic diseases. *Medical Hypotheses*, 80(5), 548-552. doi: 10.1016/j.mehy.2013.01.014.
- Mandolesi, L., Polverino, A., Montuori, S., Foti, F., Ferraioli, G., Sorrentino, P., & Sorrentino, G. (2018). Effects of physical exercise on cognitive functioning and wellbeing: biological and psychological benefits. *Frontiers in Psychology*, 9, 509. doi: 10.3389/fpsyg.2018.00509.
- Masten, C. L., Eisenberger, N. I., Borofsky, L. A., McNealy, K., Pfeifer, J. H., & Dapretto, M. (2011). Subgenual anterior cingulate responses to peer rejection: A marker of adolescents' risk for depression. *Development and Psychopathology*, 23(1), 283-292. doi:10.1017/S0954579410000799
- Mental Health Foundation. Loneliness during Corona-virus. 2020. Available at: <https://www.mentalhealth.org.uk/coronavirus/loneliness-during-coronavirus>. Accessed January 24, 2021.
- Metsios, G. S., Moe, R. H., & Kitas, G. D. (2020). Exercise and inflammation. *Best Practice & Research Clinical Rheumatology*, 34(2), 101504. doi: 10.1016/j.berh.2020.101504

- Michael, S., Graham, K. S., & Davis, G. M. (2017). Cardiac autonomic responses during exercise and post-exercise recovery using heart rate variability and systolic time intervals—a review. *Frontiers in Physiology*, 8, 301. doi: 10.3389/fphys.2017.00301.
- Michopoulos, V., Powers, A., Gillespie, C. F., Ressler, K. J., & Jovanovic, T. (2017). Inflammation in fear-and anxiety-based disorders: PTSD, GAD, and beyond. *Neuropsychopharmacology*, 42(1), 254-270. doi: 10.1038/npp.2016.146
- Miller, A. H., Haroon, E., Raison, C. L., & Felger, J. C. (2013). Cytokine targets in the brain: Impact on neurotransmitters and neurocircuits. *Depression and Anxiety*, 30(4), 297-306. doi: 10.1002/da.22084
- Miller, E. D. (2020). Loneliness in the era of COVID-19. *Frontiers in Psychology*, 11. doi: 10.3389/fpsyg.2020.02219
- Monroe, S. M., & Simons, A. D. (1991). Diathesis-stress theories in the context of life stress research: Implications for the depressive disorders. *Psychological Bulletin*, 110(3), 406-425. doi: 10.1037/0033-2909.110.3.406
- Monroe, S. M., Slavich, G. M., & Georgiades, K. (2014). The social environment and depression: The roles of life stress. In I. H. Gotlib & C. L. Hammen (Eds.), *Handbook of depression*, third edition (pp. 296-314). New York: The Guilford Press.
- Muscatell, K. A., Dedovic, K., Slavich, G. M., Jarcho, M. R., Breen, E. C., Bower, J. E., Irwin, M. R., & Eisenberger, N. I. (2015). Greater amygdala activity and dorsomedial prefrontal-amygdala coupling are associated with enhanced inflammatory responses to stress. *Brain, Behavior, and Immunity*, 43, 46-53. doi: 10.1016/j.bbi.2014.06.201
- Muscatell, K. A., Dedovic K., Slavich, G. M., Jarcho M. R., Breen, E. C., Bower, J. E., Irwin, M. R., & Eisenberger, N. I. (2016). Neural mechanisms linking social status and inflammatory responses to social stress. *Social Cognitive and Affective Neuroscience*, 11, 915-922. doi: 10.1093/scan/nsw025
- Nelson, B. W., Pettitt, A., Flannery, J. E., & Allen, N. B. (2020). Rapid assessment of psychological and epidemiological correlates of COVID-19 concern, financial strain, and health-related behavior change in a large online sample. *PloS One*, 15(11), e0241990. doi: 10.1371/journal.pone.0241990
- Nicola, M., Alsafi, Z., Sohrabi, C., Kerwan, A., Al-Jabir, A., Iosifidis, C., ... & Agha, R. (2020). The socio-economic implications of the coronavirus and COVID-19 pandemic: a review. *International Journal of Surgery*, 78, 185-193. doi: 10.1016/j.ijsu.2020.04.018

- Nusslock, R., & Miller, G. E. (2016). Early-life adversity and physical and emotional health across the lifespan: A neuroimmune network hypothesis. *Biological Psychiatry*, 80(1), 23-32. doi: 10.1016/j.biopsych.2015.05.017
- Pace, T. W., & Miller, A. H. (2009). Cytokines and glucocorticoid receptor signaling: relevance to major depression. *Annals of the New York Academy of Sciences*, 1179, 86-105. doi: 10.1111/j.1749-6632.2009.04984.x
- Pancani, L., Marinucci, M., Aureli, N., & Riva, P. (2020). Forced social isolation and mental health: A study on 1006 Italians under COVID-19 quarantine.
- Paolucci, E. M., Loukov, D., Bowdish, D. M. E., & Heisz, J. J. (2018). Exercise reduces depression and inflammation but intensity matters. *Biological Psychology*, 133, 79-84. doi: 10.1016/j.biopsycho.2018.01.015
- Passos, I. C., Vasconcelos-Moreno, M. P., Costa, L. G., Kunz, M., Brietzke, E., Quevedo, J., ... & Kauer-Sant'Anna, M. (2015). Inflammatory markers in post-traumatic stress disorder: A systematic review, meta-analysis, and meta-regression. *The Lancet Psychiatry*, 2(11), 1002-1012. doi: 10.1016/S2215-0366(15)00309-0
- Pavlov, V. A., & Tracey, K. J. (2012). The vagus nerve and the inflammatory reflex—linking immunity and metabolism. *Nature Reviews Endocrinology*, 8(12), 743. doi: 10.1038/nrendo.2012.189
- Porges, S. W. (2020). The COVID-19 Pandemic is a paradoxical challenge to our nervous system: A Polyvagal Perspective. *Clinical Neuropsychiatry*, 17, 135-8. doi: 10.36131/CN20200220.
- Porzionato, A., Emmi, A., Barbon, S., Boscolo-Berto, R., Stecco, C., Stocco, E., ... & De Caro, R. (2020). Sympathetic activation: a potential link between comorbidities and COVID-19. *The FEBS Journal*, 287(17), 3681-3688. doi: 10.1111/febs.15481.
- Quinn, M. E., Stanton, C. H., Slavich, G. M., & Joormann, J. (2020). Executive control, cytokine reactivity to social stress, and depressive symptoms: Testing the Social Signal Transduction Theory of Depression. *Stress*, 23, 60-68. doi: 10.1080/10253890.2019.1641079
- Raony, Í., de Figueiredo, C. S., Pandolfo, P., Giestal-de-Araujo, E., Oliveira-Silva Bomfim, P., & Savino, W. (2020). Psycho-neuroendocrine-immune interactions in COVID-19: Potential impacts on mental health. *Frontiers in Immunology*, 11, 1170. Doi: 10.3389/fimmu.2020.01170

- Redgrave, J., Day, D., Leung, H., Laud, P. J., Ali, A., Lindert, R., & Majid, A. (2018). Safety and tolerability of Transcutaneous Vagus Nerve stimulation in humans; a systematic review. *Brain Stimulation*, 11(6), 1225-1238. doi: 10.1016/j.brs.2018.08.010
- Remue, J., Vanderhasselt, M. A., Baeken, C., Rossi, V., Tullo, J., & De Raedt, R. (2016). The effect of a single HF-rTMS session over the left DLPFC on the physiological stress response as measured by heart rate variability. *Neuropsychology*, 30(6), 756-766. doi: 10.1037/neu0000255
- Rimmele, U., Seiler, R., Marti, B., Wirtz, P. H., Ehlert, U., & Heinrichs, M. (2009). The level of physical activity affects adrenal and cardiovascular reactivity to psychosocial stress. *Psychoneuroendocrinology*, 34(2), 190-198. doi: 10.1016/j.psyneuen.2008.08.023
- Rong, P., Liu, J., Wang, L., Liu, R., Fang, J., Zhao, J., ... & Ben, H. (2016). Effect of transcutaneous auricular vagus nerve stimulation on major depressive disorder: A nonrandomized controlled pilot study. *Journal of Affective Disorders*, 195, 172-179. doi: 10.1016/j.jad.2016.02.031
- Rosenblat, J. D., Cha, D. S., Mansur, R. B., & McIntyre, R. S. (2014). Inflamed moods: A review of the interactions between inflammation and mood disorders. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 53, 23-34. doi: 10.1016/j.pnpbp.2014.01.013
- Schiepers, O. J., Wichers, M. C., & Maes, M. (2005). Cytokines and major depression. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 29(2), 201-217. doi: 10.1016/j.pnpbp.2004.11.003.
- Seiler, A., von Känel, R., & Slavich, G. M. (2020). The psychobiology of bereavement and health: A conceptual review from the perspective of Social Signal Transduction Theory of Depression. *Frontiers in Psychiatry*, 11, 565239. doi: 10.3389/fpsy.2020.565239
- Serra, M., Pisu, M. G., Floris, I., & Biggio, G. (2005). Social isolation-induced changes in the hypothalamic–pituitary–adrenal axis in the rat. *Stress*, 8(4), 259-264. doi: 10.1080/10253890500495244
- Shields, G. S., Spahr, C. M., & Slavich, G. M. (2020). Psychosocial interventions and immune system function: A systematic review and meta-analysis of randomized clinical trials. *JAMA Psychiatry*, 77, 1031-1043. doi: 10.1001/jamapsychiatry.2020.0431
- Slavich, G. M. (2015). Understanding inflammation, its regulation, and relevance for health: A top scientific and public priority. *Brain, Behavior, and Immunity*, 45, 13-14. doi: 10.1016/j.bbi.2014.10.012

- Slavich, G. M. (2016). Life stress and health: A review of conceptual issues and recent findings. *Teaching of Psychology*, 43(4), 346-355. doi: 10.1177/0098628316662768
- Slavich, G. M. (2020a). Social safety theory: A biologically based evolutionary perspective on life stress, health, and behavior. *Annual Review of Clinical Psychology*, 16, 265-295. doi: 10.1146/annurev-clinpsy-032816-045159
- Slavich, G. M. (2020b). Psychoneuroimmunology of stress and mental health. In K. L. Harkness & E. P. Hayden (Eds.), *The Oxford handbook of stress and mental health* (pp. 519-546). New York: Oxford University Press. doi: 10.1093/oxfordhb/9780190681777.013.24
- Slavich, G. M., & Irwin, M. R. (2014). From stress to inflammation and major depressive disorder: A social signal transduction theory of depression. *Psychological Bulletin*, 140, 774-815. doi: 10.1037/a0035302
- Slavich, G. M., O'Donovan, A., Epel, E. S., & Kemeny, M. E. (2010a). Black sheep get the blues: A psychobiological model of social rejection and depression. *Neuroscience & Biobehavioral Reviews*, 35(1), 39-45. doi: 10.1016/j.neubiorev.2010.01.003
- Slavich, G. M., Tartter, M. A., Brennan, P. A., & Hammen, C. (2014). Endogenous opioid system influences depressive reactions to socially painful targeted rejection life events. *Psychoneuroendocrinology*, 49, 141-149. doi: 10.1016/j.psyneuen.2014.07.009
- Slavich, G. M., Thornton, T., Torres, L. D., Monroe, S. M., & Gotlib, I. H. (2009). Targeted rejection predicts hastened onset of major depression. *Journal of Social and Clinical Psychology*, 28, 223-243. doi: 10.1521/jscp.2009.28.2.223
- Slavich, G. M., Way, B. M., Eisenberger, N. I., & Taylor, S. E. (2010b). Neural sensitivity to social rejection is associated with inflammatory responses to social stress. *Proceedings of the National Academy of Sciences of the United States of America*, 107(33), 14817-14822. doi: 10.1073/pnas.1009164107
- Sloan, R. P., McCreath, H., Tracey, K. J., Sidney, S., Liu, K., & Seeman, T. (2007). RR interval variability is inversely related to inflammatory markers: The CARDIA study. *Molecular Medicine*, 13(3-4), 178-184. doi: 10.2119/2006-00112.Sloan
- Smith, K. (2020). The association between loneliness, social isolation and inflammation: A systematic review and meta-analysis. *Neuroscience & Biobehavioral Reviews*, 112, 519-541. doi: 10.1016/j.neubiorev.2020.02.002
- Smith, R., Thayer, J. F., Khalsa, S. S., & Lane, R. D. (2017). The hierarchical basis of neurovisceral integration. *Neuroscience & Biobehavioral Reviews*, 75, 274-296. doi:10.1016/j.neubiorev.2017.02.003

- Souza, D., Coswig, V., de Lira, C. A. B., & Gentil, P. (2020). H"IT"ting the Barriers for Exercising during Social Isolation. *Biology*, 9(9), 245. doi: 10.3390/biology9090245
- Stanley, J., Peake, J. M., and Buchheit, M. (2013). Cardiac parasympathetic reactivation following exercise: Implications for training prescription. *Sports Medicine*, 43, 1259-1277. doi: 10.1007/s40279-013-0083-4
- Stapelberg, N. J. C., Neumann, D. L., Shum, D., & Headrick, J. P. (2019). Health, pre-disease and critical transition to disease in the psycho-immune-neuroendocrine network: Are there distinct states in the progression from health to major depressive disorder?. *Physiology & Behavior*, 198, 108-119. doi: 10.1016/j.physbeh.2018.10.014.
- Stephens, A., Hamer, M., & Chida, Y. (2007). The effects of acute psychological stress on circulating inflammatory factors in humans: A review and meta-analysis. *Brain, Behavior, and Immunity*, 21(7), 901-912. doi: 10.1016/j.bbi.2007.03.011
- Streeter, C. C., Gerbarg, P. L., Saper, R. B., Ciraulo, D. A., & Brown, R. P. (2012). Effects of yoga on the autonomic nervous system, gamma-aminobutyric-acid, and allostasis in epilepsy, depression, and post-traumatic stress disorder. *Medical Hypotheses*, 78(5), 571-579. doi: 10.1016/j.mehy.2012.01.021
- Sullivan, M. B., Erb, M., Schmalzl, L., Moonaz, S., Noggle Taylor, J., & Porges, S. W. (2018). Yoga therapy and polyvagal theory: The convergence of traditional wisdom and contemporary neuroscience for self-regulation and resilience. *Frontiers in Human Neuroscience*, 12, 67. doi: 10.3389/fnhum.2018.00067
- Taquet, M., Luciano, S., Geddes, J. R., & Harrison, P. J. (2020). Bidirectional associations between COVID-19 and psychiatric disorder: Retrospective cohort studies of 62 354 COVID-19 cases in the USA. *The Lancet Psychiatry*, 8(2), 130-140. doi: 10.1016/s2215-0366(20)30462-4
- Thayer, J. F. (2009). Vagal tone and the inflammatory reflex. *Cleveland Clinic Journal of Medicine*, 76(Suppl 2), S23-26. doi: 10.3949/ccjm.76.s2.05.
- Thayer, J. F., Åhs, F., Fredrikson, M., Sollers III, J. J., & Wager, T. D. (2012). A meta-analysis of heart rate variability and neuroimaging studies: implications for heart rate variability as a marker of stress and health. *Neuroscience & Biobehavioral Reviews*, 36(2), 747-756. doi: 10.1016/j.neubiorev.2011.11.009
- Thayer, J. F., & Lane, R. D. (2009). Claude Bernard and the heart-brain connection: Further elaboration of a model of neurovisceral integration. *Neuroscience & Biobehavioral Reviews*, 33(2), 81-88. doi: 10.1016/j.neubiorev.2008.08.004

- Thayer, J. F., & Sternberg, E. M. (2010). Neural aspects of immunomodulation: focus on the vagus nerve. *Brain, Behavior, and Immunity*, 24(8), 1223-1228. doi: 10.1016/j.bbi.2010.07.247.
- Thayer, J. F., Hansen, A. L., Saus-Rose, E., & Johnsen, B. H. (2009). Heart rate variability, prefrontal neural function, and cognitive performance: The neurovisceral integration perspective on self-regulation, adaptation, and health. *Annals of Behavioral Medicine*, 37(2), 141-153. doi: 10.1007/s12160-009-9101-z
- Tornero, C., Vallejo, R., Cedeño, D., Orduña, J., Pastor, E., Belaouchi, M., ... & del Mar Garzando, M. (2020). A prospective, randomized, controlled study assessing vagus nerve stimulation using the gammaCore®-Sapphire device for patients with moderate to severe CoViD-19 Respiratory Symptoms (SAVIOR): A structured summary of a study protocol for a randomised controlled trial". *Trials*, 21(1), 576. doi: 10.1186/s13063-020-04486-w
- Tracey, K. J. (2009). Reflex control of immunity. *Nature Reviews Immunology*, 9(6), 418-428. doi: 10.1038/nri2566
- Tursich, M., Neufeld, R. W. J., Frewen, P. A., Harricharan, S., Kibler, J. L., Rhind, S. G., & Lanius, R. A. (2014). Association of trauma exposure with proinflammatory activity: A transdiagnostic meta-analysis. *Translational Psychiatry*, 4(7), e413-e413. doi: 10.1038/tp.2014.56
- Van Bavel, J. J., Baicker, K., Boggio, P. S., Capraro, V., Cichocka, A., Cikara, M., ... & Willer, R. (2020). Using social and behavioural science to support COVID-19 pandemic response. *Nature Human Behaviour*, 4(5), 460-471. doi: 10.1038/s41562-020-0884-z.
- Varatharaj, A., Thomas, N., Ellul, M. A., Davies, N. W., Pollak, T. A., Tenorio, E. L., ... & Coles, J. P. (2020). Neurological and neuropsychiatric complications of COVID-19 in 153 patients: A UK-wide surveillance study. *The Lancet Psychiatry*, 7(10), 875-882. doi: 10.1016/S2215-0366(20)30287-X
- von Haaren, B., Haertel, S., Stumpp, J., Hey, S., & Ebner-Priemer, U. (2015). Reduced emotional stress reactivity to a real-life academic examination stressor in students participating in a 20-week aerobic exercise training: A randomised controlled trial using Ambulatory Assessment. *Psychology of Sport and Exercise*, 20, 67-75. doi: /10.1016/j.psychsport.2015.04.004.
- Warren, C. M., Tona, K. D., Ouwerkerk, L., Van Paridon, J., Poletiek, F., van Steenbergen, H., ... & Nieuwenhuis, S. (2019). The neuromodulatory and hormonal effects of transcutaneous vagus nerve stimulation as evidenced by salivary alpha amylase,

- salivary cortisol, pupil diameter, and the P3 event-related potential. *Brain Stimulation*, 12(3), 635-642. doi: 10.1016/j.brs.2018.12.224.
- Wang, Z., Fang, J., Liu, J., Rong, P., Jorgenson, K., Park, J., ... & Kong, J. (2018). Frequency-dependent functional connectivity of the nucleus accumbens during continuous transcutaneous vagus nerve stimulation in major depressive disorder. *Journal of Psychiatric Research*, 102, 123-131. doi: 10.1016/j.jpsychires.2017.12.018.
- Weber, C. S., Thayer, J. F., Rudat, M., Wirtz, P. H., Zimmermann-Viehoff, F., Thomas, A., ... & Deter, H. C. (2010). Low vagal tone is associated with impaired post stress recovery of cardiovascular, endocrine, and immune markers. *European Journal of Applied Physiology*, 109(2), 201-211. doi: 10.1007/s00421-009-1341-x
- Wells, S. Y., Lang, A. J., Schmalzl, L., Groessl, E. J., & Strauss, J. L. (2016). Yoga as an intervention for PTSD: A theoretical rationale and review of the literature. *Current Treatment Options in Psychiatry*, 3(1), 60-72. doi: 10.1007/s40501-016-0068-7
- Wohleb, E. S., & Godbout, J. P. (2013). Basic aspects of the immunology of neuroinflammation. In *Inflammation in psychiatry* (Vol. 28, pp. 1-19). Karger Publishers. doi: 10.1159/000343964.
- Wohleb, E. S., McKim, D. B., Sheridan, J. F., & Godbout, J. P. (2015). Monocyte trafficking to the brain with stress and inflammation: A novel axis of immune-to-brain communication that influences mood and behavior. *Frontiers in Neuroscience*, 8, 447. doi: 10.3389/fnins.2014.00447
- Yakunina, N., Kim, S. S., & Nam, E. C. (2017). Optimization of transcutaneous vagus nerve stimulation using functional MRI. *Neuromodulation: Technology at the Neural Interface*, 20(3), 290-300. doi: 10.1111/ner.12541

Figure Captions

Fig 1. Pathways by which exposure to SARS-CoV-2 and psychosocial stressors can negatively impact mental health. (A) Exposure to SARS-CoV-2 can upregulate inflammatory activity, characterized by increased pro-inflammatory cytokine (e.g., IL-6, TNF- α) production. Via the afferent vagus nerve fibers, information regarding the presence of cytokines is conveyed to the NTS and subsequently to the hypothalamus where it can activate the HPA axis. When

this occurs, CRH released from the hypothalamus triggers ACTH release from the pituitary, and increased cortisol secretion from the adrenal cortex, leading to a generalized stress response. When transient, this response is beneficial, as cortisol generally reduces pro-inflammatory cytokine signaling in immune cells. However, when cytokine levels are frequently or persistently elevated, the HPA axis can become hyperactivated and glucocorticoid receptors on immune cells can become insensitive to the immunosuppressive effects of cortisol, leading to cortisol build-up and sustained inflammation. This chronic inflammatory state can affect neurotransmitter (e.g., serotonin) metabolism and negatively impact mental health. Afferent vagal projections are also propagated to other brain regions involved in the pathophysiology of mental health problems, including the amygdala, ACC, and mPFC. **(B)** Psychosocial stressors, such as social isolation and rejection, can also upregulate inflammatory activity and affect mental health. These stressors are processed by the brain and can activate the HPA axis, leading to increased cortisol release. Psychosocial stressors can also trigger strong cardiovascular reactivity and increased noradrenaline signaling, indicative of a shift towards SNS dominance over PNS activation. Third, these stressors have been associated with glucocorticoid insensitivity, meaning they can trigger immune cells to become resistant to the immunosuppressive effects of cortisol, which can in turn lead to a chronic inflammatory state. **(C)** Finally, it is important to note that the vagal PNS and the SNS are strongly interactive. Efferent vagal neurons project back from the NTS to cells in bodily tissues (e.g., heart, lungs, gastrointestinal system) where they release acetylcholine. Efferent vagal fibers can counteract a sustained activation of the SNS (e.g., by slowing heart rate and blood pressure). They can also reduce pro-inflammatory cytokine production through the CAP. Abbreviations: ACC, anterior cingulate cortex; ACTH, adrenocorticotrophic hormone; A, amygdala; CAP, cholinergic anti-inflammatory pathway; CRH, corticotropin-releasing hormone; HPA axis, hypothalamic-pituitary-adrenal axis; IL-6, interleukin 6; mPFC, medial prefrontal cortex; NTS, nucleus solitary tract; PNS, parasympathetic nervous system; SNS, sympathetic nervous system; TNF- α , tumor necrosis factor- α . Created with BioRender.com.

Fig 2. Pathways by which vagal function-enhancing interventions can normalize biological functioning and improve mental health. (A) tVNS applied to the afferent cervical

(neck) or auricular branch of the vagus nerve (cymba concha of the ear) modulates activity in the NTS, hypothalamus, hippocampus, amygdala, ACC, and other brain regions. tVNS may have anti-inflammatory effects by targeting the HPA axis (i.e., dashed arrow) and by activating the CAP via enhanced efferent vagal activity. Preclinical invasive VNS studies suggest that tVNS may also improve the PNS/SNS balance, thereby reducing SNS dominance and modulating noradrenaline release. In depressed patients, tVNS has been shown to improve symptomatology and functioning in neural networks and brain regions involved in cognitive and affective regulation, including the ACC, insula, NAc, mPFC, and DLPFC. **(B)** Regular physical exercise can also enhance vagal function and may improve the PNS/SNS balance. Habitual exercise can reduce pro-inflammatory activity (e.g., by activating the CAP) and improve HPA axis-related endocrine function (e.g., reduced cortisol reactivity to psychosocial stressors). Lastly, exercise improves cognitive and affective regulation, as reflected in part by improved neural functioning in relevant brain regions and networks, including the ACC and PFC.

Abbreviations: ACC, anterior cingulate cortex; A, amygdala; CAP, cholinergic anti-inflammatory pathway; DLPFC, dorsolateral prefrontal cortex; H, hippocampus; HPA axis, hypothalamic-pituitary-adrenal axis; HT, hypothalamus; mPFC, medial prefrontal cortex; NAc, nucleus accumbens; NTS, nucleus solitary tract; PFC, prefrontal cortex; PNS, parasympathetic nervous system; SNS, sympathetic nervous system; tVNS, transcutaneous vagus nerve stimulation.

Created with BioRender.com.